

**TRICEPTIN- lidocaine / menthol / methyl salicylate patch**  
**SOLA Pharmaceuticals**

-----  
**Triceptin Patch**

***Active ingredient (in each packet)***

Lidocaine 4%

***Purpose***

Topical anesthetic

***Active ingredient (in each packet)***

Menthol 1%

***Purpose***

Topical analgesic

***Active ingredient (in each packet)***

Methyl salicylate 2%

***Purpose***

Topical analgesic

***Use***

Temporarily relieves mild to moderate aches and pains of muscles and joints associated with:

- Muscle soreness
- strains
- sprains
- arthritis
- simple backache
- muscle stiffness
- bruises

***Warnings***

**For external use only**

**Stomach bleeding warning:** This product contains a NSAID, which may cause stomach bleeding. The chance is small, but higher if you

- are aged 60 or older
- have had stomach ulcers or bleeding problems
- take other drugs containing an NSAID (Aspirin, ibuprofen, Naproxen, or others)
- take a blood thinning (anticoagulant) or steroidal drug
- have three or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

***Do not use***

- More than one patch on your body at a time
- On cut, irritated or swollen skin
- On puncture wounds
- For more than one week without consulting a doctor
- If you are allergic to any active or inactive ingredients
- If pouch is damaged or opened

***When using this product***

- Use only as directed
- Read and follow all directions and warnings on this carton
- Do not allow contact with the eyes
- Do not use at the same time as other topical analgesics
- Do not bandage tightly or apply local heat (such as heating pads) to the area of use
- Do no microwave
- Dispose of used patch in manner that always keeps product away from children and pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests this patch.

***Stop use and consult a doctor if***

- Condition worsens
- Redness is present
- Irritation develops
- Symptoms persist for more than 7 days or clear up and occur again within a few days
- You experience signs of skin injury, such as pain, swelling, or blistering where the product was applied
- Stomach pain or upset gets worst or lasts
- You feel faint, vomit blood, or have bloody or black stools (these are signs of stomach bleeding).

***If pregnant or breastfeeding***

ask a healthcare professional before use.

***Keep out of reach of children and pets.***

If swallowed, get medical help or contact a Poison Control Center right away.

### ***Directions***

Adults, or children over the age of 12: Clean and dry affected area. Carefully remove backing from patch starting at a corner. Apply sticky side of patch to affected area and leave in place for 8 to 12 hours. If pain lasts after using the first patch, a second patch may be applied for up to another 8 to 12 hours. Only use one patch at a time. do not use more than 2 patches per day. Wash hands with soap and water after applying or removing patch. reseal pouch containing unused patches immediately after each use. Children under 12 years of age: Do not use.

### ***Other information***

Store at 68°F-77°F (20°C-25°C).

Store in a clean, dry place outside of direct sunlight. Protect from excessive moisture.

Some individuals may not experience pain relief until several minutes or hours after applying the patch.

### ***Inactive ingredients***

Arnica montana flower extract, Dihydroxyaluminum aminoacetate, Edetate disodium, Glycerin, Hydroxyacetophenone, Kaolin, L-Tartaric acid, Mineral oil, Polyacrylic acid, Propylene glycol, Polysorbate 80, PVP, Sodium polyacrylate, Titanium dioxide, Water.

### ***Questions or comments?***

866-747-7365

Manufactured for:

SOLA PHARMACEUTICALS LLC

Baton Rouge, LA 70810

[www.solameds.us](http://www.solameds.us)

Made in China

Triceptin


Topical Pain Patch

for temporary pain relief

NDC 70512-017-15

Qty: 15 patches

(5 per Resealable Pouch) x 3



## TRICEPTIN

lidocaine / menthol / methyl salicylate patch

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:70512-017
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>LIDOCAINE</b> (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	40 mg in 1 g
<b>METHYL SALICYLATE</b> (UNII: LAV5U5022Y) (SALICYLIC ACID - UNII:O414PZ4LPZ)	METHYL SALICYLATE	20 mg in 1 g
<b>MENTHOL</b> (UNII: L7T10EIP3A) (MENTHOL - UNII:L7T10EIP3A)	MENTHOL	10 mg in 1 g

### Inactive Ingredients

Ingredient Name	Strength
<b>ARNICA MONTANA FLOWER</b> (UNII: OZ0E5Y15PZ)	

<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)
<b>WATER</b> (UNII: 059QF0KO0R)
<b>DIHYDROXYALUMINUM AMINOACETATE</b> (UNII: DO250MG0W6)
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)
<b>MINERAL OIL</b> (UNII: T5L8T28FGP)
<b>POLYACRYLIC ACID (250000 MW)</b> (UNII: 9G2MAD7J6W)
<b>POVIDONE K90</b> (UNII: RDH86HJV5Z)
<b>SODIUM POLYACRYLATE (2500000 MW)</b> (UNII: 05I15JNI2J)
<b>HYDROXYACETOPHENONE</b> (UNII: G1L3HT4CMH)
<b>KAOLIN</b> (UNII: 24H4NWX5CO)
<b>TARTARIC ACID</b> (UNII: W4888I119H)
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)

### Product Characteristics

<b>Color</b>		<b>Score</b>	
<b>Shape</b>	RECTANGLE	<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70512-017-15	3 in 1 BOX	12/05/2022	
1		5 g in 1 POUCH; Type 2: Prefilled Drug Delivery Device/System (syringe, patch, etc.)		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	12/05/2022	

**Labeler** - SOLA Pharmaceuticals (080121345)