

## **CONZOL 1 %- miconazole spray MWI**

-----  
**Conzol™ 1%  
(miconazole nitrate)  
Lotion & Spray**

**Topical antifungal agent for dogs and cats.**

**CAUTION:** Federal law restricts this drug to use by or on the order of a licensed veterinarian.

**Not for Use in Humans  
Keep Out of Reach of Children**

Approved by FDA under ANADA # 200-362

### **DESCRIPTION:**

Conzol™ Lotion, 1% (miconazole nitrate) & Conzol™ Spray, 1% (miconazole nitrate) is a synthetic antifungal agent for use in dogs and cats. It contains: 1.15% miconazole nitrate (equivalent to 1% miconazole base by weight), polyethylene glycol 400, ethyl alcohol 55%.

### **INDICATIONS:**

Conzol™ Lotion, 1% (miconazole nitrate) & Conzol™ Spray, 1% (miconazole nitrate) is indicated for the treatment of fungal infections in dogs and cats caused by *Microsporum canis*, *Microsporum gypseum*, and *Trichophyton mentagrophytes*.

### **PRECAUTIONS:**

In the event of sensitization or irritation due to Conzol™ Lotion, 1% (miconazole nitrate) & Conzol™ Spray, 1% (miconazole nitrate), treatment should be discontinued. Avoid contact with eyes, since irritation may result. Wash hands thoroughly after administration to avoid spread of fungal infection.

### **DOSAGE AND ADMINISTRATION:**

Accurate diagnosis of the infecting organism is essential. Identification should be made either by direct microscopic examination of a mounting of infected tissue in a solution of potassium hydroxide, or by culture on an appropriate medium.

**Lotion:** Apply a light covering of Conzol™ Lotion, 1% (miconazole nitrate) to affected areas, once daily, for 2 to 4 weeks. Application is best accomplished using a gauze pad or cotton swab.

**Spray:** Spray affected areas from a distance of 2 to 4 inches to apply a light covering, once daily for 2 to 4 weeks. Do not allow pet to contact finished wood surfaces until pet

is thoroughly dried.

Medication must be continued until the infecting organism is completely eradicated as indicated by appropriate clinical or laboratory examination. If no improvement is noticed within 2 weeks, diagnosis should be re-evaluated. Difficult cases may require treatment for 6 weeks.

General measures in regard to hygiene should be observed to control sources of infection or reinfection. Clipping of hair around and over the sites of infection should be done at the start of treatment and again as necessary.

**ACTIVE INGREDIENT:**

Miconazole Nitrate.....1.15%  
(equivalent to 1% miconazole base by weight)

**STORAGE:**

Store at controlled room temperature between 15°-30°C (59°-86°F). Keep container tightly closed when not in use.

**HOW SUPPLIED:**

**Lotion:**

Conzol™ Lotion, 1% (miconazole nitrate) is available in 30 and 60 mL bottles with droppers.

**Spray:**

Conzol™ Spray, 1% (miconazole nitrate) is available in 120 mL & 240 mL bottles with spray misters.

**CAUTION:**

Federal law restricts this drug to use by or on the order of a licensed veterinarian.

**CONTACT INFORMATION:**

To report suspected adverse events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact First Priority, Inc. at (800) 650-4899 or [www.prioritycare.com](http://www.prioritycare.com). For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>

Distributed by: MWI  
Boise, ID 83705  
[www.VetOne.net](http://www.VetOne.net)  
Rev. 04/23



Approved by FDA under ANADA # 200-362



### 60 mL bottle/case label

**INDICATIONS:** For the treatment of fungal infections in dogs and cats caused by *Microsporum canis*, *Microsporum gypseum*, and *Trichophyton mentagrophytes*.

**DOSAGE:** Apply a light covering to affected areas, once daily for 2 to 4 weeks. See accompanying literature for full directions.

Avoid contact with eyes, since irritation may result.

Wash hands thoroughly after administration to avoid spread of fungal infection.

**TAKE TIME**  **OBSERVE LABEL DIRECTIONS**

NDC 13985-596-60

**VETone**  
**Conzol™ 1%**  
(miconazole nitrate)

**Lotion**

**Topical antifungal agent for dogs and cats**

**CAUTION:** Federal law restricts this drug to use by or on the order of a licensed veterinarian.

**Not for Use in Humans**  
**Keep Out of Reach of Children**

Approved by FDA under ANADA # 200-362

**V1 501008** Net Contents: 60 mL (2 fl oz)

**60 mL**

**ACTIVE INGREDIENT:**  
Miconazole Nitrate.....1.15%  
(equivalent to 1% miconazole base by weight)

**STORAGE:** Store at controlled room temperature between 15°-30°C (59°-86°F). Keep container tightly closed when not in use.

Distributed by: MWI  
Boise, ID 83705  
www.VetOne.net

Lot No.      Exp. Date

**NO VARNISH**

 3 13985 01220 4

 **MADE IN USA**  
Rev. 04/23

 **OPEN HERE**



# Conzol™ 1%

(miconazole nitrate)

## Lotion

Topical antifungal agent for dogs and cats.

**DESCRIPTION:** Conzol™ Lotion, 1% (miconazole nitrate) is a synthetic antifungal agent for use in dogs and cats. It contains: 1.15% miconazole nitrate (equivalent to 1% miconazole base by weight), polyethylene glycol 400, and ethyl alcohol 55%.

**INDICATIONS:** Conzol™ Lotion, 1% (miconazole nitrate) is indicated for the treatment of fungal infections in dogs and cats caused by *Microsporum canis*, *Microsporum gypsum*, and *Trichophyton mentagrophytes*.

**PRECAUTIONS:** In the event of sensitization or irritation due to Conzol™ Lotion, 1% (miconazole nitrate), treatment should be discontinued. Avoid contact with eyes, since irritation may result. Wash hands thoroughly after administration to avoid spread of fungal infection.

**DOSAGE AND ADMINISTRATION:** Accurate diagnosis of the infecting organism is essential. Identification should be made either by direct microscopic examination of a mounting of infected tissue in a solution of potassium hydroxide, or by culture on an appropriate medium.

**Lotion:** Apply a light covering of Conzol™ Lotion, 1% (miconazole nitrate) to affected areas, once daily, for 2 to 4 weeks. Application is best accomplished using a gauze pad or cotton swab.

Medication must be continued until the infecting organism is completely eradicated as indicated by appropriate clinical or laboratory examination.

If no improvement is noticed within 2 weeks, diagnosis should be re-evaluated. Difficult cases may require treatment for 6 weeks.

General measures in regard to hygiene should be observed to control sources of infection or reinfection. Clipping of hair around and over the sites of infection should be done at the start of treatment and again as necessary.

**HOW SUPPLIED:** Conzol™ Lotion, 1% (miconazole nitrate) is available in 30 mL & 60 mL bottles with droppers.

**CAUTION:** Federal law restricts this drug to use by or on the order of a licensed veterinarian.

**CONTACT INFORMATION:** To report suspected adverse events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact First Priority, Inc. at (800) 650-4899 or [www.prioritycare.com](http://www.prioritycare.com). For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>

Distributed by: MWM  
Boise, ID 83705  
[www.VetOne.net](http://www.VetOne.net)  
Rev. 04/23

Approved by FDA under  
ANADA # 200-362



Rectangular Ship



Quantity: **12** Bottles Conzol 1% Lotion (12 x 60 mL ea)

**V1 501008**



10 3 13985 01220 1

Lot No.

Exp. Date


**NO VARNISH**

Approved by FDA under  
ANADA # 200-362  
Rev. 04/23



Quantity: **72** Bottles Conzol 1% Lotion  
(6 inner cases of 12 x 60 mL each)

**V1 501008**



20 3 13985 01220 8

Lot No.

Exp. Date

**NO VARNISH**

Approved by FDA under  
ANADA # 200-362  
Rev. 04/23

**120 mL bottle/case label**

**INDICATIONS:** For the treatment of fungal infections in dogs and cats caused by *Microsporum canis*, *Microsporum gypseum*, and *Trichophyton mentagrophytes*.

**DOSAGE:** Apply a light covering to affected areas, once daily for 2 to 4 weeks. See accompanying literature for full directions.

Avoid contact with eyes, since irritation may result.

Wash hands thoroughly after administration to avoid spread of fungal infection.



NDC 13985-596-12

120 mL

**VET one**  
**Conzol™ 1%**  
(miconazole nitrate)

Spray

**Topical antifungal agent  
for dogs and cats**

**CAUTION:** Federal law restricts this drug to use by or on the order of a licensed veterinarian.

**Not for Use in Humans  
Keep Out of Reach of  
Children**

Approved by FDA under  
ANADA # 200-362



V1 502008

Net Contents: 120 mL (4 fl oz)

**ACTIVE INGREDIENT:**  
Miconazole Nitrate..... 1.15%  
(equivalent to 1%  
miconazole base by weight)



**STORAGE:** Store at controlled room temperature between 15°-30°C (59°- 86°F). Keep container tightly closed when not in use.

Distributed by: MWI  
Boise, ID 83705  
[www.VetOne.net](http://www.VetOne.net)  
Rev. 04/23



Lot No. Exp. Date

**NO VARNISH**





# Conzol™ 1%

(miconazole nitrate)

**Spray**

## Topical antifungal agent for dogs and cats.

**DESCRIPTION:** Conzol™ Spray, 1% (miconazole nitrate) is a synthetic antifungal agent for use in dogs and cats. It contains: 1.15% miconazole nitrate (equivalent to 1% miconazole base by weight), polyethylene glycol 400, and ethyl alcohol 55%.

**INDICATIONS:** Conzol™ Spray, 1% (miconazole nitrate) is indicated for the treatment of fungal infections in dogs and cats caused by *Microsporum canis*, *Microsporum gypseum*, and *Trichophyton mentagrophytes*.

**PRECAUTIONS:** In the event of sensitization or irritation due to Conzol™ Spray, 1% (miconazole nitrate), treatment should be discontinued. Avoid contact with eyes, since irritation may result. Wash hands thoroughly after administration to avoid spread of fungal infection.

**DOSAGE AND ADMINISTRATION:** Accurate diagnosis of the infecting organism is essential. Identification should be made either by direct microscopic examination of a mounting of infected tissue in a solution of potassium hydroxide, or by culture on an appropriate medium.

**Spray:** Spray affected areas from a distance of 2 to 4 inches to apply a light covering, once daily for 2 to 4 weeks. Do not allow pet to contact finished wood surfaces until pet is thoroughly dried.

Medication must be continued until the infecting organism is completely eradicated as indicated by appropriate clinical or laboratory examination. If no improvement is noticed within 2 weeks, diagnosis should be re-evaluated. Difficult cases may require treatment for 6 weeks.

General measures in regard to hygiene should be observed to control sources of infection or reinfection. Clipping of hair around and over the sites of infection should be done at the start of treatment and again as necessary.

**HOW SUPPLIED:** Conzol™ Spray, 1% (miconazole nitrate) is available in 120 mL & 240 mL bottles with spray misters.

**CAUTION:** Federal law restricts this drug to use by or on the order of a licensed veterinarian.

**CONTACT INFORMATION:** To report suspected adverse events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact First Priority, Inc. at (800) 650-4899 or [www.prioritycare.com](http://www.prioritycare.com). For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>

Distributed by: MWI  
Boise, ID 83705  
[www.VetOne.net](http://www.VetOne.net)  
Rev. 04/23

TAKE TIME




OBSERVE LABEL  
DIRECTIONS

Approved by FDA under  
ANADA # 200-362



Quantity: **12** Bottles Conzol 1% Spray  
(12 x 120 mL each)

**V1 502008**



10 3 13985 01221 8

Lot No.

Exp. Date

**NO VARNISH**

Approved by FDA under  
ANADA # 200-362  
Rev. 04/23



Quantity: **72** Bottles Conzol 1% Spray  
(6 inner cases of 12 x 120 mL each)

**V1 502008**



20 3 13985 01221 5

Lot No. Exp. Date

**NO VARNISH**

Approved by FDA under  
ANADA # 200-362  
Rev. 04/23

**240 mL bottle/case label**

**INDICATIONS:** For the treatment of fungal infections in dogs and cats caused by *Microsporum canis*, *Microsporum gypseum*, and *Trichophyton mentagrophytes*.

**DOSAGE:** Apply a light covering to affected areas, once daily for 2 to 4 weeks. See accompanying literature for full directions.

Avoid contact with eyes, since irritation may result.

Wash hands thoroughly after administration to avoid spread of fungal infection.

NDC 13985-596-24

240 mL

  
**Conzol™ 1%**  
(miconazole nitrate)

Spray

**Topical antifungal agent  
for dogs and cats**

**CAUTION:** Federal law restricts this drug to use by or on the order of a licensed veterinarian.

**Not for Use in Humans  
Keep Out of Reach  
of Children**

Approved by FDA under  
ANADA # 200-362



V1 503008

Net Contents: 240 mL (8 fl oz)

**ACTIVE INGREDIENT:**  
Miconazole Nitrate..... 1.15%  
(equivalent to 1% miconazole  
base by weight)



**STORAGE:** Store at controlled room temperature between 15°-30°C (59°-86°F). Keep container tightly closed when not in use.

Distributed by: MW  
Boise, ID 83705  
[www.VetOne.net](http://www.VetOne.net)  
Rev. 04/23



Lot No.      Exp. Date

**NO VARNISH**



TAKE TIME  OBSERVE LABEL DIRECTIONS



# Conzol™ 1%

(miconazole nitrate)

Spray

**Topical antifungal agent for dogs and cats.**

**DESCRIPTION:** Conzol™ Spray, 1% (miconazole nitrate) is a synthetic antifungal agent for use in dogs and cats. It contains: 1.15% miconazole nitrate (equivalent to 1% miconazole base by weight), polyethylene glycol 400, and ethyl alcohol 55%.

**INDICATIONS:** Conzol™ Spray, 1% (miconazole nitrate) is indicated for the treatment of fungal infections in dogs and cats caused by *Microsporum canis*, *Microsporum gypseum*, and *Trichophyton mentagrophytes*.

**PRECAUTIONS:** In the event of sensitization or irritation due to Conzol™ Spray, 1% (miconazole nitrate), treatment should be discontinued. Avoid contact with eyes, since irritation may result. Wash hands thoroughly after administration to avoid spread of fungal infection.

**DOSAGE AND ADMINISTRATION:** Accurate diagnosis of the infecting organism is essential. Identification should be made either by direct microscopic examination of a mounting of infected tissue in a solution of potassium hydroxide, or by culture on an appropriate medium.

**Spray:** Spray affected areas from a distance of 2 to 4 inches to apply a light covering, once daily for 2 to 4 weeks. Do not allow pet to contact finished wood surfaces until pet is thoroughly dried.

Medication must be continued until the infecting organism is completely eradicated as indicated by appropriate clinical or laboratory examination. If no improvement is noticed within 2 weeks, diagnosis should be re-evaluated. Difficult cases may require treatment for 6 weeks.

General measures in regard to hygiene should be observed to control sources of infection or reinfection. Clipping of hair around and over the sites of infection should be done at the start of treatment and again as necessary.

**HOW SUPPLIED:** Conzol™ Spray, 1% (miconazole nitrate) is available in 120 mL & 240 mL bottles with spray misters.

**CAUTION:** Federal law restricts this drug to use by or on the order of a licensed veterinarian.

**CONTACT INFORMATION:** To report suspected adverse events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact First Priority, Inc. at (800) 650-4899 or [www.prioritycare.com](http://www.prioritycare.com). For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>

Distributed by: MWI  
Boise, ID 83705  
[www.VetOne.net](http://www.VetOne.net)  
Rev. 04/23


TAKE TIME  OBSERVE LABEL  
DIRECTIONS

Approved by FDA under  
ANADA # 200-362



Quantity: **12** Bottles Conzol 1% Spray  
(12 x 240 mL each)

**V1 503008**



10 3 13985 01222 5

Lot No.

Exp. Date


**NO VARNISH**

Approved by FDA under  
ANADA # 200-362  
Rev. 04/23



Quantity: **72** Bottles Conzol 1% Spray  
(6 inner cases of 12 x 240 mL each)

**V1 503008**



20 3 13985 01222 2

Lot No. Exp. Date

**NO VARNISH**

Approved by FDA under  
ANADA # 200-362  
Rev. 04/23

## CONZOL 1 %

miconazole spray

### Product Information

<b>Product Type</b>	PRESCRIPTION ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:13985-596
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>MICONAZOLE NITRATE</b> (UNII: VW4H1CYW1K) (MICONAZOLE - UNII:7NNO0D7S5M)	MICONAZOLE NITRATE	11.5 mg in 1 mL

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:13985-596-60	72 in 1 CASE		

<b>1</b>		60 mL in 1 BOTTLE, DROPPER		
<b>2</b>	NDC:13985-596-12	72 in 1 CASE		
<b>2</b>		120 mL in 1 BOTTLE, SPRAY		
<b>3</b>	NDC:13985-596-24	72 in 1 CASE		
<b>3</b>		240 mL in 1 BOTTLE, SPRAY		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANADA	ANADA200362	09/21/2023	

**Labeler** - MWI (019926120)

**Registrant** - FIRST PRIORITY INCORPORATED (179925722)

## Establishment

Name	Address	ID/FEI	Business Operations
FIRST PRIORITY INCORPORATED		179925722	manufacture

Revised: 2/2026

MWI