CONZOL 1 %- micaonazole spray MWI

Conzol[™] 1% (miconazole nitrate) Lotion & Spray

Topical antifungal agent for dogs and cats.

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

Not for Use in Humans Keep Out of Reach of Children

Approved by FDA under ANADA # 200-362

DESCRIPTION:

Conzol[™] Lotion, 1% (miconazole nitrate) & Conzol[™] Spray, 1% (miconazole nitrate) is a synthetic antifungal agent for use in dogs and cats. It contains: 1.15% miconzole nitrate (equivalent to 1% miconzole base by weight), polyethylene glycol 400, ethyl alcohol 55%.

INDICATIONS:

Conzol[™] Lotion, 1% (miconzole nitrate) & Conzol[™] Spray, 1% (miconzole nitrate) is indicated for the treatment of fungal infections in dogs and cats caused by *Microsporum canis*, *Microsporum gypseum*, and *Trichophyton mentagrophytes*.

PRECAUTIONS:

In the event of sensitization or irritation due to Conzol[™] Lotion, 1% (miconazole nitrate) & Conzol[™] Spray, 1% (miconazole nitrate), treatment should be discontinued. Avoid contact with eyes, since irritation may result. Wash hands thoroughly after administration to avoid spread of fungal infection.

DOSAGE AND ADMINISTRATION:

Accurate diagnosis of the infecting organism is essential. Identification should be made either by direct microscopic examination of a mounting of infected tissue in a solution of potassium hydroxide, or by culture on an appropriate medium.

Lotion: Apply a light covering of $Conzol^{\text{TM}}$ Lotion, 1% (miconazole nitrate) to affected areas, once daily, for 2 to 4 weeks. Application is best accomplished using a gauze pad or cotton swab.

Spray: Spray affected areas from a distance of 2 to 4 inches to apply a light covering, once daily for 2 to 4 weeks. Do not allow pet to contact finished wood surfaces until pet

is thoroughly dried.

Medication must be continued until the infecting organism is completely eradicated as indicated by appropriate clinical or laboratory examination. If no improvement is noticed within 2 weeks, diagnosis should be re-evaluated. Difficult cases may require treatment for 6 weeks.

General measures in regard to hygiene should be observed to control sources of infection or reinfection. Clipping of hair around and over the sites of infection should be done at the start of treatment and again as necessary.

ACTIVE INGREDIENT:

Miconazole Nitrate......1.15% (equivalent to 1% miconazole base by weight)

STORAGE:

Store at controlled room temperature between 15°-30°C (59°-86°F). Keep container tightly closed when not in use.

HOW SUPPLIED:

Lotion:

Conzol $^{\text{m}}$ Lotion, 1% (miconazole nitrate) is available in 30 and 60 mL bottles with droppers.

Spray:

Conzol™ Spray, 1% (miconazole nitrate) is available in 120 mL & 240 mL bottles with spray misters.

CAUTION:

Federal law restricts this drug to use by or on the order of a licensed veterinarian.

CONTACT INFORMATION:


To report suspected adverse events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact First Priority, Inc. at (800) 650-4899 or www.prioritycare.com. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or http://www.fda.gov/reportanimalae

Distributed by: MWI Boise, ID 83705 www.VetOne.net Rev. 04/23





60 mL bottle/case label





(miconazole nitrate)



Topical antifungal agent for dogs and cats.

DESCRIPTION: Conzol** Lotion, 1% (miconazole nitrate) is a synthetic antifungal agent for use in dogs and cats. It contains: 1.15% miconazole nitrate (equivalent to 1% miconazole base by weight), polyethylene glycol 400, and ethyl alcohol 55%.

INDICATIONS: Conzol** Lotion, 1% (miconazole nitrate) is indicated for the treatment of fungal infections in dogs and cats caused by Microsporum canis, Microsporum gypseum, and Trichophyton mentagrophytes.

PRECAUTIONS: In the event of sensitization or imitation due to Conzol**
Lotion, 1% (miconazole nitrate), treatment should be discontinued. Avoid
contact with eyes, since irritation may result. Wash hands thoroughly
after administration to avoid spread of fungal infection.

DOSAGE AND ADMINISTRATION: Accurate diagnosis of the infecting organism is essential. Identification should be made either by direct microscopic examination of a mounting of infected tissue in a solution of potassium hydroxide, or by culture on an appropriate medium.

Lotion: Apply a light covering of Conzol™ Lotion, 1% (miconazole nitrate) to affected areas, once daily, for 2 to 4 weeks. Application is best accomplished using a gauze pad or cotton swab.

Medication must be continued until the infecting organism is completely eradicated as indicated by appropriate clinical or laboratory examination.

If no improvement is noticed within 2 weeks, diagnosis should be re-evaluated. Difficult cases may require treatment for 6 weeks.

General measures in regard to hygiene should be observed to control sources of infection or reinfection. Clipping of hair around and over the sites of infection should be done at the start of treatment and again as necessary.

HOW SUPPLIED: Conzol** Lotion, 1% (miconazole nitrate) is available in 30 mL & 60 mL bottles with droppers.

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

CONTACT INFORMATION: To report suspected adverse events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact First Priority, Inc. at (800) 650-4899 or oww.prioritycare.com. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or http://www.fda.gov/reportanimalae

Distributed by: MWI Boise, ID 83705 www.VetOne.net Rev. 04/23 Approved by FDA under ANADA # 200-362

23







120 mL bottle/case label

INDICATIONS: For the treatment of fungal infections in dogs and cats caused by Microsporum canis, Microsporum gypseum, and Trichophyton mentagrophytes.

DOSAGE: Apply a light covering to affected areas, once daily for 2 to 4 weeks. See accompanying literature for full directions.

Avoid contact with eyes, since irritation may result.

Wash hands thoroughly after administration to avoid spread of fungal infection.

TAKE TIME OBSERVE LABEL DIRECTIONS

NDC 13985-596-12



120 mL

(miconazole nitrate)



ACTIVE INGREDIENT:

Miconazole Nitrate..... 1.15% (equivalent to 1% miconazole base by weight)



STORAGE: Store at controlled room temperature between 15°-30°C (59°- 86°F). Keep container tightly closed when not in use.

Distributed by: MWI Boise, ID 83705 www.VetOne.net Rev. 04/23







(miconazole nitrate)



Topical antifungal agent for dogs and cats.

DESCRIPTION: Conzol™ Spray, 1% (miconazole nitrate) is a synthetic antifungal agent for use in dogs and cats. It contains: 1.15% miconazole nitrate (equivalent to 1% miconazole base by weight), polyethylene glycol 400, and ethyl alcohol 55%.

INDICATIONS: Conzol™ Spray, 1% (miconazole nitrate) is indicated for the treatment of fungal infections in dogs and cats caused by Microsporum canis, Microsporum gypseum, and Trichophyton mentagrophytes.

PRECAUTIONS: In the event of sensitization or irritation due to Conzol™ Spray, 1% (miconazole nitrate), treatment should be discontinued. Avoid contact with eyes, since irritation may result. Wash hands thoroughly after administration to avoid spread of fungal infection.

DOSAGE AND ADMINISTRATION: Accurate diagnosis of the infecting organism is essential. Identification should be made either by direct microscopic examination of a mounting of infected tissue in a solution of potassium hydroxide, or by culture on an appropriate medium.

Spray: Spray affected areas from a distance of 2 to 4 inches to apply a light covering, once daily for 2 to 4 weeks. Do not allow pet to contact finished wood surfaces until pet is thoroughly dried.

Medication must be continued until the infecting organism is completely eradicated as indicated by appropriate clinical or laboratory examination. If no improvement is noticed within 2 weeks, diagnosis should be re-evaluated. Difficult cases may require treatment for 6 weeks.

General measures in regard to hygiene should be observed to control sources of infection or reinfection. Clipping of hair around and over the sites of infection should be done at the start of treatment and again as necessary.

HOW SUPPLIED: Conzol™ Spray, 1% (miconazole nitrate) is available in 120 mL & 240 mL bottles with spray misters.

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

CONTACT INFORMATION: To report suspected adverse events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact First Priority, Inc. at (800) 650-4899 or www.prioritycare.com. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or http://www.fda.gov/reportanimalae

Distributed by: MWI Boise, ID 83705 www.VetOne.net Rev. 04/23



Approved by FDA under ANADA # 200-362



Quantity: **12** Bottles Conzol 1% Spray (12 x 120 mL each)

V1 502008



10 3 13985 01221 8

Lot No.

Exp. Date

NO VARNISH

Approved by FDA under ANADA # 200-362 Rev. 04/23



Quantity: **72** Bottles Conzol 1% Spray (6 inner cases of 12 x 120 mL each)

V1 502008



20 3 13985 01221 5

NO VARNISH

Approved by FDA under ANADA # 200-362 Rev. 04/23

240 mL bottle/case label

INDICATIONS: For the treatment of fungal infections in dogs and cats caused by Microsporum canis, Microsporum gypseum, and Trichophyton mentagrophytes.

DOSAGE: Apply a light covering to affected areas, once daily for 2 to 4 weeks. See accompanying literature for full directions.

Avoid contact with eyes, since irritation may result.

Wash hands thoroughly after administration to avoid spread of fungal infection. NDC 13985-596-24

240 mL




(miconazole nitrate)



ACTIVE INGREDIENT: Miconazole Nitrate.......1.15% (equivalent to 1% miconazole base by weight)



STORAGE: Store at controlled room temperature between 15°-30°C (59°-86°F). Keep container tightly closed when not in use.



Distributed by: MWI Boise, ID 83705 www.VetOne.net Rev. 04/23









Conzol™ 1%

(miconazole nitrate)



Topical antifungal agent for dogs and cats.

DESCRIPTION: Conzol™ Spray, 1% (miconazole nitrate) is a synthetic antifungal agent for use in dogs and cats. It contains: 1.15% miconazole nitrate (equivalent to 1% miconazole base by weight), polyethylene glycol 400, and ethyl alcohol 55%.

INDICATIONS: Conzol™ Spray, 1% (micronazole nitrate) is indicated for the treatment of fungal infections in dogs and cats caused by Microsporum canis, Microsporum gypseum, and Trichophyton mentagrophytes.

PRECAUTIONS: In the event of sensitization or irritation due to Conzol™ Spray, 1% (miconazole nitrate), treatment should be discontinued. Avoid contact with eyes, since irritation may result. Wash hands thoroughly after administration to avoid spread of fungal infection.

DOSAGE AND ADMINISTRATION: Accurate diagnosis of the infecting organism is essential. Identification should be made either by direct microscopic examination of a mounting of infected tissue in a solution of potassium hydroxide, or by culture on an appropriate medium.

Spray: Spray affected areas from a distance of 2 to 4 inches to apply a light covering, once daily for 2 to 4 weeks. Do not allow pet to contact finished wood surfaces until pet is thoroughly dried.

Medication must be continued until the infecting organism is completely eradicated as indicated by appropriate clinical or laboratory examination. If no improvement is noticed within 2 weeks, diagnosis should be re-evaluated. Difficult cases may require treatment for 6 weeks.

General measures in regard to hygiene should be observed to control sources of infection or reinfection. Clipping of hair around and over the sites of infection should be done at the start of treatment and again as necessary.

HOW SUPPLIED: Conzol™ Spray, 1% (miconazole nitrate) is available in 120 mL & 240 mL bottles with spray misters.

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

CONTACT INFORMATION: To report suspected adverse events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact First Priority, Inc. at (800) 650-4899 or www.prioritycare.com. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or http://www.fda.gov/reportanimalae

Distributed by: MWI Boise, ID 83705 www.VetOne.net Rev. 04/23



Approved by FDA under ANADA # 200-362



Quantity: **12** Bottles Conzol 1% Spray (12 x 240 mL each)

V1 503008



10 3 13985 01222 5

Lot No.

Exp. Date


NO VARNISH

Approved by FDA under ANADA # 200-362 Rev. 04/23



Quantity: **72** Bottles Conzol 1% Spray (6 inner cases of 12 x 240 mL each)

V1 503008



20 3 13985 01222 2

NO VARNISH

Approved by FDA under ANADA # 200-362 Rev. 04/23

CONZOL 1 %

micaonazole spray

Product Information

Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:13985-596
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
	MICONAZOLE NITRATE	11.5 mg in 1 mL

F	Packaging			
#	tem Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:13985-596-60	72 in 1 CASE		

1		60 mL in 1 BOTTLE, DROPPER
2	NDC:13985-596-12	72 in 1 CASE
2		120 mL in 1 BOTTLE, SPRAY
3	NDC:13985-596-24	72 in 1 CASE
3		240 mL in 1 BOTTLE, SPRAY

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANADA	ANADA200362	09/21/2023	

Labeler - MWI (019926120)

Registrant - FIRST PRIORITY INCORPORATED (179925722)

Establishment			
Name	Address	ID/FEI	Business Operations
FIRST PRIORITY INCORPORATED		179925722	manufacture

Revised: 9/2023 MW