

SODIUM FLUORIDE CHEWABLE- sodium fluoride tablet Bryant Ranch Prepack

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Sodium Fluoride Chewable Tablets

Sodium Fluoride Chewable Tablets

This insert is for 1.0 mg strength tablets, 0.5 mg strength tablets, and 0.25 mg strength tablets.

Rx Only

This product is a prescription product for the clinical dietary management of the metabolic processes of caries prophylaxis.

Water F⁻ Content			
Ages	0 ppm F⁻ to <0.3 ppm F⁻	0.3 ppm F⁻ to 0.6ppm F⁻	>0.6ppm F⁻
3 yrs. to 6 yrs.	0.5mg*	0.25mg*	0
>6yrs. to 16 yrs.	1.0mg*	0.5mg*	0

* Per day

Active Ingredient:Fluoride (as Sodium Fluoride)

0.25 mg / 0.5 mg / 1.0 mg

Inactive Ingredients:citric acid, magnesium stearate, malic acid, microcrystalline cellulose, orange flavor, sucralose, talc, xylitol.

Caution:Do not eat or drink dairy products within one hour of fluoride administration. Incompatibility of fluoride with dairy foods has been reported due to formation of calcium fluoride which is poorly absorbed.

CLINICAL PHARMACOLOGY

Sodium Fluoride acts systemically (before tooth eruption) and topically (post-eruption) by increasing tooth resistance to acid dissolution, by promoting remineralization, and by inhibiting the cariogenic microbial process.

INDICATIONS AND USAGE

For once daily self-applied systemic use as a dental caries preventive in pediatric patients. It has been established that ingestion of fluoridated drinking water (1 ppm F⁻) during the period of tooth development results in a significant decrease in the incidence

of dental caries. Sodium Fluoride Chewable Tablets were developed to provide systemic fluoride for use as a supplement in pediatric patients from age 3 years to age 16 years and older living in areas where the drinking water fluoride contents does not exceed 0.6 ppm F⁻.

CONTRAINDICATIONS

Fluoride 1.0 mg Tablets are contraindicated when the fluoride content of drinking water is >0.3 ppm F⁻ and should not be administered to pediatric patients under 6 years. Fluoride 0.5 mg Tablets are contraindicated when the fluoride content of drinking water is more than 0.6 ppm F⁻ and should not be administered to pediatric patients under age 6 when the fluoride content of drinking water is 0.3 ppm F⁻ or more or to pediatric patients under age 3 years. Fluoride 0.25 mg Tablets are contraindicated when the fluoride content of drinking water is more than 0.6 ppm F⁻ and should not be administered to pediatric patients under age 3 years when the fluoride content of drinking water is 0.3 ppm F⁻ or more. Do not administer Sodium Fluoride Chewable Tablets (any strength) to pediatric patients under age 3 years. Sodium Fluoride Chewable Tablets (any strength) are not indicated for use in adults.

OVERDOSAGE

Prolonged daily ingestion of excessive fluoride will result in varying degrees of fluorosis. Accidental ingestion of large amounts of fluoride may result in acute burning in the mouth and sore tongue. Nausea, vomiting, and diarrhea may occur soon after ingestion (within 30 minutes) and are accompanied by salivation, hematemesis, and epigastric cramping abdominal pain. These symptoms may persist for 24 hours. If less than 5 mg fluoride/kg body weight (i.e., less than 2.3 mg fluoride/lb body weight) have been ingested, give calcium (e.g., milk) orally to relieve gastrointestinal symptoms and observe for a few hours. If more than 5 mg fluoride/kg body weight (i.e., more than 2.3 mg fluoride/lb body weight) have been ingested, induce vomiting, give orally soluble calcium (e.g., milk, 5% calcium gluconate or calcium lactate solution) and immediately seek medical assistance. For accidental ingestion of more than 15 mg fluoride/kg body weight (i.e., more than 6.9 mg fluoride/lb body weight), induce vomiting and admit immediately to a hospital facility. A treatment dose of Sodium Fluoride Chewable Tablets contains 0.25, 0.5 or 1.0 mg fluoride. The treatment of choice depends upon the age of the child and the water fluoride content. A bottle of 120 0.25 mg tablets contains 30 mg fluoride. A bottle of 120 0.5 mg tablets contains 60 mg fluoride. A bottle of 120 1.0 mg tablets contains 120 mg fluoride. [The total amount of sodium fluoride in a bottle of 120 Fluoride Chewable Tablets (all strengths) conforms with the recommended amount of the American Dental Association for the maximum to be dispensed at one time for safety purposes.]

WARNING

Prolonged daily ingestion of quantities greater than the recommended amount may result in various degrees of dental fluorosis in pediatric patients under age 6 years, especially if the water fluoridation exceeds 0.6 ppm. Read directions carefully before using. This product, as all chewable tablets, is not recommended for children under age 3 due to risk of choking. Keep out of the reach of infants and children.

PRECAUTIONS

General

Please refer to the **CONTRAINDICATIONS, WARNINGS** and **OVERDOSAGE** sections for overdose concerns. Use in pediatric patients below the age of 3 years is not recommended by current American Dental Associations and American Academy of Pediatrics guidelines.

Carcinogenesis, Mutagenesis, Impairment of Fertility

In a study conducted in rodents, no carcinogenesis was found in male and female rats treated with fluoride at dose levels ranging from 4.1 to 9.1 mg/kg of body weight. Equivocal evidence of carcinogenesis was reported for male rats treated with 2.5 and 4.1 mg/kg of body weight. In a second study, no carcinogenesis was observed in rats, males or females, treated with fluoride up to 11.3 mg/kg of body weight. This dose is at least 400 times greater than the recommended daily dose of Sodium Fluoride Chewable Tablets. Fluoride ion is not mutagenic in standard bacterial systems. It has been shown that fluoride ion has potential to induce chromosome aberrations in cultured human and rodent cells at doses much higher than those to which humans are exposed. In vivo data is conflicting. Some studies report chromosome damage in rodents while other studies using similar protocols report negative results. Potential adverse reproductive effects of fluoride exposure in humans has not been adequately evaluated. Adverse effects on reproduction were reported for rats, mice, fox, and cattle exposed to 100 ppm or greater concentrations of fluoride in their diet or drinking water. Other studies conducted in rats demonstrated that lower doses of fluoride (5 mg/kg of body weight) did not result in impaired fertility and reproductive capabilities. This dose is approximately 200 times greater than the recommended daily dose of Sodium Fluoride Chewable Tablets.

Pregnancy

Teratogenic Effects

Pregnancy Category B

It has been shown that fluoride crosses the placenta of rats, but only 0.01% of the amount administered is incorporated in fetal tissue. Animal studies (rats, mice, rabbits) have shown that fluoride is not a teratogen. Maternal exposure to 12.2 mg fluoride/kg of body weight (rats) or 13.1 mg/kg of body weight (rabbits) did not affect the litter size or fetal weight and did not increase the frequency of skeletal or visceral malformations. Epidemiological studies conducted in areas with high levels of naturally fluoridated water showed no increase in birth defects. Heavy exposure to fluoride during in utero development may result in skeletal fluorosis which becomes evident in childhood.

Nursing Mothers

It is not known if fluoride is excreted in human milk. However, many drugs are excreted in human milk and caution should be exercised when Sodium Fluoride Chewable Tablets are administered to a nursing woman. Reduced milk production was reported in farm raised fox when the animals were fed a diet containing a high concentration of fluoride

(98-137 mg/kg of body weight). No adverse effects on parturition, lactation, or offspring were seen in rats administered fluoride up to 5 mg/kg of body weight. This dose is at least 200 times greater than the recommended daily dose of Sodium Fluoride Chewable Tablets.

Geriatric Use

Sodium Fluoride Chewable Tablets (any strength) are not indicated for use in geriatric patients.

Pediatric Use

The use of Sodium Fluoride Chewable Tablets as a caries preventive in pediatric age groups 3 years to 16 years is supported by evidence from adequate and well controlled studies on fluoride supplementation from birth through adolescence.

ADVERSE REACTIONS

Allergic rash and other idiosyncrasies have been rarely reported.

DOSAGE AND ADMINISTRATION

(for dosage see table)

Dissolve in the mouth or chew before swallowing, preferably at bedtime after brushing teeth. See schedule below to determine dosage.

HOW SUPPLIED

Each tablet contains 0.5 mg Fluoride from 1.1 mg Sodium Fluoride. White to off-white, orange flavor, round-shaped chewable tablet. Debossed with "M161" on one side.

NDC 63629-1132-1 (120 count).

Repackaged/Relabeled by:
Bryant Ranch Prepack, Inc.
Burbank, CA 91504

STORAGE

Store at 25° C (77° F); excursions permitted to 15° - 30° C (59° - 86° F). See USP Controlled Room Temperature. Protect from light and moisture.

REFERENCES

1. Accepted Dental Therapeutics, Ed. 40, American Dental Association, Chicago, 399-402 (1984).
2. J. Jakush, New Fluoride Schedule Adopted, ADA News, 12, 14 (May 16, 1994).
3. Aasenden, R., and Peebles, T.C. "Effects of Fluoride Supplementation From Birth on

Dental Caries and Fluorosis in Teenaged Children", Arch. Oral, Biol., 23, 111-115 (1974).

4. Hamber, L. "Controlled Trial of Fluoride Vitamin Drops for Prevention of Caries in Children", Lancet, 1, 441-442 (1971).

5. Hennon, D.K. Stookey, G.K. and Veiswanter, B.B. "Fluoride-Vitamin Supplements: Effects on Dental Caries and Fluorosis When Used in Areas with Suboptimum Fluoride in the Water Supply", JADA, 95, 965-971 (1977).

Dispense in a tight, light-resistant container with a child-resistant closure as defined in the USP/NF.

Warning: Keep this and all medications out of the reach of children. In case of accidental overdose, seek professional assistance or contact a poison control center immediately.

All prescription substitutions using this product shall be pursuant to state statutes as applicable. This is not an Orange Book product.

Manufactured by:

Method Pharmaceuticals, LLC
Fort Worth, TX 76118

Rev. 07/2018

Sodium Fluoride Tablet Chewable 1.1 #120



GTIN
Lot
Exp
SN

Each tablet contains: Fluoride ion, 0.5 mg from 1.1 mg Sodium Fluoride. Sugar & dye free. Inactive Ingredients: citric acid, magnesium stearate, malic acid, microcrystalline cellulose, orange flavor, sucralose, talc, xylitol.

Scan Package Insert QR Code for full prescribing information.

Store in a cool & dry place at a controlled room temperature 20°-25°C (68°-77°F) excursions permitted between 15°-30°C (59°-86°F)



Package
Insert

Store in original container.

NDC 63629-1132-1

Sodium Fluoride Chewable
Tablets

0.5 mg



Relabeled by:
Bryant Ranch Prepack, Inc.
Burbank, CA 91504 USA

Rx only
120 Tablets
Manufactured by:
Method
Pharmaceutical LLC



Extended label

Description: White to off-white, orange flavor, round shaped chewable tablet. Debossed with "M161" on one side. **Indications and Usage:** Each 1.1 mg Sodium Fluoride Tablet provides 0.5 mg of fluoride in chewable tablet form for children 3-6 years of age in areas where the concentration of fluoride in drinking water is less than 0.3 ppm, and for children 6-16 years of age where drinking water contains between 0.3 to 0.6 ppm of fluoride. **Dosage and Administration:** Scan Package Insert QR Code to determine the dosage. Dissolve in mouth or chew before swallowing preferably at bedtime after brushing teeth. **Caution:** Do not eat or drink dairy products within one hour of fluoride administration. **WARNING: KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN. IN CASE OF ACCIDENTAL OVERDOSAGE, SEEK PROFESSIONAL ASSISTANCE OR CONTACT A POISON CONTROL CENTER IMMEDIATELY.** Dispense in a tight, light-resistant container with a child-resistant closure (USP/NF). All prescription substitutions using this product shall be pursuant to state statutes as applicable. **This is not an Orange Book product. Call your doctor for medical advice about side effects. You may report side effects to Method Pharmaceuticals, LLC at 877-250-3427 or to the FDA at 1-800-FDA-1088.**

SODIUM FLUORIDE CHEWABLE

sodium fluoride tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:63629-1132(NDC:58657-161)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SODIUM FLUORIDE (UNII: 8ZYQ1474W7) (FLUORIDE ION - UNII:Q80VPU408O)	FLUORIDE ION	0.5 mg

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
MAGNESIUM STEARATE (UNII: 70097M6130)	
MALIC ACID (UNII: 817L1N4CKP)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

TALC (UNII: 7SEV7J4R1U)

XYLITOL (UNII: VCQ006KQ1E)

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	8mm
Flavor	ORANGE	Imprint Code	M161
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63629-1132-1	120 in 1 BOTTLE; Type 0: Not a Combination Product	10/21/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		08/17/2018	

Labeler - Bryant Ranch Prepack (171714327)

Registrant - Bryant Ranch Prepack (171714327)

Establishment

Name	Address	ID/FEI	Business Operations
Bryant Ranch Prepack		171714327	repack(63629-1132) , relabel(63629-1132)

Revised: 5/2024

Bryant Ranch Prepack