

**CHILDRENS IBUPROFEN- ibuprofen suspension**  
**Padagis Israel Pharmaceuticals Ltd**

-----  
**Children's Ibuprofen Drug Facts**

**Active ingredient (in each 5 mL)**

Ibuprofen 100 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

**Purposes**

Pain reliever/fever reducer

**Uses**

temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more

than directed or for longer than directed.

**Sore throat warning:** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

### **Do not use**

- if the child has ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

### **Ask a doctor before use if**

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

### **Ask a doctor or pharmacist before use if the child is**

- under a doctor's care for any serious condition
- taking any other drug

### **When using this product**

- give with food or milk if stomach upset occurs

### **Stop use and ask a doctor if**

- child experiences any of the following signs of stomach bleeding:
  - feels faint
  - vomits blood
  - has bloody or black stools
  - has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- the child does not get any relief within first day (24 hours) of treatment

- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed**
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.
- if needed, repeat dose every **6-8 hours**
- do not use more than **4 times a day**
- replace original bottle cap to maintain child resistance
- wash dosage cup after each use

**Dosing Chart**

<b>Weight (lb)</b>	<b>Age (yr)</b>	<b>Dose (mL)**</b>
under 24 lbs	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

\*\*or as directed by a doctor

**Other information**

- **each 5 mL contains:** sodium 2 mg
- **do not use if printed neckband is broken or missing**
- store at 20-25°C (68-77°F)
- do not freeze

**Inactive ingredients**

anhydrous citric acid, FD&C red #40, glycerin, high fructose corn syrup, hypromellose, natural and artificial bubblegum flavors, polysorbate 80, purified water, sodium benzoate, sorbitol solution, xanthan gum

## **Questions or comments?**

**1-800-719-9260**

### **Principal Display Panel**

Compare to Children's Motrin<sup>®</sup> active ingredient

Children's Ibuprofen Oral Suspension

100 mg per 5mL

Pain Reliever/Fever Reducer (NSAID)


For Ages 2 to 11 Years

- Lasts up to 8 Hours
- Alcohol Free
- Gluten Free

Bubble Gum Flavor

4 FL OZ (118 mL)

Padagis<sup>®</sup>



**Drug Facts (continued)**

- right before or after heart surgery
- Ask a doctor before use if**
  - stomach bleeding warning applies to your child
  - child has a history of stomach problems, such as heartburn
  - child has problems or serious side effects from taking pain relievers or fever reducers
  - child has not been drinking fluids
  - child has lost a lot of fluid due to vomiting or diarrhea
  - child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
  - child has asthma
  - child is taking a diuretic

- Ask a doctor or pharmacist before use if the child is**
  - under a doctor's care for any serious condition
  - taking any other drug

- When using this product**
  - give with food or milk if stomach upset occurs

- Stop use and ask a doctor if**
  - child experiences any of the following signs of stomach bleeding:
    - feels faint
    - vomits blood
    - has bloody or black stools
    - has stomach pain that does not get better
  - child has symptoms of heart problems or stroke:
    - chest pain
    - trouble breathing
    - weakness in one part or side of body
    - slurred speech
    - leg swelling
  - the child does not get any relief within first day (24 hours) of treatment
  - fever or pain gets worse or lasts more than 3 days
  - redness or swelling is present in the painful area
  - any new symptoms appear

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

- Directions**
  - this product does not contain directions or complete warnings for adult use
  - do not give more than directed

**Drug Facts (continued)**

- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.
- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day
- replace original bottle cap to maintain child resistance
- wash dosage cup after each use

**Dosing Chart**

Weight (lb)	Age (yr)	Dose (mL)**
under 24 lbs	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

\*\*or as directed by a doctor

**Other information**

- each 5 mL contains: sodium 2 mg
- do not use if printed neckband is broken or missing
- store at 20-25°C (68-77°F)
- do not freeze

**Inactive ingredients** anhydrous citric acid, FD&C red #40, glycerin, high fructose corn syrup, hypromellose, natural and artificial bubblegum flavors, polysorbate 80, purified water, sodium benzoate, sorbitol solution, xanthan gum

**Questions or comments?**  
1-800-719-9260

**Padagis.**  
Manufactured for Padagis®  
Minneapolis, MN 55427  
www.padagis.com

**OTC** via Rx  
If a prescription is written for this product it may be covered by your health plan.

Children's  
**Ibuprofen**  
Oral Suspension  
100 mg per 5 mL  
Pain Reliever/Fever Reducer  
(NSAID)

NDC 45802-140-26

NDC 45802-140-26

Compare to  
Children's Motrin®  
active ingredient

Children's  
**Ibuprofen**  
Oral Suspension

100 mg per 5 mL

Pain Reliever/Fever Reducer  
(NSAID)

- For Ages 2 to 11 Years
- Lasts up to 8 Hours
- Alcohol Free
- Gluten Free



Bubble Gum Flavor

4 FL OZ (118 mL)

**Padagis.**

**Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.**

**Drug Facts**

**Active ingredient** *Purposes*  
(in each 5 mL)  
Ibuprofen 100 mg (NSAID)\*.....Pain reliever/  
fever reducer

\*nonsteroidal anti-inflammatory drug

**Uses temporarily:**

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash

If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

takes more or for a longer time than directed **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **Sore throat warning:** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

**Do not use**

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer



16626 RT C5

# CHILDRENS IBUPROFEN

ibuprofen suspension

## Product Information

**Product Type**

HUMAN OTC DRUG

**Item Code (Source)**

NDC:45802-140

**Route of Administration**

ORAL

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	100 mg in 5 mL

Inactive Ingredients	
Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics			
Color	PINK (light)	Score	
Shape		Size	
Flavor	BUBBLE GUM	Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:45802-140-26	1 in 1 CARTON	04/09/2014	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074937	04/09/2014	

**Labeler** - Padagis Israel Pharmaceuticals Ltd (600093611)