

**VICKS DAYQUIL AND VICKS NYQUIL HIGH BLOOD PRESSURE COLD AND FLU-  
acetaminophen, dextromethorphan hydrobromide, doxylamine succinate  
The Procter & Gamble Manufacturing Company**

-----  
**VICKS ® DayQuil™ NyQuil™ HIGH BLOOD PRESSURE COLD & FLU  
Liquid Convenience Pack**

**NyQuil HIGH BLOOD PRESSURE Cold & Flu**

***Drug Facts***

**Active ingredients (in each 30 mL)**

**Acetaminophen 650 mg**  
Dextromethorphan HBr 30 mg  
Doxylamine succinate 12.5 mg

**Purpose**

Pain reliever/ fever reducer  
Cough suppressant  
Antihistamine

**Uses**

temporarily relieves common cold/flu symptoms:

- cough due to minor throat & bronchial irritation
- sore throat
- headache
- minor aches & pains
- fever
- runny nose and sneezing

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses (30 mL each) in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy Alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland

**Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

**When using this product**

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

---

adults & children 12 yrs & over  
children 4 to under 12 yrs  
children under 4 yrs

---

30 mL every 6 hrs  
ask a doctor  
do not use

### **Other information**

- **each 30 mL contains:** sodium 28.23 mg
- store at no greater than 25°C and do not refrigerate

### **Inactive ingredients**

citric acid, FD&C blue no. 1, FD&C red no. 40, flavor, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum

### **Questions?**

**1-800-362-1683**

**DIST. BY: PROCTER & GAMBLE,  
CINCINNATI, OH 45202**

**DayQuil HIGH BLOOD PRESSURE Cold & Flu**

### ***Drug Facts***

#### **Active ingredients (in each 30 mL)**

Acetaminophen 650 mg  
Dextromethorphan HBr 20 mg

#### **Purpose**

Pain reliever/Fever reducer  
Cough suppressant

#### **Uses**

temporarily relieves common cold/flu symptoms:

- cough due to minor throat & bronchial irritation
- sore throat
- headache
- minor aches & pains
- fever

## Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses (30 mL each) in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy Alert:** Acetaminophen may cause severe skin reactions.

Symptoms may include: • skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

## Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

## Ask a doctor before use if you have

- liver disease
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

## Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

## Stop use and ask a doctor if

- pain or cough get worse or last more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts.

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

## Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not

notice any signs or symptoms.

### Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

---

adults & children 12 yrs & over  
children 4 to under 12 yrs  
children under 4 yrs

---

30 mL every 4 hrs  
ask a doctor  
do not use

---

### Other information

- **each 30 mL contains:** sodium 26.06 mg
- store at no greater than 25°C and do not refrigerate

### Inactive ingredients

citric acid, FD&C yellow no. 6, flavor, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum

### Questions?

**1-800-362-1683**

**TAMPER EVIDENT: Do not use if printed shrinkband seal around the neck is broken or missing**

**DIST. BY: PROCTER & GAMBLE,  
CINCINNATI, OH 45202**

### PRINCIPAL DISPLAY PANEL - Kit Carton

FAST RELIEF

DAY & NIGHT PACK

Vicks DayQuil and NyQuil HIGH BLOOD PRESSURE

Vicks DayQuil HIGH BLOOD PRESSURE

COLD & FLU

**Acetaminophen, Dextromethorphan HBr**

Headache, Fever, Sore Throat, Minor Aches & Pains

Cough

DECONGESTANT FREE

SPECIALLY FORMULATED

HBP

Vicks NyQuil HIGH BLOOD PRESSURE

COLD & FLU

Acetaminophen, Doxylamine Succinate, Dextromethorphan HBr

Headache, Fever, Sore Throat, Minor Aches & Pains

Sneezing, Runny Nose

Cough

DECONGESTANT FREE

2 BOTTLES - 1 DAYQUIL/1 NYQUIL; 8 FL OZ (236 mL) EACH; TOTAL 16 FL OZ (472 mL)

**DayQuil** **NyQuil**

**SUGAR FREE** **ALCOHOL FREE** **VALUE PACK** **SUGAR FREE**

**DayQuil** **NyQuil**

**HIGH BLOOD PRESSURE COLD & FLU**

**SPECIALLY FORMULATED**

Effective cold & flu relief for people with **HIGH BLOOD PRESSURE**.

**SUGAR, ALCOHOL, & DECONGESTANT FREE**

**HBP**

TAMPER EVIDENT: Do not use if printed shrinkband seal around the neck is broken or missing

2 BOTTLES - 1 DAYQUIL/1 NYQUIL; 8 FL. OZ (236 mL) EACH; TOTAL 16 FL. OZ (472 mL)

Headache, Fever, Sore Throat, Minor Aches & Pains  
Cough  
Sneezing, Runny Nose

PARENTS: Learn about teen medicine abuse [www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

P&G Patents: [www.pg.com/patents](http://www.pg.com/patents)

21416290  
439000451011

**NyQuil™ HBP Cold & Flu**

**Drug Facts**

**Active ingredients (in each 30 mL)**

Active ingredients (in each 30 mL)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 30 mg	Cough suppressant
Doxylamine succinate 12 mg	Sedative

**Uses** temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4 doses (30 mL each) in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy Alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: skin redness, blisters, rash. If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have:**

- liver disease
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema
- trouble urinating due to enlarged prostate gland

**Ask a doctor or pharmacist before use if you are:**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

**When using this product:**

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if:**

- pain or cough get worse or last more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults & for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs
- adults & children 12 yrs & over: 30 mL per dose every 6 hrs
- children 4 to under 12 yrs: ask a doctor
- children under 4 yrs: do not use

**Other information**

- each 30 mL contains: sodium 28.23 mg
- store at no greater than 25°C and do not refrigerate

**Inactive ingredients:** citric acid, FD&C blue no. 1, FD&C red no. 40, flavor, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucrose, water, xanthan gum

**Questions? 1-800-362-1683**

**DayQuil™ HBP Cold & Flu**

**Drug Facts**

**Active ingredients (in each 30mL)**

Active ingredients (in each 30mL)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 30 mg	Cough suppressant
Doxylamine succinate 12 mg	Sedative

**Uses** temporarily relieves common cold/flu symptoms:

- cough due to minor throat & bronchial irritation
- sore throat
- headache
- minor aches & pains
- fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4 doses (30 mL each) in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy Alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: skin redness, blisters, rash. If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have:**

- liver disease
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.**

**Stop use and ask a doctor if:**

- pain or cough get worse or last more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults & for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed
- only use dose cup provided
- do not exceed 4 doses per 24 hrs
- adults & children 12 yrs & over: 30 mL per dose every 4 hrs
- children 4 to under 12 yrs: ask a doctor
- children under 4 yrs: do not use

**Other information**

- each 30 mL dose contains: sodium 29 mg
- store at no greater than 25°C and do not refrigerate

**Inactive ingredients:** citric acid, FD&C yellow no. 6, flavor, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucrose, water, xanthan gum

**Questions? 1-800-362-1683**

**DayQuil**

**HIGH BLOOD PRESSURE COLD & FLU**

**SPECIALLY FORMULATED**

Effective cold & flu relief for people with **HIGH BLOOD PRESSURE**.

**SUGAR, ALCOHOL, & DECONGESTANT FREE**

TAMPER EVIDENT: Do not use if printed shrinkband seal around the neck is broken or missing

ALSO SOLD AS DAYQUIL/NYQUIL FOR PEOPLE WITH DIABETES

**DayQuil**

**HIGH BLOOD PRESSURE COLD & FLU**

Acetaminophen, Doxylamine Succinate, Dextromethorphan HBr

**NyQuil**

**HIGH BLOOD PRESSURE COLD & FLU**

Acetaminophen, Doxylamine Succinate, Dextromethorphan HBr

**NyQuil**

**HIGH BLOOD PRESSURE COLD & FLU**

**POWERFUL RELIEF**

**VICKS®** has been trusted for over **125 YEARS** to treat your cold & flu symptoms.

**SUGAR, ALCOHOL & DECONGESTANT FREE**

PARENTS: Learn about teen medicine abuse [www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

P&G Patents: [www.pg.com/patents](http://www.pg.com/patents)

**VICKS DAYQUIL AND VICKS NYQUIL HIGH BLOOD PRESSURE COLD AND FLU**  
acetaminophen, dextromethorphan hydrobromide, doxylamine succinate kit

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:69423-974
---------------------	----------------	---------------------------	---------------

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69423-974-16	1 in 1 PACKAGE; Type 0: Not a Combination Product	07/12/2022	
2	NDC:69423-974-12	1 in 1 PACKAGE; Type 0: Not a Combination Product	07/09/2024	

**Quantity of Parts**

Part #	Package Quantity	Total Product Quantity
Part 1	1 BOTTLE, PLASTIC	236 mL
Part 2	1 BOTTLE, PLASTIC	236 mL

**Part 1 of 2****VICKS DAYQUIL HIGH BLOOD PRESSURE COLD AND FLU**

acetaminophen, dextromethorphan liquid

**Product Information**

<b>Item Code (Source)</b>	NDC:69423-972
---------------------------	---------------

<b>Route of Administration</b>	ORAL
--------------------------------	------

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 30 mL

**Inactive Ingredients**

Ingredient Name	Strength
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	

<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM CITRATE</b> (UNII: 1Q73Q2JULR)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	

### Product Characteristics

<b>Color</b>	orange	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	CITRUS	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69423-972-08	236 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:69423-972-06	177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/12/2022	

## Part 2 of 2

### VICKS NYQUIL HIGH BLOOD PRESSURE COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, and doxylamine succinate liquid

### Product Information

<b>Item Code (Source)</b>	NDC:69423-973
<b>Route of Administration</b>	ORAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM CITRATE</b> (UNII: 1Q73Q2JULR)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	

## Product Characteristics

<b>Color</b>	red	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	CHERRY	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69423-973-08	236 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:69423-973-06	177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/12/2022	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/12/2022	

**Labeler** - The Procter & Gamble Manufacturing Company (004238200)