

ASPIRIN EXTRA STRENGTH- aspirin tablet, coated
Time-Cap Labs, Inc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

382R ASA 500 MG

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have lost a lot of fluid due to vomiting or diarrhea
- you have not been drinking fluids

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout, or arthritis
- taking any other drug
- under a doctor's care for any serious condition

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding
- feel faint
- have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear
- ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- drink a full glass of water with each dose
 - adults and children 12 years and over: take 1 to 2 tablets every 4 hours while symptoms last. Do not take more than 12 tablets in 24 hours unless directed by a doctor
 - children under 12 years: consult a doctor
-

-

Other information

- store at 25° C (77° F) excursions permitted between 15°-30° C (59°-86° F)
 - use by expiration date on package
-

-

corn starch, croscarmellose, hypromellose, microcrystalline cellulose, mineral oil, titanium dioxide

Use for the temporary relief of minor aches and pains due to:

headache, colds, muscle pain, menstrual pain, toothache, minor pain of arthritis or as directed by your doctor

Pain Reliever/fever reducer

Warnings:

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include: hives, facial swelling, shock, asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you are age 60 or older, have had stomach ulcers or bleeding problems, take a blood thinning (anticoagulant) or steroid drug, take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen or others), have 3 or more alcoholic drinks every day while using this product, take more or for a longer time than directed.

In each Tablet Aspirin 500 mg (NSAID*) *non-steroidal anti-inflammatory drug

Adhesive Area

Drug Facts (continued)

(aspirin, ibuprofen, naproxen or others)
 ■ have 3 or more alcoholic drinks every day while using this product. ■ take more or for a longer time than directed

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you are taking a diuretic ■ you have asthma ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
 ■ you have lost a lot of fluid due to vomiting or diarrhea
 ■ you have not been drinking fluids

Ask a doctor or pharmacist before use if you are
 ■ taking a prescription drug for diabetes, gout, or arthritis
 ■ under a doctor's care for any serious condition

Stop use and ask a doctor if
 ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood
 ■ have bloody or black stools
 ■ have stomach pain that does not get better
 ■ pain gets worse or lasts more than 10 days
 ■ redness or swelling is present in the painful area
 ■ fever gets worse or lasts more than 3 days
 ■ any new symptoms appear
 ■ ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless explicitly directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ drink a full glass of water with each dose
 ■ adults and children 12 years and over: take 1 to 2 tablets every 4 to 6 hours, while symptoms last. Do not exceed 8 tablets in 24 hours
 ■ children under 12 years: consult a doctor

Other information
 ■ each tablet contains: sodium 1 mg (VERY LOW SODIUM)
 ■ store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F) ■ use by expiration date on package

Inactive ingredients corn starch, croscarmellose sodium, hypromellose, microcrystalline cellulose, mineral oil, titanium dioxide

STOP PEELING

Time-Cap Labs, Inc. NDC 49483-382-01
 'Compare to the active ingredient in Bayer® Extra Strength

Extra Strength ASPIRIN ASPIRIN REGIMEN
 Safety Coated

500mg
 Pain Reliever (NSAID)
 Fever Reducer

100 COATED TABLETS New Improved Formula

This product is not manufactured or distributed by Bayer Healthcare LLC, owner of the registered trademark Bayer, Extra Strength, 382R, 1112. Manufactured by Time-Cap Labs, Inc., 7 Michael Avenue, Farmingdale, NY 11735

Drug Facts
Active ingredient (in each tablet)
 Aspirin 500 mg (NSAID)* Pain reliever/fever reducer
 *nonsteroidal anti-inflammatory drug

Uses temporarily relieves ■ headache ■ muscle pain
 ■ pain and fever of colds ■ menstrual pain
 ■ minor pain of arthritis ■ toothache

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Aspirin may cause a severe allergic reaction which may include ■ hives ■ shock ■ facial swelling ■ asthma (wheezing)
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs

LOT #: _____
EXP. DATE: _____

DO NOT PEEL HERE TO REMOVE DRUG FACTS

ASPIRIN EXTRA STRENGTH

aspirin tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49483-382
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	500 mg in 500

Inactive Ingredients

Ingredient Name	Strength
-----------------	----------

MODIFIED CORN STARCH (1-OCTENYL SUCCINIC ANHYDRIDE) (UNII: 461P5CJN6T)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
MINERAL OIL (UNII: T5L8T28FGP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics			
Color	white	Score	no score
Shape	ROUND	Size	12mm
Flavor		Imprint Code	TCL382
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49483-382-01	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/17/2018	
2	NDC:49483-382-00	100000 in 1 CARTON; Type 0: Not a Combination Product	12/17/2018	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part343	02/11/2013	

Labeler - Time-Cap Labs, Inc (037052099)

Establishment			
Name	Address	ID/FEI	Business Operations
Time-Cap Labs, Inc		037052099	manufacture(49483-382)