TUSSIN DM- dextromethorphan hydrobromide, guaifenes in solution H E B

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

HEB Tussin Drug Facts

Active ingredients (in each 10 mL)

Dextromethorphan HBr, USP 20 mg Guaifenesin, USP 200 mg

Purposes

Cough suppressant Expectorant

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Stop use and ask a doctor if

cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age

age	dose	
adults and children	10 mL every 4 hours	
12 years and over	_	
children under 12 years	do not use	

Other information

- each 10 mL contains: sodium 6 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

anhydrous citric acid, FD&C red no. 40, flavor, glycerin, high fructose corn syrup, menthol, propylene glycol, purified water, sodium benzoate, sodium citrate, sucralose

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to Robitussin® Cough + Chest Congestion DM active ingredients

Tussin

Dextromethorphan HBr/Cough Suppressant

Guaifenesin/Expectorant

Cough & Chest Congestion DM

Non-Drowsy

Adults/For Ages 12 & Over

Relief of:

Cough

Mucus

Chest Congestion

8 FL OZ (237 mL)



Drug Facts

Active ingredients Purposes

Congh & Chest Congestion Div Guaifenesin/Expectorant Condh Suppressant Dextro methor phan HBr/

(in each 10 mL)
Dextromethorphan HBr,
USP 20 mg......Cou
Guaifenesin, USP 200 mg.... .Cough suppressantExpectorant

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- cough that occurs with too much phleam (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Stop use and ask a doctor if cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

If pregnant or breast-feeding, ask a health

professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

LOT NO. EXP. # 35934 37 C32

- do not take more than 6 doses in any 24-hour period
 measure only with dosing cup provided
 keep dosing cup with product

Drug Facts (continued)

- mL = mill iliter
 this adult product is not intended for use in children under 12 years of age

age	dose	
adults and children 12 years and over	10 mL every 4 hours	
children under 12 years	do not use	

Other information

- each 10 mL contains: sodium 6 mg store at 20-25°C (68-77°F). Do not refrigerate.
- Inactive ingredients anhydrous citric acid, FD&C red no. 40, flavor, glycerin, high fructose corn syrup, menthol, propylene glycol, purified water, sodium benzoate, sodium citrate, sucralose

Questions or comments?

*This product is not manufactured or distributed by Pfizer, distributor of Robitussin® Cough + Chest Congestion DM.

GLUTEN FREE

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING

MADE WITH PRIDE & CARE FOR H-E-B® SAN ANTONIO, TX 78204





TUSSIN DM

dextromethorphan hydrobromide, guaifenesin solution

Product Information HUMAN OTC DRUG Product Type Item Code (Source) NDC:37808-359 ORAL **Route of Administration**

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9 D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 10 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 10 mL

Inactive Ingredients		
Strength		

Product Characteristics			
Color	RED (Orange-Red)	Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

P	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37808-359-26	1 in 1 CARTON	12/13/1998	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:37808-359-34	1 in 1 CARTON	08/09/1991	
2		237 mL in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:37808-359-40	1 in 1 CARTON	09/20/2002	
3		355 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	08/09/1991	

Labeler - HEB (007924756)

Revised: 5/2018 HE B