GOOD NEIGHBOR PHARMACY LOPERAMIDE HYDROCHLORIDE- loperamide hcl suspension Amerisource Bergen

Amerisource Bergen Loperamide Hydrochloride Oral Suspension Drug Facts

Active ingredient (in each 7.5 mL)

Loperamide HCl 1 mg

Purpose

Anti-diarrheal

Use

controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Do not use

if you have bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease

Ask a doctor or pharmacist before use if you are

taking antibiotics

When using this product

• tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging.

These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- shake well before using
- only use attached measuring cup to dose product

adults and children 12 years and over	30 mL (6 tsp) after the first loose stool; 15 mL (3 tsp) after each subsequent loose stool; but no more than 60 mL (12 tsp) in 24 hours
children 9-11 years (60-95 lbs)	15 mL (3 tsp) after the first loose stool; 7.5 mL (1 1/2 tsp) after each subsequent loose stool; but no more than 45 mL (9 tsp) in 24 hours
children 6-8 years (48-59 lbs)	15 mL (3 tsp) after the first loose stool; 7.5 mL (1 1/2 tsp) after each subsequent loose stool; but no more than 30 mL (6 tsp) in 24 hours
children under 6 years (up to 47 lbs)	ask a doctor

Other information

- each 30 mL (6 tsp) contains: sodium 15 mg
- store between 20-25°C (68-77°F)
- see side panel for lot number and expiration date

Inactive ingredients


anhydrous citric acid, carboxymethylcellulose sodium, D&C yellow no. 10, FD&C blue no. 1, glycerin, microcrystalline cellulose, natural and artificial mint flavor, propylene glycol, purified water, simethicone, sodium benzoate, sucralose, titanium dioxide, xanthan gum

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

Compare to Imodium® A-D active ingredient Loperamide Hydrochloride Oral Suspension anti-diarrheal Controls the Symptoms of Diarrhea mint flavor 1 mg loperamide hydrochloride per 7.5 mL



<i>Drug Facts</i> (continued)	Drug Facts (continued)		
Ask a doctor before use if you have ■ fever ■ mucus in the stool ■ a history of liver disease		ht away. (1-800-222-1222)	
Ask a doctor or pharmacist before use if you are taking antibiotics	Directions ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea ■ find right dose on chart. If possible, use weight		
When using this product matter tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.	to dose; otherwise	e use age. ■ shake well only use attached measuring	
Stop use and ask a doctor if ■ symptoms get worse ■ diarrhea lasts for more than 2 days ■ you get abdominal swelling or bulging. These may be signs of a serious condition.		30 mL (6 tsp) after the first loose stool; 15 mL (3 tsp) after each subsequent loose stool; but no more than 60 mL (12 tsp) in 24 hours	
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison	children 9-11 years (60-95 lbs)	15 mL (3 tsp) after the first loose stool; 7.5 mL (1 1/2 tsp) after each subsequent loose stool; but no more than 45 mL (9 tsp) in 24 hours	

	JUUU		$\left \right\rangle$
children 6-8 years15 mL (3 tsp) after the first loose stool; 7.5 mL (1 1/2 tsp)		Drug Facts (continued) ■ see side panel for lot number and expiration date	
	after each subsequent loose stool; but no more than 30 mL (6 tsp) in 24 hours	Inactive ingredients anhydrous citric acid, carboxymethylcellulose sodium, D&C yellow no. 10, FD&C blue no. 1, glycerin,	
children under 6 years (up to 47 lbs)	ask a doctor	microcrystalline cellulose, natural and artificial mint flavor, propylene glycol, purified water, simethicone, sodium benzoate, sucralose,	
Other information ■ each 30 mL (6 tsp) contains: sodium 15 mg ■ store between 20-25°C (68-77°F)		titanium dioxide, xanthan gum <i>Questions or comments?</i> 1-800-719-9260	

operamide hcl suspension						
Product Information						
Product T ype	HUMAN OTC DRUG	Item Code (Source)	NDC:46	DC:46122-111	
Route of Administration	ORAL					
Active Ingredient/Active Mo	piety					
Ing	redient Name		Basis of Strength		Strengtl	
L OPERAMIDE HYDROCHLORIDE (UNII:6 X9 OC 3H4II)	UNII: 77TI35393C) (LOPERAMIDE -		LOPERAMIDE HYDROCHLORIDE		1 mg in 7.5 mL	
Inactive Ingredients	Ingradiant Name				Strongth	
ANHYDROUS CITRIC ACID (UNII: X	Ingredient Name				Strength	
CARBOXYMETHYLCELLULOSE S						
D&C YELLOW NO. 10 (UNII: 35SW5	, , ,					
FD&C BLUE NO. 1 (UNII: H3R47K3T						
GLYCERIN (UNII: PDC6A3C0OX)						
CELLULOSE, MICROCRYSTALLIN	IE (UNII: OP1R32D61U)					
PROPYLENE GLYCOL (UNII: 6DC9	Q167V3)					
WATER (UNII: 059QF0KO0R)						
SODIUM BENZOATE (UNII: OJ245F)	E5EU)					
SUCRALOSE (UNII: 96K6UQ3ZD4)						
TITANIUM DIO XIDE (UNII: 15FIX9V)	210)					

Color	GREEN (opaque, viscous)	Score		
Shape		Size		
Flavor	MINT	Imprint Code		
Contains				
Packaging				
# Item Code	Package Description	Marketing Start Date	Marketing End Date	
1 NDC:46122-111-26	120 mL in 1 BOTTLE; Type 0: Not a Combination Product	02/17/2012		
Marketing Inf	ormation			
Marketing Categor	y Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA091292	02/17/2012		

Labeler - Amerisource Bergen (007914906)

Revised: 7/2018

Amerisource Bergen