

**ROFENAID 40- sulfadimethoxine and ormetoprim powder**  
**Zoetis Inc.**

-----

**RofenAid 40**

**DESCRIPTION**

RofenAid  
40

Sulfadimethoxine & Ormetoprim

Type A Medicated Article

Antibacterial

Anticoccidial

For use in broiler, replacement (breeder and layer) chickens, turkeys, ducks and chukar partridges only.

**Active Drug Ingredients**

Sulfadimethoxine . . . . . 113.5 grams per pound (25%),

Ormetoprim . . . . . 68.1 grams per pound (15%)

in a carrier suitable for incorporation in feed.

**For use in manufacturing medicated feeds only**

**Chickens**

Broiler and replacement (breeder and layer) chickens

**Indications:**

As an aid in the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti* and *E. mivati*; as an aid in the prevention of bacterial infections caused by *Haemophilus gallinarum* (infectious coryza), *Escherichia coli* (colibacillosis) and *Pasteurella multocida* (fowl cholera).

**Use directions:**

Add 1 lb of RofenAid 40 to one ton of feed to provide 0.02% (113.5 g/ton [0.0125%] sulfadimethoxine and 68.1 g/ton [0.0075%] ormetoprim) concentration in feed. Feed continuously as the sole ration.

**Warning:**

Do not feed to chickens over 16 weeks (112 days) of age.

**Turkeys**

**Indications:**

As an aid in the prevention of coccidiosis caused by *Eimeria adenoides*, *E. gallopavonis*, and *E. meleagrimitis*; for the prevention of bacterial infections caused by *Pasteurella multocida* (fowl cholera).

**Use directions:**

Add 1/2 lb of RofenAid 40 to one ton of feed to provide 0.01% (56.75 g/ton [0.00625%] sulfadimethoxine and 34.05 g/ton [0.00375%] ormetoprim) concentration in feed. Feed continuously as

the sole ration.

## **Ducks**

### **Ducks, including breeding ducks**

#### **Indications:**

As an aid in the control of bacterial infections due to *Pasteurella multocida* (fowl cholera) in ducks including breeding ducks.

#### **Use directions:**

Add 2 lbs of RofenAid 40 to one ton of feed to provide 0.04% (227 g/ton [0.025%] sulfadimethoxine and 136.2 g/ton [0.015%] ormetoprim) concentration in feed. Medication should be started at the first signs of infection. Feed for 7 days as the sole ration.

## **Ducks**

#### **Indications:**

As an aid in the control of bacterial infections due to *Escherichia coli*, *Riemerella anatipestifer* (strains 2, 3 & 4) and severe challenge of *P. multocida* (fowl cholera).

#### **Use directions:**

Add 4 lbs of RofenAid 40 to one ton of feed to provide 0.08% (454 g/ton [0.050%] sulfadimethoxine and 272.4 g/ton [0.030%] ormetoprim) concentration in feed. Medication should be started at the first signs of infection. Feed for 7 days as the sole ration.

#### **Caution:**

Not for breeding ducks. The safety of feeding sulfadimethoxine and ormetoprim at levels higher than 0.025% and 0.015%, respectively, in breeding ducks has not been established.

## **Chukar Partridges**

#### **Indications:**

For the prevention of coccidiosis caused by *Eimeria kofoidi* and *E. legionensis*.

#### **Use directions:**

Add 1 lb of RofenAid 40 to one ton of feed to provide 0.02% (113.5 g/ton [0.0125%] sulfadimethoxine and 68.1 g/ton [0.0075%] ormetoprim) concentration in feed. Feed continuously as the sole ration to young birds up to 8 weeks of age.

#### **Caution:**

The safety of feeding RofenAid 40 to breeding stock has not been established.

## **Warning**

**Do not feed to turkeys and ducks producing eggs for food. Withdraw 5 days prior to slaughter of broiler and replacement chickens, turkeys and ducks.**

## **Note**

Manufacture of Type B or C feeds from this product requires a Medicated Feed Mill License.

Labels for feeds containing RofenAid 40 must contain appropriate indications, limitations and warnings as well as required feed ingredient information.

Restricted Drug (California) - Use only as directed. Not for human use.  
Approved by FDA under NADA # 040-209  
Store below 25°C (77°F), excursions permitted to 37°C (99°F).


zoetis

Distributed by:  
Zoetis Inc.  
Kalamazoo, MI 49007

Product of China

10010602

#### PRINCIPAL DISPLAY PANEL



**ROFENAID 40**

sulfadimethoxine and ormetoprim powder

### Product Information

<b>Product Type</b>	VFD TYPE A MEDICATED ARTICLE ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:54771-1446
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SULFADIMETHOXINE (UNII: 30CPC5LDEX) (SULFADIMETHOXINE - UNII:30CPC5LDEX)	SULFADIMETHOXINE	113.5 g in 0.45 kg
ORMETOPRIM (UNII: M3EFS94984) (ORMETOPRIM - UNII:M3EFS94984)	ORMETOPRIM	68.1 g in 0.45 kg

### Product Characteristics

<b>Color</b>	BROWN (beige to light tan)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54771-1446-0	22.68 kg in 1 BAG		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA040209	12/01/2018	

**Labeler** - Zoetis Inc. (828851555)

### Establishment

Name	Address	ID/FEI	Business Operations
Zoetis LLC		069370112	MANUFACTURE

Revised: 4/2020

Zoetis Inc.