

PANACUR- fenbendazole paste
Merck Sharp & Dohme Corp.


panacur®
(fenbendazole)

Paste 10% (100 mg/g) Equine Dewormer

DESCRIPTION:

Panacur® (fenbendazole) Paste 10% contains the active anthelmintic, fenbendazole. The chemical name of fenbendazole is methyl 5-(phenylthio)-2-benzimidazole carbamate.

The chemical structure is:



Each gram of Panacur® Paste 10% contains 100 mg of fenbendazole and is flavored with artificial apple-cinnamon liquid.

ACTIONS:

The antiparasitic action of Panacur® Paste 10% is believed to be due to the inhibition of energy metabolism in the parasite.

INDICATIONS:

Panacur® Paste 10% is indicated for the treatment and control of large strongyles (*Strongylus edentatus*, *S. equinus*, *S. vulgaris*), encysted early 3rd stage (hypobiotic), late 3rd stage and 4th stage cyathostome larvae, small strongyles, pinworms (*Oxyuris equi*), ascarids (*Parascaris equorum*), and for the control of arteritis caused by 4th stage larvae of *Strongylus vulgaris* in horses.

PRECAUTIONS:

Side effects associated with Panacur® Paste 10% could not be established in well-

controlled safety studies in horses with single doses as high as 454 mg/lb (1,000 mg/kg) and 15 consecutive daily doses of 22.7 mg/lb (50 mg/kg). Particularly with higher doses, the lethal action of fenbendazole may cause the release of antigens by the dying parasites. This phenomenon may result in either a local or systemic hypersensitivity reaction. As with any drug, these reactions should be treated symptomatically.

Panacur® Paste 10% has been evaluated for safety in pregnant mares during all stages of gestation with doses as high as 11.4 mg/lb (25 mg/kg) and in stallions with doses as high as 11.4 mg/lb (25 mg/kg). No adverse effects on reproduction were detected. The recommended dose for control of 4th stage *Strongylus vulgaris* larvae, 4.6 mg/lb (10 mg/kg) daily for 5 consecutive days, has not been evaluated for safety in stallions or pregnant mares.

WARNINGS:

NOT FOR USE IN HUMANS. KEEP OUT OF REACH OF CHILDREN. The Safety Data Sheet (SDS) contains more detailed occupational safety information. For customer service, adverse effects reporting, and/or a copy of the SDS, call 1-800-211-3573. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDAVETS, or <http://www.fda.gov/reportanimalae>.

OTHER WARNINGS: Do not use in horses intended for human consumption.

Parasite resistance may develop to any dewormer, and has been reported for most classes of dewormers. Treatment with a dewormer used in conjunction with parasite management practices appropriate to the geographic area and the animal(s) to be treated may slow the development of parasite resistance. Fecal examinations or other diagnostic tests and parasite management history should be used to determine if the product is appropriate for the herd, prior to the use of any dewormer. Following the use of any dewormer, effectiveness of treatment should be monitored (for example, with the use of a fecal egg count reduction test or another appropriate method). A decrease in a drug's effectiveness over time as calculated by fecal egg count reduction tests may indicate the development of resistance to the dewormer administered. Your parasite management plan should be adjusted accordingly based on regular monitoring.

DOSAGE:

For foals and weanlings (less than 18 months of age) where ascarids are a common problem, the recommended dose is 4.6 mg/lb (10 mg/kg); one syringe will deworm a 550 lb horse.

For the control of large strongyles, small strongyles, and pinworms, the recommended dose is 2.3 mg/lb (5 mg/kg). One syringe will deworm a 1,100 lb horse.

For control of hypobiotic (encysted early 3rd stage), late 3rd stage, and 4th stage cyathostome larvae, as well as 4th stage *Strongylus vulgaris* larvae, the recommended dose is 4.6 mg/lb (10 mg/kg) daily for 5 consecutive days; administer one syringe for each 550 lb body weight per day.

Do not underdose. Ensure each animal receives a complete dose based on a current body weight. Underdosing may result in ineffective treatment, and encourage the development of parasite resistance.

DOSAGE: Panacur® Paste 10% is administered orally at a rate of 2.3 mg/lb (5 mg/kg) for the control of large strongyles, small strongyles, and pinworms. One syringe will deworm two 1,250 lb (568 kg) horses at a dose of 5 mg/kg. For foals and weanlings (less than 18 months of age) where ascarids are a common problem, the recommended dose is 4.6 mg/lb (10 mg/kg); one syringe will deworm a 1,250 lb horse.

For control of hypobiotic (encysted early 3rd stage), late 3rd stage, and 4th stage cyathostome larvae, as well as 4th stage *Strongylus vulgaris* larvae, the recommended dose is 4.6 mg/lb (10 mg/kg) daily for 5 consecutive days; administer one syringe for each 1,250 lb body weight per day.

Do not underdose. Ensure each animal receives a complete dose based on a current body weight. Underdosing may result in ineffective treatment, and encourage the development of parasite resistance.

DIRECTIONS FOR USE:

1. Determine the weight of the horse.
2. Remove syringe tip.
3. Turn the dial ring until the edge of the ring nearest the tip lines up with zero.
4. Depress plunger to advance paste to tip.
5. Set the dial ring at the graduation nearest the weight of the horse for the dosage rate of 5 mg/kg. For the dosage rate of 10 mg/kg, set the dial ring at two times (double) the horse's weight.
6. Horse's mouth should be free of food.
7. Insert nozzle of syringe through the interdental space and deposit the paste on the back of the tongue by depressing the plunger.

CONSULT YOUR VETERINARIAN FOR ASSISTANCE IN THE DIAGNOSIS, TREATMENT AND CONTROL OF PARASITISM.

HOW SUPPLIED:

Panacur® Paste 10% Equine Dewormer is supplied in 25 gram syringes.

Store at or below 25°C (77°F).

HOW SUPPLIED: Panacur® Paste 10% Equine Dewormer is supplied in 57 gram syringes, 5 per carton.

Store at or below 25°C (77°F).

Fenbendazole (active ingred.) made in: see imprint. Formulated in France.

Distributed by:

Intervet Inc. (d/b/a Merck Animal Health)
Rahway, NJ 07065

Approved by FDA under NADA #120-648

©2023 Merck & Co., Inc., Rahway, NJ, USA and its affiliates. All rights reserved.

Rev. 7/23

MERCK
Animal Health

PRINCIPAL DISPLAY PANEL - 25 gram Syringe Carton

panacur®
(fenbendazole)

Equine
Dewormer

25 gram Paste 10%
(100 mg/g)

MERCK
Animal Health

panacur®

(fenbendazole)



panacur®

(fenbendazole)

Equine
Dewormer
25 gram Paste 10%
(100 mg/g)



DOSAGE:

For foals and yearlings less than 18 months of age where a coccid is a common problem, the recommended dose is 4.4 mg/lb (10 mg/kg), one syringe will deworm 350 lb horse.

For the control of large strongyles, small strongyles, and pinworms, the recommended dose is 8.3 mg/lb (5 mg/kg). One syringe will deworm a 1,100 lb horse.

For control of hydatids (encysted early 3rd stage), tetra- or tricho-, and 4th stage cyathostome larvae, as well as 4th stage Strongylus vulgaris larvae, the recommended dose is 5.6 mg/lb (10 mg/kg daily for 5 consecutive days, administered one syringe for each 550 lb body weight per day).

Do not overdose. Ensure each animal receives a complete dose based on a current body weight.

Underdosing may result in ineffective treatment, and encourage the development of parasite resistance.

DIRECTIONS FOR USE:

1. Determine the weight of the horse.
2. Remove syringe tip.
3. Turn the dial (mg/kg) on the side of the ring at the top then up with a coin.
4. Turn the dial (mg/kg) on the side of the ring at the top then up with a coin.
5. Insert the plunger in a nostril tube to the nostril.
6. Horse's mouth should be free of food.
7. Insert 1/2 oz of syringe through the nostril space and deposit the paste on the back of the tongue by depressing the plunger.

CONSULT YOUR VETERINARIAN
FOR ASSISTANCE IN THE
DIAGNOSIS, TREATMENT AND
CONTROL OF PARASITISM.

PRECAUTIONS:

SIDE EFFECTS: Associated with Panacur® Paste 10% may not be established in well-controlled safety studies in horses with single doses as high as 54 mg/lb (100 mg/kg) and 15 consecutive daily doses of 22.7 mg/lb (50 mg/kg). Particularly with higher doses, the interaction of fenbendazole may cause a release of sulfurous gas in either a local or systemic hypersensitivity reaction. As with any drug, these reactions should be identified promptly.

Panacur® Paste 10% has been evaluated for safety in pregnant mares during all stages of gestation with doses as high as 11.4 mg/lb (25 mg/kg) and in stallions treated with up to 1.4 mg/lb (25 mg/kg). No adverse effects in stallions were described. The recommended dose for control of 4th stage Strongyle larvae (4.4 mg/lb (10 mg/kg daily for 5 consecutive days), has not been evaluated for safety in stallions or pregnant mares.

WARNINGS: NOT FOR USE IN HUMANS. KEEP OUT OF REACH OF CHILDREN. The Safety Data Sheet (SDS) contains more detailed occupational safety information. For customer service, a adverse effects reporting, and/or a copy of the SDS, call 1-800-211-3573. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS, or

HOW SUPPLIED:

Panacur Paste 10% Equine

Dewormer is supplied in 25 gram

syringes.

Store at or below 25°C (77°F).

Fenbendazole (active ingredient) 10%.

In: Imprint. Formulated in France.

Distributed by:
Intervet Inc. (d/b/a Merck Animal
Health)
Rahway, NJ 07065

Approved by FDA under NADA #

120-948

©2023 Merck & Co., Inc., Rahway,

NJ, USA and its affiliates. All rights

reserved.

Rev. 7/23



panacur®

(fenbendazole)

Paste 10% (100 mg/g) Equine Dewormer

DESCRIPTION:

Panacur® (fenbendazole) Paste 10% contains the active anti-helmintic, fenbendazole. The chemical name is 1-(phenylmethyl)-5-(phenoxy)-2-benzimidazole carboxamide. The chemical structure is:



Each gram of Panacur® Paste 10% contains 100 mg of fenbendazole and is flavored with artificial apple-flavored oil liquid.

ACTIONS:

The anthelmintic action of Panacur® Paste 10% is mainly 5-phenoxy-2-benzimidazole carboxamide. The chemical structure is:



Panacur® Paste 10% is indicated for the treatment and control of large strongyles (*Strongylus edentatus*, *S. equinus*, *S. vulgaris*), encysted early 3rd stage (hypobiotic), late 3rd stage and 4th stage cyathostome larvae, small strongyles, pinworms (*Oxyuris equi*), ascarids (*Parascaris equorum*) and the control of *Strongylus vulgaris* in horses.

PRECAUTIONS:

SIDE EFFECTS: Associated with Panacur® Paste 10% may not be established in well-controlled safety studies in horses with single doses as high as 54 mg/lb (100 mg/kg) and 15 consecutive daily doses of 22.7 mg/lb (50 mg/kg). Particularly with higher doses, the interaction of fenbendazole may cause a release of sulfurous gas in either a local or systemic hypersensitivity reaction. As with any drug, these reactions should be identified promptly.

Panacur® Paste 10% has been evaluated for safety in pregnant mares during all stages of gestation with doses as high as 11.4 mg/lb (25 mg/kg) and in stallions treated with up to 1.4 mg/lb (25 mg/kg). No adverse effects in stallions were described. The recommended dose for control of 4th stage Strongyle larvae (4.4 mg/lb (10 mg/kg daily for 5 consecutive days), has not been evaluated for safety in stallions or pregnant mares.

WARNINGS: NOT FOR USE IN HUMANS. KEEP OUT OF REACH OF CHILDREN. The Safety Data Sheet (SDS) contains more detailed occupational safety information. For customer service, a adverse effects reporting, and/or a copy of the SDS, call 1-800-211-3573. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS, or

HOW SUPPLIED:

Panacur Paste 10% Equine

Dewormer is supplied in 25 gram

syringes.

Store at or below 25°C (77°F).

Fenbendazole (active ingredient) 10%.

In: Imprint. Formulated in France.

Distributed by:
Intervet Inc. (d/b/a Merck Animal
Health)

Rahway, NJ 07065

Approved by FDA under NADA #


120-948


©2023 Merck & Co., Inc., Rahway,

NJ, USA and its affiliates. All rights

reserved.

Rev. 7/23





PRINCIPAL DISPLAY PANEL - 57 gram Syringe Carton

panacur®
(fenbendazole)

POWERPAC

Equine Dewormer

Controls Encysted EL₃ Small Strongyle Larvae

Controls both larval & adult parasites

MERCK
Animal Health

API Model
Lot:
Exp.:

Import aera
vamish free zone



- *S. equinus*, *S. vulgaris*
- Large strongyles (*Strongylus edentatus*)
- Small strongyles (*Cyathostomes*)
- *LL₃* - encysted late 3rd stage and 4th stage mucosal cyathostome larvae
- *EL₃* - encysted (hypobiotic) early 3rd stage
- Ascariids (*Parascaris equorum*)
- Pinworms (*Oxyuris equi*)

Panacur® (fenbendazole) is indicated for the control of

panacur® POWERPAC

POWERPAC

panacur®
(fenbendazole)
POWERPAC

Equine Dewormer

Controls Encysted EL₃ Small Strongyle Larvae

Controls both larval & adult parasites



MERCK

panacur®

(fenbendazole)

Paste 10% (100 mg/g)
Equine Dewormer

DESCRIPTION:

Panacur® (Fenbendazole) Paste 10% contains the active anthelmintic fenbendazole. The chemical name of fenbendazole is methyl 5-(phenylthio)-2-benzimidazole carbamate. The chemical structure is:



Each gram of Panacur® Paste 10% contains 100 mg of fenbendazole and is flavored with artificial apple-cinnamon liquid.

ACTIONS:

The antiparasitic action of Panacur® Paste 10% is believed to be due to the inhibition of energy metabolism in the parasite.

INDICATIONS:

Panacur® Paste 10% is indicated for the treatment and control of large strongyles (*Strongylus edentatus*, *S. equinus*, *S. vulgaris*), encysted early 3rd stage (hypobiotic), late 3rd stage and 4th stage cyathostome larvae, small strongyles, pinworms (*Oxyuris equi*), sarcoids (Panacur® Equimax), and control of arteritis caused by 4th stage larvae of *Strongylus vulgaris* in horses.

PRECAUTIONS:

Side effects associated with Panacur® Paste 10% could not be established in well-controlled safety studies in horses with single doses as high as 454 mg/lb (1,000 mg/kg) and 15 consecutive daily doses of 227 mg/lb (50 mg/kg). Particularly with higher doses, the lethal action of fenbendazole may cause the release of antigens by the dying parasites. This phenomenon may result in either a local or systemic hypersensitivity reaction. As with any drug, these reactions should be treated symptomatically.

Panacur® Paste 10% has been evaluated for safety in pregnant mares during all stages of gestation with doses as high as 11.4 mg/lb (25 mg/kg) and in stallions with doses as high as 11.4 mg/lb (25 mg/kg). No adverse effects on reproduction

were detected. The recommended dose for control of 4th stage *Strongylus vulgaris* larvae, 4.6 mg/lb (10 mg/kg) daily for 5 consecutive days, has not been evaluated for safety in stallions or pregnant mares.

WARNINGS: NOT FOR USE IN HUMANS. KEEP OUT OF REACH OF CHILDREN. The Safety Data Sheet (SDS) contains more detailed occupational safety information. For customer service, adverse effects reporting, and/or a copy of the SDS, call 1-800-211-3573. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS, or <http://www.fda.gov/reportadverse>.

OTHER WARNINGS: Do not use in horses intended for human consumption.

Parasite resistance may develop to any dewormer and has been reported for most classes of dewormers. Treatment with a dewormer used in conjunction with parasite management practices appropriate to the geographic area and the animal(s) to be treated may slow the development of parasite resistance. Fecal examinations or other diagnostic tests and parasite management history should be used to determine if the product is appropriate for the herd, prior to the use of any dewormer. Following the use of any dewormer, effectiveness of treatment should be monitored (for example, with the use of a fecal egg count reduction test or another appropriate method). A decrease in a drug's effectiveness over time as calculated by fecal egg count reduction tests may indicate the development of resistance to the dewormer administered. Your parasite management plan should be adjusted accordingly based on regular monitoring.

DOSAGE: Panacur® Paste 10% is administered orally at a rate of 2.3 mg/lb (5 mg/kg) for the control of large strongyles, small strongyles, and pinworms. One syringe will deworm two 1,250 lb (568 kg) horses at a dose of 5 mg/kg. For foals and weanlings less than 18 months of age where ascards are a common problem, the recommended dose is 4.6 mg/lb (10 mg/kg); one syringe will deworm a 1,250 lb horse.

For control of hypobiotic (encysted early 3rd stage), late 3rd stage, and 4th stage cyathostome larvae, as well as 4th stage *Strongylus vulgaris* larvae, the recommended dose is 4.6 mg/lb (10 mg/kg) daily for 5 consecutive days; administer one syringe for each 1,250 lb body weight per day.

Do not underdose. Ensure each animal receives a complete dose based on a current body weight. Underdosing may result in ineffective treatment, and encourage the development of parasite resistance.

DIRECTIONS FOR USE:

1. Determine the weight of the horse.
2. Remove syringe tip.
3. Turn the dial ring until the edge of the ring nearest the tip lines up with zero.
4. Depress plunger to advance paste to tip.
5. Now set the dial ring at the graduation nearest the weight of the horse.
6. Horse's mouth should be free of food.
7. Insert nozzle of syringe through the interlabial space and deposit the paste on the back of the tongue by depressing the plunger.

CONSULT YOUR VETERINARIAN FOR ASSISTANCE IN THE DIAGNOSIS, TREATMENT AND CONTROL OF PARASITISM.

HOW SUPPLIED: Panacur® Paste 10% Equine Dewormer is supplied in 57 gram syringes, 5 per carton.

Store at or below 25°C (77°F).

Fenbendazole (active ingred.) made in: see Imprint. Formulated in France.

Distributed by:

Intervet Inc. (db/a Merck Animal Health)

Rahway, NJ 07065

Approved by FDA under NADA # 120-648

©2023 Merck & Co., Inc., Rahway, NJ, USA and its affiliates. All rights reserved.

Rev. 07/23



panacur®


(fenbendazole)

POWERPAC

Equine Dewormer

Controls both larval & adult parasites

798933 R1



PSF020F14 05 R1
124x102x248mm
CMYK

PANACUR

fenbendazole paste

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:57926-081
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FENBENDAZOLE (UNII: 621BVT9M36) (FENBENDAZOLE - UNII:621BVT9M36)	FENBENDAZOLE	100 mg in 1 g

Product Characteristics

Color		Score	
Shape		Size	
Flavor	APPLE, CINNAMON	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:57926-081-44	1 in 1 CARTON		
1		25 g in 1 SYRINGE, PLASTIC		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA120648	05/10/2010	

PANACUR

fenbendazole paste

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:57926-082
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FENBENDAZOLE (UNII: 621BVT9M36) (FENBENDAZOLE - UNII:621BVT9M36)	FENBENDAZOLE	100 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
CARBOMER HOMOPOLYMER TYPE B (ALLYL PENTAERYTHRITOL CROSSLINKED) (UNII: HHT01ZNK31)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
GLYCERIN (UNII: PDC6A3C00X)	
SORBITOL (UNII: 506T60A25R)	
WATER (UNII: 059QF0KO0R)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	

Product Characteristics

Color	Score
Shape	Size

Flavor	APPLE, CINNAMON	Imprint Code
Contains		

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:57926-082-48	1 in 1 CARTON		
1		57 g in 1 SYRINGE, PLASTIC		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA120648	07/22/2011	

Labeler - Merck Sharp & Dohme Corp. (001317601)

Revised: 11/2024

Merck Sharp & Dohme Corp.