

**GOOD SENSE MIGRAINE FORMULA- acetaminophen, aspirin, caffeine tablet, film coated
L. Perrigo Company**

Perrigo Migraine Formula Drug Facts

Active ingredients (in each caplet)

Acetaminophen 250 mg

Aspirin 250 mg (NSAID*)

Caffeine 65 mg

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever

Pain reliever aid

Use

- treats migraine

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 2 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: his product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Medication overuse headache warning: Headaches may worsen if this product is used for 10 or more days per month.

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

Do not use

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

Ask a doctor before use if

- you have never had migraines diagnosed by a health professional
- you have a headache that is different from your usual migraines
- you have the worst headache of your life
- you have fever and stiff neck
- you have headaches beginning after or caused by head injury, exertion, coughing or bending
- you experienced your first headache after the age of 50
- you have daily headaches
- you have a migraine so severe as to require bed rest
- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have vomiting with your migraine headache

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for:
 - diabetes
 - gout

- arthritis
- under a doctor's care for any serious condition
- taking any other drug
- taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - your migraine is not relieved or worsens after first dose
 - new or unexpected symptoms occur
 - ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed**
- adults: take 2 caplets with a glass of water
- if symptoms persist or worsen, ask your doctor
- do not take more than 2 caplets in 24 hours, unless directed by a doctor
- under 18 years of age: ask a doctor

Other information

- store at 20°-25°C (68°-77°F)
- close cap tightly after use
- read all product information before using. Keep this box for important information.

Inactive ingredients

carnauba wax, crospovidone, FD&C blue #1 aluminum lake, hypromellose, microcrystalline cellulose, povidone, pregelatinized starch, propylene glycol, saccharin sodium, stearic acid, titanium dioxide

Questions comments?

1-800-719-9260

Principal Display Panel

Treats Migraine

See New Warnings Information

Migraine Formula

Acetaminophen, Aspirin (NSAID) and Caffeine Tablets

Pain Reliever/Pain Reliever Aid

Compare to active ingredients of Excedrin® Migraine

100% SATISFACTION GUARANTEED

100 Coated Caplets**

**Capsule-Shaped Tablets

GOODSENSE[®]

NDC 0113-0374-78

Treats Migraine

See New Warnings Information

Migraine Formula



Acetaminophen, Aspirin (NSAID) and Caffeine Tablets
Pain Reliever/Pain Reliever Aid

Compare to active ingredients of
Excedrin[®] Migraine

100%
SATISFACTION
GUARANTEED

100 Coated Caplets** **Capsule-Shaped Tablets

37478 C2 C8



Drug Facts

Active ingredients (in each caplet)

Acetaminophen 250 mg.....Pain reliever
Aspirin 250 mg (NSAID).....Pain reliever
Caffeine 65 mg.....Pain reliever aid
*nonsteroidal anti-inflammatory drug

Use ■ treats migraine

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

■ hives ■ facial swelling ■ asthma (wheezing) ■ shock
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

■ more than 2 caplets in 24 hours, which is the maximum daily amount

■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

■ are age 60 or older
■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed

Drug Facts (continued)

Medication overuse headache warning: Headaches may worsen if this product is used for 10 or more days per month.
Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

Do not use

■ if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
■ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

Ask a doctor before use if

■ you have never had migraines diagnosed by a health professional
■ you have a headache that is different from your usual migraines
■ you have the worst headache of your life
■ you have fever and stiff neck
■ you have headaches beginning after or caused by head injury, exertion, coughing or bending
■ you experienced your first headache after the age of 50
■ you have daily headaches
■ you have a migraine so severe as to require bed rest
■ you have liver disease
■ stomach bleeding warning applies to you
■ you have a history of stomach problems, such as heartburn
■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
■ you are taking a diuretic
■ you have asthma
■ you have problems or serious side effects from taking pain relievers or fever reducers
■ you have vomiting with your migraine headaches

Ask a doctor or pharmacist before use if you are

■ taking a prescription drug for:
■ diabetes ■ gout ■ arthritis
■ under a doctor's care for any serious condition
■ taking any other drug
■ taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

Drug Facts (continued)

Stop use and ask a doctor if

■ an allergic reaction occurs. Seek medical help right away.
■ you experience any of the following signs of stomach bleeding:
■ feel faint ■ vomit blood ■ have bloody or black stools
■ have stomach pain that does not get better
■ your migraine is not relieved or worsens after first dose
■ new or unexpected symptoms occur
■ ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ do not use more than directed
■ adults: take 2 caplets with a glass of water
■ if symptoms persist or worsen, ask your doctor
■ do not take more than 2 caplets in 24 hours, unless directed by a doctor
■ under 18 years of age: ask a doctor

Other information

■ store at 20°-25°C (68°-77°F)
■ close cap tightly after use
■ read all product information before using. Keep this box for important information.

Inactive ingredients camellia wax, croscopolidone, FD&C blue #1 aluminum lake, hypromellose, microcrystalline cellulose, polydextrose, pregelatinized starch, propylene glycol, saccharin sodium, stearic acid, titanium dioxide

Questions or comments? 1-800-719-9260

KEEP BOX FOR REFERENCE

DO NOT USE IF PRINTED FOIL UNDER CAP IS BROKEN OR MISSING

Gluten Free

Distributed By
Perrigo[®]
Allergan, MI 49010

GOOD SENSE MIGRAINE FORMULA

acetaminophen, aspirin, caffeine tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0113-0374
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
CROSPVIDONE (15 MPA.S AT 5%) (UNII: 68401960MK)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	WHITE	Score	no score
Shape	OVAL	Size	17mm
Flavor		Imprint Code	L374
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0113-0374-62	1 in 1 CARTON	04/17/2002	
1		24 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:0113-0374-71	1 in 1 CARTON	04/17/2002	
2		50 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:0113-0374-78	1 in 1 CARTON	04/17/2002	
3		100 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075794	04/17/2002	

Labeler - L. Perrigo Company (006013346)

Revised: 12/2019

L. Perrigo Company