

**HEALTHY ACCENTS ASPIRIN- aspirin tablet, chewable**  
**DZA Brands LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**DZA Brands, LLC Aspirin Drug Facts**

**Active ingredient (in each tablet)**

Aspirin 81 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever

**Uses**

for the temporary relief of minor aches and pains

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

**Ask a doctor before use if**

- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug for diabetes, gout or arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
  - pain gets worse or lasts for more than 10 days
  - fever gets worse or lasts for more than 3 days
  - redness or swelling is present
  - new symptoms occur
  - ringing in the ears or loss of hearing occurs

These could be signs of a serious condition

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

**Other information**

- store at 20-25°C (68-77°F)

**Inactive ingredients**

corn starch, dextrose excipient, FD&C yellow no. 6 aluminum lake, flavors, saccharin sodium

## Questions or comments?

1-866-322-2439

## Principal Display Panel

Compare to Bayer® Chewable Aspirin active ingredient

low dose

aspirin

pain reliever (NSAID)

81 mg

low dose

aspirin regimen

orange flavor

81 mg each

actual size

**DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING**

7  
LOT NO. 25439193432  
9

EXP. : 4 5 7 5 8 HA CH

healthy accents™  
low dose  
aspirin  
pain reliever (NSAID) • 81 mg  
low dose • aspirin regimen\*\*  
orange flavor

NDC 55316-467-68  
Compare to Bayer®  
Chewable Aspirin  
active ingredient

**36 chewable tablets • 81 mg each**

actual size

**Drug Facts (continued)**  
**Inactive ingredients** corn starch, dextrose excipient, FD&C yellow no. 6 aluminum lake, flavors, saccharin sodium

**Questions or comments?**  
1-866-322-2439

Distributed by: DZA Brands, LLC  
2110 Executive Drive  
Salisbury, NC 28147

For product questions or concerns  
contact us at 1-866-322-2439.  
Please include UPC number  
and code from package.

**Other information**  
store at 20-25°C (68-77°F)

**Drug Facts (continued)**

- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or loss of hearing occurs

These could be signs of a serious condition

**if pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

**Other information**

- store at 20-25°C (68-77°F)

\*\*Aspirin is not appropriate for everyone, so be sure to talk to your doctor before you begin an aspirin regimen.

### Drug Facts

Active ingredient (in each tablet)	Purpose
Aspirin 81 mg (NSAID)*.....	Pain reliever
*nonsteroidal anti-inflammatory drug	

**Uses** for the temporary relief of minor aches and pains

### Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug



### Drug Facts (continued)

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer  
■ for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor


### Ask a doctor before use if


- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if you are taking a prescription drug for diabetes, gout or arthritis**

### Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better





## HEALTHY ACCENTS ASPIRIN

aspirin tablet, chewable

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55316-467
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

### Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	

### Product Characteristics

<b>Color</b>	ORANGE	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	8mm
<b>Flavor</b>	ORANGE	<b>Imprint Code</b>	L467
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55316-467-68	1 in 1 CARTON	02/04/2008	09/30/2020
1		36 in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	02/04/2008	09/30/2020

