

CITROMA- magnesium citrate liquid
Shopko Stores Operating Co LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredient

Magnesium citrate 1.745g per fl oz

Purpose

Saline laxative

use

for relief of occasional constipation (irregularity). Generally produces bowel movement in 1/2 to 6 hours

Warnings

for this product

Do Not use

if you are on a low salt diet

Ask A Doctor before use if you have

- kidney disease
- a magnesium or sodium restricted diet
- abdominal pain, nausea, or vomiting
- noticed a sudden change in bowel habits that persists over a period of 2 weeks
- already used a laxative for a period longer than 1 week

Ask a doctor or pharmacist

before use if you are taking any other drug. Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use

and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

if pregnant or breast-feeding

ask help professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away

Directions

- drink a full glass (8 ounces) of liquid with each dose
- may be taken as a single daily dose or in divided doses

adults and children 12 years of age and over - 6.5 to 10 fl oz in 24 hours

children 6 to under 12 years of age - 3 to 7 fl oz maximum 7 fl oz in 24 hours

children 2 to under 6 years of age - 2 to 3 fl oz

children under 2 years of age - ask a doctor

discard unused product within 24 hours of opening bottle

other information

- magnesium content 290 mg per 1 fl oz (30 mL)
- sodium content 48 mg per 1 fl oz (30 mL)
- store at temperatures between 46° and 86° F (8° and 30° C)

Inactive ingredients

citric acid, lemon oil, polyethylene glycol, purified water, sodium bicarbonate, sodium saccharin, sucrose

Adverse Reaction

Manufactured by Vi-Jon, Inc.,

One Swan Dr., Smyrna, TN 37167

Shopko Sotres Operating Co. LLC

www.shopko.com

For Consumer Questions

call 1-888-593-0593

667.001/AG

principal display panel

SHOPKO

Lemon Flavor

Magnesium

Citrate

Oral Solution

Saline Laxative

For Fast Relief of Occasional Constipation

Do not use if tamper-evident twist-off cap is missing, broken or separated from neckring

10 FL OZ (296 mL)

SHOPKO

Lemon Flavor

Magnesium Citrate

Oral Solution

Saline Laxative

For Fast Relief of Occasional Constipation

Do not use if tamper evident twist-off cap is missing, broken or separated from necking

10 FL OZ (296 mL)

Drug Facts

Active ingredient

Magnesium citrate 1.745 g per fl oz

Purpose

Saline laxative

Use for relief of occasional constipation (irregularity). Generally produces bowel movement in 1/2 to 6 hours

Warnings

Do not use if you are on a low salt diet

Ask a doctor before use if you have ■ kidney disease

■ a magnesium or sodium-restricted diet ■ abdominal pain, nausea, or vomiting

■ noticed a sudden change in bowel habits that persists over a period of 2 weeks

■ already used a laxative for a period longer than 1 week

Ask a doctor or pharmacist before use if you are taking any other drug. Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ drink a full glass (8 ounces) of liquid with each dose

■ may be taken as a single daily dose or in divided doses

adults and children 12 years of age and over 6.5 to 10 fl oz maximum 10 fl oz in 24 hours

children 6 to under 12 years of age 3 to 7 fl oz maximum 7 fl oz in 24 hours

children 2 to under 6 years of age 2 to 3 fl oz maximum 3 fl oz in 24 hours

children under 2 years of age ask a doctor

discard unused product within 24 hours of opening bottle

Other information ■ magnesium content 290 mg per 1 fl oz (30 mL) ■ sodium content 45 mg per 1 fl oz (30 mL) ■ store at temperatures between 46° and 86° F (8° and 30° C)

Inactive ingredients citric acid, lemon oil, polyethylene glycol, purified water, sodium bicarbonate, sodium saccharin, sucrose

SHOPKO
Satisfaction
Guaranteed

By Return Or Exchange

Manufactured by W-Jon, Inc.,
One Swan Dr., Smyrna, TN 37167
©2012 Shopko Stores Operating Co., LLC
www.shopko.com
For consumer questions
call 1-888-593-0593

L0003078SB

UPC 4-00014-51400-3



CITROMA

magesium citrate liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:370 12-667
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MAGNESIUM CITRATE (UNII: RHO26O1T9V) (MAGNESIUM CATION - UNII:T6V3LHY838)	MAGNESIUM CITRATE	1.745 g in 29.6 mL

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
LEMON OIL (UNII: I9GRO824LL)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
WATER (UNII: 059QF0K00R)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SUCROSE (UNII: C151H8M554)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37012-667-38	296 mL in 1 BOTTLE, GLASS; Type 0: Not a Combination Product	07/15/1989	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	07/15/1989	

Labeler - Shopko Stores Operating Co LLC (023252638)

Registrant - Vi-Jon (790752542)

Establishment

Name	Address	ID/FEI	Business Operations
Vi-Jon		790752542	manufacture(37012-667)

Revised: 5/2019

Shopko Stores Operating Co LLC