

LAXATIVE- sennosides tablet
WALGREENS

195R 0363 9978 Walgreens Sennosides USP, 25 mg Tablets

Drug Facts

Active ingredient (in each tablet)

Sennosides USP, 25 mg

Purpose

Stimulant laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in 6 to 12 hours

Warnings

Do not use laxative products when abdominal pain, nausea, or vomiting are present unless directed by a doctor

Ask a doctor before use if you have noticed a sudden change in bowel habits that persists over a period of 2 weeks.

Ask a doctor or pharmacist before use if you are taking any other drug. Laxatives may affect how other drugs work. Take this product 2 or more hours before or after other drugs.

When using this product do not use for a period longer than 1 week

Stop use and ask a doctor if rectal bleeding or failure to have a bowel movement occur after use of a laxative. These may be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- swallow tablet (s) with a glass of water.
- swallow tablet (s) whole, do not crush, break, or chew.

adults and children 12 years of age and older
children 6 to under 12 years of age
children under 6 years of age

2 tablets once or twice daily
1 tablet once or twice daily
ask a doctor

Other information

- **each tablet contains:** calcium 40 mg, sodium 2 mg (VERY LOW SODIUM)
- store between 20°-25°C (68°-77°F)

VALUE SIZE

Walgreens

NDC 0363-9978-90

Compare to the active ingredient in
ex-lax® Maximum Strength*

EASY-OPEN

Laxative Tablets

SENNOSIDES USP, 25 mg / STIMULANT LAXATIVE

Maximum Strength

90
TABLETS

NOT FOR HOUSEHOLDS
WITH YOUNG CHILDREN

ACTUAL SIZE

IMPROMPTU: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each tablet) Purpose
Sennosides USP, 25 mg..... Stimulant laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in 6 to 12 hours

Warnings

Do not use laxative products when abdominal pain, nausea, or vomiting are present unless directed by a doctor

Ask a doctor before use if you have noticed a sudden change in bowel habits that persists over a period of 2 weeks.

Ask a doctor or pharmacist before use if you are taking any other drug. Laxatives may affect how other drugs work. Take this product 2 or more hours before or after other drugs.

When using this product do not use for a period longer than 1 week

Drug Facts (continued under label)

Walgreens Pharmacist Recommended
Our pharmacists recommend the Walgreens brand.
We invite you to compare to national brands.
*This product is not manufactured or distributed by
GSK Consumer Healthcare, owner of the registered
trademark ex-lax®.

DISTRIBUTED BY:
WALGREEN CO.
2700 N. WYMAN
DEERFIELD, IL 60015
100% SATISFACTION
GUARANTEED
walgreens.com
© 2023 Walgreen Co.

W3ORG0223-F

ITEM 613502 W00000-0000-0



3 11917 00708 3

196R 0123

Lot No.:

Exp. Date:



Drug Facts (continued)

Stop use and ask a doctor if rectal bleeding or failure to have a bowel movement occur after use of a laxative. These may be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- swallow tablet(s) with a glass of water.
 - swallow tablet(s) whole, do not crush, break or chew.
- | | |
|---|-------------------------------|
| adults and children 12 years of age and older | 2 tablets once or twice daily |
| children 6 to under 12 years of age | 1 tablet once or twice daily |
| children under 6 years of age | ask a doctor |

Other information

- each tablet contains: calcium 40 mg, sodium 2 mg (VERY LOW SODIUM)
- store between 20°-25°C (68°-77°F)

Inactive ingredients acacia, calcium sulfate anhydrous, carnauba wax, colloidal silicon dioxide, corn starch, croscarmellose sodium, dibasic calcium phosphate dihydrate, FD&C blue #1 aluminum lake, iron oxide, iron oxide black, iron oxide yellow, kaolin, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, povidone, shellac, stearic acid, sugar, talc, titanium dioxide.

Questions or comments? call 1-877-290-4008



new2recycle.info

Directions ■ swallow tablet(s) with a glass of water.
 adults and children 12 years of age and older twice daily
 children 6 to under 12 years of age twice daily
 children under 6 ask a doctor

Warnings ■ Do not use laxative products for more than 7 days unless directed by a doctor. If you have a history of heart disease, kidney disease, or other medical conditions, ask your doctor before using this product. Laxatives may affect how other drugs work. Take other drugs, laxatives may affect how other drugs work. Take other drugs, laxatives may affect how other drugs work. Take other drugs, laxatives may affect how other drugs work.

Drug Facts (continued)
 Ask a doctor or pharmacist before use if you are taking any other drug. Laxatives may affect how other drugs work. Take other drugs, laxatives may affect how other drugs work. Take other drugs, laxatives may affect how other drugs work.
 this product 2 or more hours before or after other drugs. When using this product do not use for a period longer than 1 week. Stop use and ask a doctor if rectal bleeding or failure to have a bowel movement occur after use of laxative. These may be signs of a serious condition.
 If pregnant or breastfeeding, ask a health professional before use.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions ■ swallow tablet(s) with a glass of water.
 adults and children 12 years of age and older twice daily
 children 6 to under 12 years of age twice daily
 children under 6 ask a doctor

Other Information
 ■ each tablet contains: calcium 40 mg, sodium 2 mg (NERYL W/SODIUM)
 ■ store between 20° - 25° C (68° - 77° F)

Active ingredient (in each tablet) Purpose
 Sennosides USP, 25 mg Stimulant laxative

Warnings
 Do not use laxative products when abdominal pain, nausea, or vomiting are present unless directed by a doctor.

Uses ■ relieves occasional constipation (irregularly) ■ generally produces bowel movement in 6 to 12 hours

Drug Facts (continued)
 Ask a doctor or pharmacist before use if you have a condition that changes in bowel habits that persists over a period of 2 weeks.

Warnings
 Do not use laxative products when abdominal pain, nausea, or vomiting are present unless directed by a doctor.

Uses ■ relieves occasional constipation (irregularly) ■ generally produces bowel movement in 6 to 12 hours

Drug Facts (continued)
 Ask a doctor or pharmacist before use if you have a condition that changes in bowel habits that persists over a period of 2 weeks.

Warnings
 Do not use laxative products when abdominal pain, nausea, or vomiting are present unless directed by a doctor.

Uses ■ relieves occasional constipation (irregularly) ■ generally produces bowel movement in 6 to 12 hours

Drug Facts (continued)
 Ask a doctor or pharmacist before use if you have a condition that changes in bowel habits that persists over a period of 2 weeks.


Warnings
 Do not use laxative products when abdominal pain, nausea, or vomiting are present unless directed by a doctor.

Uses ■ relieves occasional constipation (irregularly) ■ generally produces bowel movement in 6 to 12 hours

Drug Facts (continued)
 Ask a doctor or pharmacist before use if you have a condition that changes in bowel habits that persists over a period of 2 weeks.

Warnings
 Do not use laxative products when abdominal pain, nausea, or vomiting are present unless directed by a doctor.

Uses ■ relieves occasional constipation (irregularly) ■ generally produces bowel movement in 6 to 12 hours



ITEM 613551 W000000-000-0
 119177100711
 3

W3ORG0223-F

Directions ■ swallow tablet(s) with a glass of water.
 adults and children 12 years of age and older twice daily
 children 6 to under 12 years of age twice daily
 children under 6 ask a doctor

Warnings ■ Do not use laxative products for more than 7 days unless directed by a doctor. If you have a history of heart disease, kidney disease, or other medical conditions, ask your doctor before using this product. Laxatives may affect how other drugs work. Take other drugs, laxatives may affect how other drugs work. Take other drugs, laxatives may affect how other drugs work.

Drug Facts (continued)
 Ask a doctor or pharmacist before use if you are taking any other drug. Laxatives may affect how other drugs work. Take other drugs, laxatives may affect how other drugs work. Take other drugs, laxatives may affect how other drugs work.
 this product 2 or more hours before or after other drugs. When using this product do not use for a period longer than 1 week. Stop use and ask a doctor if rectal bleeding or failure to have a bowel movement occur after use of laxative. These may be signs of a serious condition.
 If pregnant or breastfeeding, ask a health professional before use.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions ■ swallow tablet(s) with a glass of water.
 adults and children 12 years of age and older twice daily
 children 6 to under 12 years of age twice daily
 children under 6 ask a doctor

Other Information
 ■ each tablet contains: calcium 40 mg, sodium 2 mg (NERYL W/SODIUM)
 ■ store between 20° - 25° C (68° - 77° F)

Active ingredient (in each tablet) Purpose
 Sennosides USP, 25 mg Stimulant laxative

Warnings
 Do not use laxative products when abdominal pain, nausea, or vomiting are present unless directed by a doctor.

Uses ■ relieves occasional constipation (irregularly) ■ generally produces bowel movement in 6 to 12 hours

Drug Facts (continued)
 Ask a doctor or pharmacist before use if you have a condition that changes in bowel habits that persists over a period of 2 weeks.

Warnings
 Do not use laxative products when abdominal pain, nausea, or vomiting are present unless directed by a doctor.

Uses ■ relieves occasional constipation (irregularly) ■ generally produces bowel movement in 6 to 12 hours

Drug Facts (continued)
 Ask a doctor or pharmacist before use if you have a condition that changes in bowel habits that persists over a period of 2 weeks.

Warnings
 Do not use laxative products when abdominal pain, nausea, or vomiting are present unless directed by a doctor.

Uses ■ relieves occasional constipation (irregularly) ■ generally produces bowel movement in 6 to 12 hours

Drug Facts (continued)
 Ask a doctor or pharmacist before use if you have a condition that changes in bowel habits that persists over a period of 2 weeks.

Warnings
 Do not use laxative products when abdominal pain, nausea, or vomiting are present unless directed by a doctor.

Uses ■ relieves occasional constipation (irregularly) ■ generally produces bowel movement in 6 to 12 hours

Drug Facts (continued)
 Ask a doctor or pharmacist before use if you have a condition that changes in bowel habits that persists over a period of 2 weeks.

Warnings
 Do not use laxative products when abdominal pain, nausea, or vomiting are present unless directed by a doctor.

Uses ■ relieves occasional constipation (irregularly) ■ generally produces bowel movement in 6 to 12 hours

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

Lot No.:
 Exp Date:



Compare to the active ingredient in
 ex-lax® Maximum Strength¹

Laxative Tablets

SENNOSIDES USP, 25 mg / STIMULANT LAXATIVE

Maximum Strength

- Gentle, dependable constipation relief

24 TABLETS



ACTUAL SIZE

E2603

LAXATIVE
 sennosides tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-9978
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SENNOSIDES (UNII: 3FYP5M0IJX) (SENNOSIDES - UNII:3FYP5M0IJX)	SENNOSIDES	25 mg

Inactive Ingredients

Ingredient Name	Strength
CALCIUM PHOSPHATE, DIBASIC, DIHYDRATE (UNII: O7TSZ97GEP)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
STARCH, CORN (UNII: O8232NY3SJ)	
KAOLIN (UNII: 24H4NWX5CO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
RAW SUGAR (UNII: 8M707QY5GH)	
TALC (UNII: 7SEV7J4R1U)	
ACACIA (UNII: 5C5403N26O)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
POVIDONE (UNII: FZ989GH94E)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
CALCIUM SULFATE ANHYDROUS (UNII: E934B3V59H)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SHELLAC (UNII: 46N107B71O)	

Product Characteristics

Color	blue	Score	no score
Shape	ROUND	Size	11mm
Flavor		Imprint Code	TCL083
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-9978-48	4 in 1 CARTON	04/15/2023	

1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product	
2	NDC:0363-9978-24	2 in 1 CARTON	04/15/2023
2		12 in 1 BLISTER PACK; Type 0: Not a Combination Product	
3	NDC:0363-9978-90	90 in 1 BOTTLE; Type 0: Not a Combination Product	04/15/2023

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	04/15/2023	

Labeler - WALGREENS (008965063)

Registrant - TIME CAP LABORATORIES INC. (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
TIME CAP LABORATORIES INC.		037052099	manufacture(0363-9978)

Revised: 1/2025

WALGREENS