ACID REDUCER- ranitidine tablet Cardinal Health

Drug Facts

ACTIVE INGREDIENT (IN EACH TABLET)

Ranitidine 150 mg (as ranitidine hydrochloride USP, 168 mg)

PURPOSE

Acid reducer

USES

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

WARNINGS

Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers
- if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness**, **sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding

Ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

DIRECTIONS

- adults and children 12 years and over:
 - to **relieve** symptoms, swallow 1 tablet with a glass of water
 - to **prevent** symptoms, swallow 1 tablet with a glass of water **30 to 60 minutes before** eating food or drinking beverages that cause heartburn
 - can be used up to twice daily (do not take more than 2 tablets in 24 hours)
 - children under 12 years: ask a doctor

OTHER INFORMATION

- TAMPER EVIDENT: DO NOT USE IF THE CARTON OR INDIVIDUAL BLISTER UNIT IS OPEN OR TORN.
- store at 20° 25° C (68° 77° F)
- avoid excessive heat or humidity
- this product is sodium and sugar free

INACTIVE INGREDIENTS

Colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, talc, titanium dioxide

QUESTIONS?

Call 1-800-406-7984

PRINCIPAL DISPLAY PANEL

NDC 49781-027-24

LEADER®

Compare to Zantac 150® active ingredient†

MAXIMUM STRENGTH

Acid Reducer

Ranitidine Tablets, USP 150 mg

Acid Reducer

Prevents & Relieves Heartburn Associated with Acid Indigestion & Sour Stomach

24 TABLETS

DISTRIBUTED BY CARDINAL HEALTH

5104348/0713



ACID REDUCER

ranitidine tablet

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49781-027
Route of Administration	ORAL		

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
RANITIDINE HYDRO CHLO RIDE (UNII: BK76465IHM) (RANITIDINE - UNII:884KT10 YB7)	RANITIDINE	150 mg		

Inactive Ingredients			
Ingredient Name	Strength		
SILICON DIO XIDE (UNII: ETJ7Z6 XBU4)			
CROSCARMELLOSE SODIUM (UNII: M28 OL1HH48)			
HYPROMELLOSES (UNII: 3NXW29V3WO)			

FERRIC OXIDE RED (UNII: 1K09F3G675)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICRO CRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIO XIDE (UNII: 15FIX9 V2JP)	

Product Characteristics			
Color	pink	Score	no score
Shape	OVAL (Oval Shaped)	Size	12mm
Flavor		Imprint Code	9 R
Contains			

F	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49781-027-24	3 in 1 CARTON		
1		8 in 1 BLISTER PACK		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA200536	09/25/2013	

Labeler - Cardinal Health (097537435)

Registrant - Ranbaxy Pharmaceuticals Inc. (937890044)

Establishment				
Name	Address	ID/FEI	Business Operations	
Shasun Pharmaceuticals Limited		915786829	manufacture (49 78 1-0 27)	

Revised: 12/2013 Cardinal Health