NASAL DECONGESTANT- pseudoephedrine hcl tablet, film coated L.N.K. International, Inc.

Quality Plus 44-112

Active ingredient (in each tablet)

Pseudoephedrine HCl 30 mg

Purpose

Nasal decongestant

Uses

- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily relieves sinus congestion and pressure

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- diabetes
- heart disease
- high blood pressure
- thyroid disease
- difficulty in urination due to enlargement of the prostate gland

When using this product

do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or occur with fever

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

take 2
tablets
every
4 to 6
hours;
do
not
take
more
than 8
tablets
in 24
hours
take 1
tablet
every
4 to 6
hours;
do not
take
more
than 4
tablets
in 24
hours
do not
use
450

Other information

- each tablet contains: calcium 15 mg
- TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

croscarmellose sodium, dibasic calcium phosphate dihydrate, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, silicon dioxide, titanium dioxide, triacetin

Questions or comments?

1-800-426-9391

Principal Display Panel

QUALITY +PLUS

NDC 50844-112-46

*Compare to active ingredient in Sudafed® Sinus Congestion

MAXIMUM STRENGTH NASAL DECONGESTANT

Pseudoephedrine HCl 30 mg • NASAL DECONGESTANT

- NASAL & SINUS CONGESTION
- SINUS PRESSURE •

96 Tablets

NON-DROWSY

ACTUAL SIZE

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

*This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Sudafed® Sinus Congestion.

50844 REV0619A11246

Distributed by **LNK INTERNATIONAL, INC.** 60 Arkay Drive, Hauppauge, NY 11788 USA

Unestions or comments? 1-800-426-9391

dioxide, titanium dioxide, triacetin microcrystalline cellulose, polydextrose, polyethylene glycol, silicon yellow #6 aluminum lake, hypromellose, magnesium stearate, Inactive ingredients croscarmellose sodium, dibasic calcium phosphate dihydrate, FD&C red #40 aluminum lake, FD&C red #40 aluminum lake, TD&C red #40 aluminum satesrate

- see end flap for expiration date and lot number (4.98-69)
- store at 25°C (77°F); excursions permitted between 15°-30°C
 - OPENED OR BLISTER IS TORN OR BROKEN ■ tech tablet contains: calcium 15 mg

 TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS

Other information

children under 6 years	esu fon ob
children ages 6 to 1 1 years	take 1 tablet every 4 to 6 hours; do not take more than 4 tablets in 24 hours
adults and children 12 years and over	take 2 tablets every 4 to 6 hours; do not take more than 8 tablets in 24 hours

Nasal decongestant

help or contact a Poison Control Center (1-800-222-1222) right Keep out of reach of children. In case of overdose, get medical **Drug Facts** (continued)

If pregnant or breast-feeding, ask a health professional before

- symptoms do not improve within 7 days or occur with fever
 - nervousness, dizziness, or sleeplessness occur Stop use and ask a doctor if

When using this product do not exceed recommended dosage.

- difficulty in urination due to enlargement of the prostate gland
 - thyroid disease high blood pressure a diabetes ■ ■ heart disease
 - Ask a doctor before use if you have

etore taking this product.

prescription drug contains an MAOI, ask a doctor or pharmacist after stopping the MAOI drug. If you do not know if your or emotional conditions, or Parkinson's disease), or for 2 weeks oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric Do not use if you are now taking a prescription monoamine Warnings

- temporarily relieves sinus congestion and pressure
- femborarily relieves nasal congestion due to the common cold SƏSN

Purpose

Pseudo ephedrine HCI 30 mg. Active ingredient (in each tablet)

KEEP OUTER PACKAGE FOR

Drug Facts



MAXIMUM STRENGTH ASAL DECONGESTANT

B-1603-112-46-R REV0619A11246



NDC 50844-112-46

*Compare to active ingredient in Sudafed® Sinus Congestion

MAXIMUM STRENGTH ASAL DECONGESTANT

Pseudoephedrine HCl 30 mg • NASAL DECONGESTANT

NASAL & SINUS CONGESTION

SINUS PRESSURE



96 Tablets

NON-DROWSY

ACTUAL SIZE

977111778051110

Hauppauge, VV 11788 ASU 60 Arkay Drive Distributed by LNK INTERNATIONAL, INC.

REV0619A11246

Corporation, owner of the registered trademark Sudafed® Sinus Congestion. *This product is not manufactured or distributed by Johnson & Johnson

Quality Plus 44-112

NASAL DECONGESTANT

QUALITY

PLUS

MAXIMUM STRENGTH NA

DECONGEST

pseudoephedrine hcl tablet, film coated

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

No Print Area

Lot no & Expiration Date

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50844-112
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
	PSEUDOEPHEDRINE HYDROCHLORIDE	30 mg

Inactive Ingredients	
Ingredient Name	Strength
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
FD&C RED NO. 40 ALUMINUM LAKE (UNII: 6T47AS764T)	
FD&C YELLOW NO. 6 ALUMINUM LAKE (UNII: GYP6Z2JR6Q)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3MJQ0SDW1A)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	

Product Characteristics				
Color	red	Score	no score	
Shape	ROUND	Size	7mm	
Flavor		Imprint Code	44;112	
Contains				

Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:50844- 112-08	2 in 1 CARTON	08/25/1981		
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product			
2	NDC:50844- 112-22	4 in 1 CARTON	08/25/1981		
2		12 in 1 BLISTER PACK; Type 0: Not a Combination Product			
3	NDC:50844- 112-46	8 in 1 CARTON	08/25/1981		
3		12 in 1 BLISTER PACK; Type 0: Not a Combination Product			

Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
OTC Monograph Drug	M012	08/25/1981			

Labeler - L.N.K. International, Inc. (038154464)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(50844-112)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(50844-112) , pack(50844-112)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(50844-112)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(50844-112)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(50844-112)

Establishment					
Name	Address	ID/FEI	Business Operations		
LNK International, Inc.		117025878	manufacture(50844-112)		

Revised: 8/2025 L.N.K. International, Inc.