

INVERSEDINE- atipamezole hydrochloride injection, solution
MWI Veterinary Supply

VETONE®

Inversedine®

(atipamezole hydrochloride)

5.0 mg/mL

Sterile Injectable Solution

Dexmedetomidine and Medetomidine Reversing Agent


For intramuscular use in dogs only

CAUTION:

Federal law restricts this drug to use by or on the order of a licensed veterinarian.

DESCRIPTION:

Inversedine® (atipamezole hydrochloride) is a synthetic α_2 -adrenergic antagonist. The chemical name is 4-(2-ethyl-2,3-dihydro-1H-inden-2-yl)-1H-imidazole hydrochloride. The molecular formula is $C_{14}H_{16}N_2 \cdot HCl$ and structural formula is:



Each mL of Inversedine® contains 5.0 mg atipamezole hydrochloride, 1.0 mg methylparaben (NF), 8.5 mg sodium chloride (USP), and water for injection (USP).

INDICATIONS:

Inversedine® is indicated for the reversal of the sedative and analgesic effects of dexmedetomidine hydrochloride, and medetomidine hydrochloride in dogs.

DOSAGE AND ADMINISTRATION:

Inversedine® is administered intramuscularly (IM) for reversal of sedation and analgesia regardless of the route used for dexmedetomidine hydrochloride or medetomidine hydrochloride. The atipamezole dose for the reversal of IV dexmedetomidine

hydrochloride or medetomidine hydrochloride is 3750 mcg/m². The atipamezole dose for the reversal of IM dexmedetomidine hydrochloride or medetomidine hydrochloride is 5000 mcg/m².

The dosage of Inversedine® is calculated based on body surface area. Use the following tables to determine the correct injection volume or the correct Inversedine® dosage on the basis of kilograms of body weight.

Note that the mcg/kg dosage decreases as body weight increases.

Table 1: Atipamezole dosing for reversal of IV dexmedetomidine hydrochloride- or medetomidine hydrochloride-induced sedation/analgesia:

Dose table for Inversedine® (3750 mcg/m²) when dexmedetomidine hydrochloride or medetomidine hydrochloride is given IV

For # lb	For # kg	dose = mcg/kg	Volume = mL Inversedine®
4-7	2-3	300	0.1
7-9	3-4	250	0.15
9-11	4-5	230	0.2
11-22	5-10	200	0.3
22-33	10-15	170	0.4
33-44	15-20	150	0.5
44-55	20-25	140	0.6
55-66	25-30	130	0.7
66-81	30-37	120	0.8
81-99	37-45	110	0.9
99-110	45-50	105	1.0
110-132	50-60	100	1.1
132-143	60-65	95	1.2
143-165	65-75	93	1.3
165-176	75-80	91	1.4
>176	>80	90	1.5

Table 2: Atipamezole dosing for reversal of IM dexmedetomidine hydrochloride- or medetomidine hydrochloride- induced sedation/analgesia:

Dose table for Inversedine® (5000 mcg/m²) when dexmedetomidine hydrochloride or medetomidine hydrochloride is given IM

For # lb	For # kg	dose = mcg/kg	volume = mL Inversedine®
4-7	2-3	400	0.15
7-9	3-4	350	0.2
9-11	4-5	300	0.3
11-22	5-10	250	0.4
22-29	10-13	230	0.5
29-33	13-15	210	0.6

33-44	15-20	200	0.7
44-55	20-25	180	0.8
55-66	25-30	170	0.9
66-73	30-33	160	1.0
73-81	33-37	150	1.1
81-99	37-45	145	1.2
99-110	45-50	140	1.3
110-121	50-55	135	1.4
121-132	55-60	130	1.5
132-143	60-65	128	1.6
143-154	65-70	125	1.7
154-176	70-80	123	1.8
>176	>80	120	1.9

CONTRAINDICATIONS:

Since atipamezole is always used concomitantly with dexmedetomidine or medetomidine, it should not be used in dogs with the following conditions: cardiac disease, respiratory disorders, liver or kidney diseases, dogs in shock, severely debilitated dogs, or dogs stressed due to extreme heat, cold or fatigue.

Administration of atipamezole is contraindicated in dogs with a known hypersensitivity to the drug.

HUMAN WARNINGS:

Not for human use. Keep out of reach of children.

Atipamezole hydrochloride can be absorbed and may cause irritation following direct exposure to skin, eyes, or mouth. In case of accidental eye exposure, flush with water for 15 minutes. In case of accidental skin exposure, wash with soap and water. Remove contaminated clothing. If irritation or other adverse reaction occurs (for example, increased heart rate, tremor, muscle cramps), seek medical attention.

In case of accidental oral exposure or injection, seek medical attention. Caution should be used while handling and using filled syringes.

Users with cardiovascular disease (for example, hypertension or ischemic heart disease) should take special precautions to avoid any exposure to this product.

The safety data sheet (SDS) contains more detailed occupational safety information.

To report adverse reactions in users or to obtain a copy of the SDS for this product call 1-800-887-2763.

Note to Physician: This product contains an α_2 -adrenergic antagonist.

PRECAUTIONS:

1. Handling: Atipamezole hydrochloride can produce an abrupt reversal of sedation; therefore, dogs that have recently received atipamezole hydrochloride should be

handled with caution. The potential for apprehensive or aggressive behavior should be considered in the handling of dogs emerging from sedation, especially in dogs predisposed to nervousness or fright. Also, avoid situations where a dog might fall.

2. Sedation relapse: While atipamezole does reverse the clinical signs associated with medetomidine or dexmedetomidine sedation, complete physiologic return to pretreatment status may not be immediate or may be temporary, and dogs should be monitored for sedation relapse. Sedation relapse is more likely to occur in dogs that receive an alpha₂-agonist by the IV route, compared to dogs that are sedated using the IM route. Animals should be monitored closely for persistent hypothermia, bradycardia, and depressed respiration, until signs of recovery persist.

3. Analgesia reversal: Atipamezole reverses analgesic effects as well as sedative effects. Additional procedures for the control of pain may be required.

4. Debilitated dogs: The safety of atipamezole has not been evaluated in dogs with compromised health. Geriatric, debilitated, and ill dogs are more likely to experience adverse reactions associated with the administration of alpha₂-antagonists (as well as alpha₂-agonists). Dogs with abnormalities associated with the cardiovascular system are especially at risk.

5. Breeding dogs: Atipamezole hydrochloride has not been evaluated in breeding dogs; therefore, the drug is not recommended for use in pregnant or lactating dogs, or in dogs intended for breeding.

6. Minimum age and weight: Atipamezole hydrochloride has not been evaluated in dogs less than four months of age or in dogs weighing less than 4.4 lbs (2 kg).

ADVERSE REACTIONS:

Occasional vomiting may occur. At times, a period of excitement or apprehensiveness may be seen in dogs treated with atipamezole. Other effects of atipamezole include hypersalivation, diarrhea, and tremors.

Contact Information: To report suspected adverse drug events, for technical assistance or to obtain a copy of the Safety Data Sheet, contact Parnell at 1-800-887-2763. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

CLINICAL PHARMACOLOGY:

Atipamezole is a potent alpha₂-antagonist which selectively and competitively inhibits alpha₂-adrenergic receptors. The result of atipamezole administration in the dog is the rapid recovery from the sedative and analgesic effects produced by the alpha₂-adrenergic agonists dexmedetomidine or medetomidine. Atipamezole does not reverse the effects of other classes of sedatives, anesthetics, or analgesics.

Atipamezole is rapidly absorbed following intramuscular injection; maximum serum concentration is reached in approximately 10 minutes. Onset of arousal is usually apparent within 5 to 10 minutes of injection, depending on the depth and duration of dexmedetomidine- or medetomidine-induced sedation. Elimination half-life from serum is less than 3 hours. Atipamezole undergoes extensive hepatic biotransformation, with excretion of metabolites primarily in urine.

Dexmedetomidine or medetomidine activation of peripheral and central α_2 -adrenergic receptors induces a pattern of pharmacological responses that include sedation, reduction of anxiety, analgesia, and bradycardia.

Blood pressure is initially increased due to peripheral vasoconstriction and thereafter drops to normal or slightly below normal levels.

A transient decrease in systolic blood pressure occurs immediately after administration of atipamezole to dexmedetomidine- or medetomidine-sedated dogs, followed by a transient increase in arterial pressure within 10 minutes compared to pre-atipamezole levels. This is the opposite of the response to α_2 -agonist treatment, and is probably due to atipamezole-induced peripheral vasodilation.

Atipamezole administration rapidly abolishes dexmedetomidine- or medetomidine-induced bradycardia, usually within 3 minutes. The magnitude of the effect of atipamezole on heart rate is greater when dexmedetomidine is administered intravenously compared to intramuscularly. Dogs receiving medetomidine or IM dexmedetomidine may not return to pre-sedative heart rates after atipamezole administration and some dogs briefly show heart rate elevations above baseline. Respiratory rate increases following atipamezole injection.

EFFECTIVENESS:

One hundred and nine dogs received atipamezole in the field study (55 dogs received the reversal agent following dexmedetomidine; 54 following medetomidine). The mean age was 5.9 years and ranged between 17 weeks and 16 years. The mean weight was 45.5 lbs (20.7 kg), ranging from 4.8 lbs to 117 lbs (2.2 kg to 53.2 kg). Atipamezole was administered by the IM route of administration, within a range of 39-57 minutes after administration of either dexmedetomidine (IV and IM) or medetomidine (IV and IM).

Atipamezole reversed the effects of dexmedetomidine and medetomidine in all cases. In dexmedetomidine treated dogs, the onset of reversal was evident within 5 minutes after administration of atipamezole (57% could stand). Within 15 minutes, 96% of dexmedetomidine treated dogs were standing, 92% responded normally to sound, 86% had a normal muscle tone of jaw, and >90% had a normal pedal reflex response.

Responses in dogs treated with medetomidine were similar or slightly later.

Following atipamezole, heart rate increased between 0 and 5 minutes following either α_2 -agonist (IV dexmedetomidine dogs had heart rates from 60 to 85 bpm, and IV medetomidine dogs from 51 to 67 bpm; IM dexmedetomidine dogs had heart rates from 45 to 73 bpm, and IM medetomidine dogs from 52 to 79 bpm). Bradycardia resolved more slowly in the IM treatment groups. The body temperature remained at the same level during the 120 minutes of follow-up after atipamezole administration. Respiratory rates increased toward normal between 0 and 5 minutes after the administration of atipamezole in all treatment groups. Mucous membranes were described as normal after 5 minutes in 91% of dexmedetomidine dogs (IV or IM). By 120 minutes, 96% were normal (after IV dexmedetomidine) or 100% were normal (after IM dexmedetomidine).

Many physiological responses were slightly slower to return toward normal when dogs were treated with medetomidine IV or IM.

No adverse events were reported in the atipamezole treated dogs.

ANIMAL SAFETY:

Atipamezole was tolerated in healthy dogs receiving 10X the recommended dose and in dogs receiving repeated doses at 1, 3, and 5X the recommended dose, in the absence of an α_2 -agonist. Signs were dose-related and included excitement, panting, trembling, vomiting, soft or liquid feces and scleral injection. At 10X the recommended dose, increases in creatine kinase, AST, and ALT were noted. Creatine kinase also increased in 3 (of 6) dogs in the 3X treatment group. Localized skeletal muscle injury was seen at the injection site but no associated clinical signs or complications were observed. Dogs receiving the recommended atipamezole dose in the absence of medetomidine or dexmedetomidine exhibited no adverse clinical signs. In additional safety studies, adverse events were absent up to the 3X dose of atipamezole when its administration followed medetomidine or dexmedetomidine sedation.

In a separate safety study using a crossover design, 5 dogs received atipamezole after dexmedetomidine (IV and IM). Dexmedetomidine's effects on blood pressure, heart rate, respiratory rate, and cardiac conduction times were reversed by atipamezole. However, heart rate and cardiac conduction times did not return to predexmedetomidine values.

Heart rate increases after atipamezole were closer to baseline values in dogs treated with dexmedetomidine IV (compared to IM).

STORAGE INFORMATION:

Store at room temperature 20° to 25°C (68-77°F), with excursions permitted to 15° to 30°C (59-86°F). Use within 90 days of first puncture.

HOW SUPPLIED:

Inversedine® is supplied in 10-mL, multidose vials containing 5.0 mg of atipamezole hydrochloride per mL.

Approved by FDA under ANADA # 200-772

Manufactured by:
PARNELL TECHNOLOGIES PTY. LTD.
4/476 Gardeners Road
Alexandria NSW 2015 Australia

Distributed by: MWI
Boise, ID 83705
www.VetOne.net

10 mL: 50524-01-Jan26



Iss. 01/2026

NDC 86136-151-31

VETONE®

Inversedine® (atipamezole hydrochloride)

5.0 mg/mL

Sterile Injectable Solution


Dexmedetomidine and Medetomidine Reversing Agent

For intramuscular use in dogs only

Caution: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

Approved by FDA under ANADA # 200-772

Net Contents: 10 mL



VETONE®
 Inversedine® (atipamezole hydrochloride)

5.0 mg/mL
 Sterile Injectable Solution

Dexmedetomidine and Medetomidine Reversing Agent

For intramuscular use in dogs only

Caution: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

Approved by FDA under ANADA # 200-772

Net Contents: 10 mL


Inversedine[®]
(atipamezole hydrochloride)
5.0 mg/mL
Sterile Injectable Solution
Dexmedetomidine and Medetomidine
Reversing Agent
For intramuscular use in dogs only
Caution: Federal law restricts this drug to use by or on the order of a licensed veterinarian.
V1 503026 **Net contents: 10 mL**

NDC 86136-151-31
Storage: Store at room temperature 20° to 25°C (68-77°F), with excursions permitted to 15° to 30°C (59-86°F). Use within 90 days of first puncture.
WARNING: Keep Out of Reach of Children Not For Human Use
Read Package Insert Before Using This Drug
Approved by FDA under ANADA # 200-772
Distributed by: MWI Manufactured by:
Boise, ID 83705
www.VetOne.net
Iss. 01/2026
50522-01-Jan26

PARNELL


batch & expiry

INVERSEDINE

atipamezole hydrochloride injection, solution

Product Information

Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:86136-151
Route of Administration	INTRAMUSCULAR		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ATIPAMEZOLE HYDROCHLORIDE (UNII: 2W4279571X) (ATIPAMEZOLE - UNII:03N9U5JAF6)	ATIPAMEZOLE HYDROCHLORIDE	5 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X)	8.5 mg in 1 mL

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:86136-151-31	10 mL in 1 VIAL, GLASS		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANADA	ANADA200772	03/13/2026	

Labeler - MWI Veterinary Supply (019926120)

Revised: 3/2026

MWI Veterinary Supply