

**MAXIMUM STRENGTH SEVERE CONGESTION AND COUGH MAX-
dextromethorphan hydrobromide, guaifenesin, phenylephrine
hydrochloride liquid
Discount Drug Mart**

Drug Facts

Active ingredients (in each 20 mL)

Dextromethorphan HBr, USP 20 mg

Guaifenesin, USP 400 mg

Phenylephrine HCL, USP 10 mg

Purpose

Cough Suppressant

Expectorant

Nasal Decongestant

Keep out of reach of children.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Uses

- help loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
- cough due to minor throat and bronchial irritation
- nasal congestion

Warnings

DO NOT USE IF PRINTED SEAL UNDER CAP IS TORN OR MISSING.

Do not use

- for children under 12 years of age
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- thyroid disease

- high blood pressure
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

When using this product

- **do not use more than directed**

Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur
- symptoms do not get better within 7 days or are accompanied by a fever
- Cough comes back, or occurs with fever, rash or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Directions

- take only as recommended
- use dosage cup
- mL = milliliter
- do not take more than 6 doses in any 24-hour period

Age	Dose
Adults and children 12 years and older	20 mL every 4 hours
Children under 12 years of age	Do not use

Other information

- each 20 mL contains: sodium 10 mg
- dosage cup provided
- store between 15°-30°C (59°-86°F)
- do not refrigerate

Inactive ingredients

Anhydrous citric acid, edetate disodium, FD and C Blue # 1, FD and Red # 40, glycerin, flavors, glycerin, , propylene glycol, propyl gallate, purified water, sodium benzoate, sorbitol, sucralose, Xanthan gum

Questions?

Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944

DISCOUNT drug mart FOOD FAIR MAXIUM STRENGTH SEVERE CONGESTION & COUGH MAX product label

*COMPARE TO THE ACTIVE INGREDIENTS IN MUCINEX® FAST-MAX™ SEVERE CONGESTION AND COUGH

LF-011

DISCOUNT

drug mart

FOOD FAIR

MAXIUM STRENGTH SEVERE CONGESTION & COUGH MAX

Dextromethorphan HBr/ Guaifenesin / Phenylephrine HCL

COUGH SUPPRESSANT/

EXPECTORANT/

NASAL DECONGESTANT

Controls Cough

Thins & Loosens mucus

Relieves Nasal & Chest Congestion

Adults

For Ages 12 & Over

6 FL OZ (177 mL)

* This product is not manufactured or distributed by Reckitt Benckiser Inc., distributor of MUCINEX® FAST-MAX™ SEVERE CONGESTION AND COUGH

LB - 011 LOT: EXP:

Peel Corner to Read Complete Drug Facts and Information

SATISFACTION GUARANTEED

IF DISSASIFIED , RETURN UNUSED PORTION AND PACKAGE TO STORE WHERE PURCHASED. IF UNABLE TO RETURN TO STORE, SEND REASON FOR DISSATISFACTION, NAME, ADDRESS AND EMPTY PACKAGE TO: DISCOUNT DRUG MART, 211 COMMERCE DRIVE, MEDINA, OHIO 44256



*COMPARE TO THE
ACTIVE INGREDIENTS IN
MUCINEX® FAST-MAX™
SEVERE CONGESTION & COUGH
LF-011

**MAXIMUM STRENGTH
SEVERE CONGESTION
& COUGH MAX**

Dextromethorphan HBr /
Guaifenesin / Phenylephrine HCl

COUGH SUPPRESSANT /
EXPECTORANT /
NASAL DECONGESTANT

Controls Cough
Thins & Loosens Mucus
Relieves Nasal & Chest Congestion

Adults
For Ages 12 & Over

6 FL OZ (177 mL)

Peel Corner to Read Complete
Drug Facts and Information



DO NOT USE IF PRINTED SEAL
UNDER CAP IS TORN OR MISSING

LB-011



Lot:
Exp:

Drug Facts	
Active ingredients (in each 20 mL)	Purpose
Dextromethorphan HBr 20 mg.....	Cough suppressant
Guaifenesin 400 mg.....	Expectorant
Phenylephrine HCl 10 mg.....	Nasal Decongestant
Uses	
<ul style="list-style-type: none"> ■ helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes ■ temporarily relieves these symptoms occurring with a cold: <ul style="list-style-type: none"> ■ cough due to minor throat and bronchial irritation ■ nasal congestion 	
Warnings	
<p>Do not use ■ for children under 12 years of age</p> <ul style="list-style-type: none"> ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a 	

Drug Facts (continued)
doctor or pharmacist before taking this product.
Ask a doctor before use if you have
<ul style="list-style-type: none"> ■ heart disease ■ thyroid disease ■ high blood pressure ■ diabetes ■ trouble urinating due to an enlarged prostate gland ■ cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema. ■ cough that occurs with too much phlegm (mucus)
When using this product
<ul style="list-style-type: none"> ■ do not use more than directed
Stop use and ask a doctor if
<ul style="list-style-type: none"> ■ nervousness, dizziness or sleeplessness occur ■ symptoms do not get better within 7 days or are accompanied by a fever ■ cough comes back, or occurs with fever, rash or persistent headache. These could be signs of a serious condition.

Facts (continued)

Pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take only as recommended
- use dosage cup
- mL = milliliter
- do not take more than 6 doses in any 24-hour period

Age	Dose
Adults & children 12 years and older	20 mL every 4 hours
Children under 12 years of age	Do not use

Other information

- each 20 mL contains: sodium 10 mg
- dosage cup provided
- store between 15-30°C (59-86°F)
- do not refrigerate

Drug Facts (continued)

Inactive ingredients

anhydrous citric acid, edetate disodium, FD&C Blue #1, FD&C Red #40, flavors, glycerin, propylene glycol, propyl gallate, purified water, sodium benzoate, sorbitol, sucralose, Xanthan gum

Questions? Call weekdays from 9:30 AM to 4:30 PM EST at 1-877-798-5944

* This product is not manufactured or distributed by Reckitt Benckiser Inc., distributor of Mucinex® FAST-MAX™ Severe Congestion & Cough

SATISFACTION GUARANTEED
IF DISSATISFIED, RETURN UNUSED PORTION AND PACKAGE TO STORE WHERE PURCHASED. IF UNABLE TO RETURN TO STORE, SEND REASON FOR DISSATISFACTION, NAME, ADDRESS AND EMPTY PACKAGE TO:
DISCOUNT DRUG MART, 211 COMMERCE DRIVE, MEDINA, OHIO 44256

res

MAXIUM STRENGTH SEVERE CONGESTION AND COUGH MAX

dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53943-517
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYL GALLATE (UNII: 8D45NN7V92)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53943-517-25	177 mL in 1 BOTTLE; Type 0: Not a Combination Product	12/10/2013	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/10/2013	

Labeler - Discount Drug Mart (047741335)