

FAMOTIDINE- famotidine tablet SAFEWAY

**Maximum Strength
Acid Controller
Famotidine Tablets USP, 20mg
Acid Reducer**

**Just one Tablet!
Prevents & Relieves Heartburn due to acid indigestion**

Active Ingredient (in each tablet)

Famotidine Tablets USP, 20mg

Purpose

Acid Reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Allergy alert

Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chestpain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding,

ask a health professional before use

Keep out of reach of children

In case of overdose, get medical help or contact a poison control center right away. (1-800-222-1222)

Directions

- adults and children 12 years and over:
- to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn.
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other Information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25 C (68-77F)
- protect from moisture


Inactive Ingredients

hydroxypropyl cellulose, hypromellose, macrogol, magnesium stearate, microcrystalline cellulose, pre-gelatinized starch, sodium starch glycolate, talc, titanium dioxide, triacetin

Questions or Comments?

call **1-877-770-3183** Mon-Fri 8:00 AM EST to 5:00 PM PST

PDP



Inactive Ingredients

Ingredient Name	Strength
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
MAGNESIUM STEARATE (UNII: 70097M6I3O)	
TALC (UNII: 7SEV7J4R1U)	
HYDROXYPROPYL CELLULOSE (110000 WAMW) (UNII: 5Y0974F5PW)	
TRIACETIN (UNII: XHX3C3X673)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSE 2910 (6 MPA.S) (UNII: 0WZ8WG20P6)	
POLYETHYLENE GLYCOL 4000 (UNII: 4R4HFI6D95)	

Product Characteristics

Color	white (White to off-white)	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	V;15
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-191-20	200 in 1 BOTTLE; Type 0: Not a Combination Product	06/30/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA215822	06/30/2023	

Labeler - SAFEWAY (009137209)

Revised: 12/2025

SAFEWAY