

**FLU HBP MAXIMUM STRENGTH- acetaminophen, chlorpheniramine maleate,
dextromethorphan hbr tablet, film coated
Rite Aid Corporation**

Rite Aid 44-738

Active ingredients (in each caplet)

Acetaminophen 325 mg
Chlorpheniramine maleate 2 mg
Dextromethorphan HBr 10 mg

Purpose

Pain reliever/fever reducer
Antihistamine
Cough suppressant

Uses

- temporarily relieves these common cold and flu symptoms:
 - minor aches and pains
 - sore throat
 - sneezing
 - headache
 - runny nose
 - cough
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters
- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a

doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- difficulty in urination due to enlargement of the prostate gland
- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- avoid alcoholic beverages
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- excitability may occur, especially in children
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**
- adults and children 12 years and over: take 2 caplets every 4 hours while symptoms persist. Do not take more than 10 caplets in 24 hours.
- children under 12 years: ask a doctor

Other information

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- protect from excessive moisture
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, crospovidone, FD&C red #40 aluminum lake, lecithin, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Questions or comments?

1-800-426-9391

Principal display panel

NDC 11822-0738-8

Compare to the active ingredients in
**Coricidin® HBP Maximum Strength
Multi-Symptom Flu***

**MAXIMUM STRENGTH
FLU HBP**

ACETAMINOPHEN

PAIN RELIEVER / FEVER REDUCER

CHLORPHENIRAMINE MALEATE

ANTIHISTAMINE

DEXTROMETHORPHAN HBr

COUGH SUPPRESSANT

PAIN RELIEVER / FEVER REDUCER

ANTIHISTAMINE

COUGH SUPPRESSANT

Relieves body aches, pains & headache,
fever, cough, runny nose, sore throat, sneezing

For people with high blood

pressure

Decongestant-free

ACTUAL SIZE

24

CAPLETS

**TAMPER EVIDENT: DO NOT USE IF
PACKAGE IS OPENED OR IF BLISTER
UNIT IS TORN, BROKEN OR SHOWS
ANY SIGNS OF TAMPERING**

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Coricidin®


HBP Maximum Strength Multi-Symptom Flu.

50844

REV1019B73808

DISTRIBUTED BY: RITE AID, 200 NEWBERRY COMMONS,
ETTERS, PA 17319 www.riteaid.com

SATISFACTION GUARANTEE: If you're not satisfied,
we'll happily refund your money



Rite aid 44-738

FLU HBP MAXIMUM STRENGTH

acetaminophen, chlorpheniramine maleate, dextromethorphan hbr tablet, film coated

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|----------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:11822-0738 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------------------|----------|
| ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D) | ACETAMINOPHEN | 325 mg |
| CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U) | CHLORPHENIRAMINE MALEATE | 2 mg |
| DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 10 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|-----------------|----------|
| | |

| |
|---|
| STARCH, CORN (UNII: O8232NY3SJ) |
| CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561) |
| FD&C RED NO. 40 ALUMINUM LAKE (UNII: 6T47AS764T) |
| LECITHIN, SOYBEAN (UNII: 1DI56QDM62) |
| MAGNESIUM STEARATE (UNII: 70097M6I30) |
| MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U) |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) |
| POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990) |
| POVIDONE, UNSPECIFIED (UNII: FZ989GH94E) |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4) |
| SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2) |
| STEARIC ACID (UNII: 4ELV7Z65AP) |
| TALC (UNII: 7SEV7J4R1U) |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) |

Product Characteristics

| | | | |
|-----------------|------|---------------------|----------|
| Color | red | Score | no score |
| Shape | OVAL | Size | 16mm |
| Flavor | | Imprint Code | 44;738 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:11822-0738-8 | 2 in 1 CARTON | 07/24/2020 | |
| 1 | | 12 in 1 BLISTER PACK; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M012 | 07/24/2020 | |

Labeler - Rite Aid Corporation (014578892)

Establishment

| Name | Address | ID/FEI | Business Operations |
|-------------------------|---------|-----------|--|
| LNK International, Inc. | | 832867837 | manufacture(11822-0738) , pack(11822-0738) |

Establishment

| Name | Address | ID/FEI | Business Operations |
|-------------------------|---------|-----------|-------------------------|
| LNK International, Inc. | | 832867894 | manufacture(11822-0738) |

Establishment

| Name | Address | ID/FEI | Business Operations |
|-------------------------|---------|-----------|-------------------------|
| LNK International, Inc. | | 117025878 | manufacture(11822-0738) |

Revised: 7/2024

Rite Aid Corporation