

**WALGREEN HIGH BLOOD PRESSURE NIGHTTIME COLD AND FLU-  
acetaminophen, dextromethorphan hydrobromide, and doxylamine  
succinate liquid  
WALGREENS CO.**

-----

**Walgreens High Blood Pressure Nighttime COLD & FLU**

***Drug Facts***

<b><i>Active ingredients (in each 30 mL dose cup)</i></b>	<b><i>Purpose</i></b>
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 30 mg	Cough suppressant
Doxylamine succinate 12.5 mg	Antihistamine

**Uses**

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

**Warnings**

**Liver warning**

This product contains acetaminophen.  
Severe liver damage may occur if you take

- more than 4 doses (30 mL each dose) in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy Alert:** Acetaminophen may cause severe skin reactions.  
Symptoms may include:

- Skin reddening
- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away

### **Sore throat warning**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema
- trouble urinating due to enlarged prostate gland

### **Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

### **When using this product**

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

### **Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.**

### **Overdose warning**

In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- take only as directed
- use dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL per dose every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

### **Other information**

- **each 30 mL contains:** sodium 16 mg
- store at room temperature

### **Inactive ingredients**

Anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavor, propylene glycol, propyl gallate, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose

### **Questions?**

**1-866-467-2748**

**Distributed by**

### **PRINCIPAL DISPLAY PANEL - 236 ml Bottle Label**

Compare to the active ingredients in Vicks® NyQuil™ High Blood Pressure Cold & Flu\*

NDC 0363-6210-08

**High Blood**

**Pressure**

**Nighttime**

**Cold & Flu**

**Acetaminophen, Doxylamine Succinate, Dextromethorphan HBr**

Headache, Fever, Sore Throat,

Minor Aches & Pains

Sneezing, Runny Nose

Cough

Sugar Free & No Added Alcohol

8 FL OZ (236 mL)

\*This product is not manufactured or distributed by Procter & Gamble, the distributor of Vicks® NyQuil™ High Blood Pressure Cold & Flu.

6.75" Width

3.25" Height

NDC 0363-6210-08

**TAMPER EVIDENT: DO NOT USE IF PRINTED SHRINK BAND IS MISSING OR BROKEN**

**Drug Facts**

Active ingredients (in each 30 mL)	Purpose
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 30 mg	Cough suppressant
Doxylamine succinate 12.5mg	Antihistamine

**Uses**

- temporarily relieves common cold/flu symptoms:
  - cough due to minor throat & bronchial irritation
  - sore throat
  - minor aches & pains
  - runny nose and sneezing

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take more than 4 doses (30 mL each dose) in 24 hours, which is the maximum daily amount for this product with other drugs containing acetaminophen or 3 or more alcoholic drinks every day while using this product.

**Allergy Alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: Skin redness, blisters, rash. If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.



Compare to the active ingredients in Vicks® NyQuil™ High Blood Pressure Cold & Flu\*

**NIGHTTIME**

# Cold & Flu High Blood Pressure

ACETAMINOPHEN / PAIN RELIEVER / FEVER REDUCER  
DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT  
DOXYLAMINE SUCCINATE / ANTIHISTAMINE

**Sugar Free**

• Headache, fever, sore throat, minor aches & pains, cough, sneezing, runny nose

8 FL OZ (236 mL)

**Drug Facts (continued)**

**Do not use** with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have** liver disease, glaucoma, cough that occurs with too much phlegm (mucus), a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema, trouble urinating due to an enlarged prostate gland.

**Ask a doctor or pharmacist before use if you are** taking sedatives or tranquilizers, taking the blood thinning drug warfarin.

Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands. \*This product is not manufactured or distributed by The Procter & Gamble Company, distributor of Vicks® NyQuil™ High Blood Pressure Cold & Flu.

DISTRIBUTED BY: WALGREEN CO.  
200 WILMOT RD., DEERFIELD, IL 60015  
100% SATISFACTION GUARANTEED  
walgreens.com  
©2023 Walgreen Co.

ITEM 642758 W10017-0423-F

3 11917 00833 2

W3ORG0423-F 62108WAGLR

LOT: \_\_\_\_\_

EXP: \_\_\_\_\_

**PEEL BACK HERE FOR DRUG FACTS ►**

**Drug Facts (continued)**

**When using this product** ■ excitability may occur, especially in children ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery ■ alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if** ■ pain or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** ■ take only as directed ■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL per dose every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

**Other information**

- each 30 mL contains: sodium 16 mg
- store at room temperature

**Inactive ingredients** anhydrous citric acid, edetate disodium, FD&C Blue No. 1, FD&C Red No. 40, flavor, propylene glycol, propyl gallate, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose.

**Questions or comments?** 1-866-467-2748

# WALGREEN HIGH BLOOD PRESSURE NIGHTTIME COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, and doxylamine succinate liquid

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0363-6210
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>PROPYL GALLATE</b> (UNII: 8D45NN7V92)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM CITRATE, UNSPECIFIED FORM</b> (UNII: 1Q73Q2JULR)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-6210-08	236 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/03/2023	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/03/2023	

**Labeler** - WALGREENS CO. (008965063)

