

**ZYRTEC-D- cetirizine hydrochloride and pseudoephedrine hydrochloride tablet, film coated, extended release**  
**Kenvue Brands LLC**

-----  
**ZYRTEC-D**

**Drug Facts**

<b>Active ingredients (in each extended release tablet)</b>	<b>Purpose</b>
Cetirizine HCl 5 mg	Antihistamine
Pseudoephedrine HCl 120 mg	Nasal decongestant

**Uses**

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
  - runny nose
  - sneezing
  - itchy, watery eyes
  - itching of the nose or throat
  - nasal congestion
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily restores freer breathing through the nose

**Warnings**

**Do not use**

- if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- heart disease
- thyroid disease
- diabetes
- glaucoma
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- liver or kidney disease. Your doctor should determine if you need a different dose.

**Ask a doctor or pharmacist before use if you are** taking tranquilizers or sedatives.

## When using this product

- **do not use more than directed**
- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

## Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- you get nervous, dizzy, or sleepless
- symptoms do not improve within 7 days or are accompanied by fever

## If pregnant or breast-feeding:

- if breast-feeding: not recommended
- if pregnant: ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

## Directions

- do not break or chew tablet; swallow tablet whole

adults and children 12 years and over	take 1 tablet every 12 hours; do not take more than 2 tablets in 24 hours.
adults 65 years and over	ask a doctor
children under 12 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

## Other information

- store between 20° to 25°C (68° to 77°F)
- **do not use if blister unit is torn or broken**

## Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide

## Questions?

call **1-800-343-7805** (toll-free) or **215-273-8755** (collect)

**PRINCIPAL DISPLAY PANEL**

***Original Prescription Strength***

***New look***

***Same***

***relief***

NDC 50580-719-24

**ZYRTEC-D**®

*Cetirizine HCl* **5 mg**/antihistamine

*Pseudoephedrine HCl* **120 mg**/ nasal decongestant

*Extended Release Tablets*

Indoor &

Outdoor

Allergies

ALLERGY

+ SINUS

NASAL CONGESTION

+ SINUS PRESSURE

**12**

***Hour***

***relief of***

- ***Sneezing***
- ***Runny Nose***
- ***Itchy, Watery Eyes***
- ***Itchy Throat or Nose***

***Actual Size***

**24**

***Extended***

***Release***

***Tablets***



Made in Belgium  
 Distributed by:  
**JOHNSON & JOHNSON CONSUMER INC.**  
 McNeil Consumer Healthcare Division  
 Fort Washington, PA 19034 USA  
 ©J&JCI 2024  
 www.zyrtec.com

**Drug Facts (continued)**  
**Inactive ingredients** colloidal silicon dioxide, croscarmellose sodium, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide  
**Questions?** call 1-800-343-7805 (toll-free) or 215-273-8755 (collect)

**Drug Facts (continued)**  
 Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.  
**When using this product** do not use more than directed.  
 ■ avoid alcoholic drinks  
 ■ drowsiness may occur  
 ■ alcohol, sedatives, and tranquilizers may increase drowsiness  
 ■ be careful when driving a motor vehicle or operating machinery  
**Stop use and ask a doctor if**  
 ■ an allergic reaction to this product occurs. Seek medical help right away.  
 ■ you get nervous, dizzy, or sleepless  
 ■ symptoms do not improve within 7 days or are accompanied by fever  
**If pregnant or breast-feeding:**  
 ■ if pregnant: ask a health professional before use.  
 ■ if breast-feeding: not recommended.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)  
**Directions**  
 ■ do not break or chew tablet; swallow tablet whole  
 adults and children take 1 tablet every 12 hours; do not take more than 2 tablets in 24 hours.  
 adults 65 years and over ask a doctor.  
 children under 12 years of age ask a doctor.  
 consumers with liver or kidney disease ask a doctor.  
**Other information** ■ store between 20° to 25°C (68° to 77°F)  
 ■ do not use if blister unit is torn or broken

**Drug Facts**  
**Active ingredients (in each extended release tablet)** Purpose  
 Cetirizine HCl 5 mg.....Antihistamine  
 Pseudoephedrine HCl 120 mg.....Nasal decongestant  
**Uses**  
 ■ temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itchy, watery eyes ■ itching of the nose or throat  
 ■ nasal congestion  
 ■ redness/swelling of nasal passages  
 ■ temporarily relieves sinus congestion and pressure  
 ■ temporarily restores free breathing through the nose  
**Warnings**  
 ■ if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine (MAOI) you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
**Ask a doctor before use if you have**  
 ■ heart disease ■ thyroid disease ■ diabetes ■ glaucoma ■ trouble urinating due to an enlarged prostate gland ■ liver or kidney disease. Your doctor should determine if you need a different dose.  
**Do not use**  
 ■ if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine (MAOI)

The trade dress of this ZYRTEC® package is subject to trademark protection. 30059503 www.kenvuepats.com



**Original Prescription Strength** New look Same relief

NDC 50580-719-24

# ZYRTEC-D®

Cetirizine HCl 5 mg/antihistamine  
 Pseudoephedrine HCl 120 mg/nasal decongestant  
 Extended Release Tablets

**ALLERGY + SINUS**

**NASAL CONGESTION + SINUS PRESSURE**

**24 Extended Release Tablets**

12 Hour relief of

Sneezing

Runny Nose

Itchy, Watery Eyes

Itchy Throat or Nose

Actual Size

**ZYRTEC-D**  
 cetirizine hydrochloride and pseudoephedrine hydrochloride tablet, film coated, extended release

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:50580-719
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>CETIRIZINE HYDROCHLORIDE</b> (UNII: 64O047KTOA) (CETIRIZINE - UNII:YO7261ME24)	CETIRIZINE HYDROCHLORIDE	5 mg
<b>PSEUDOEPHEDRINE HYDROCHLORIDE</b> (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	120 mg

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>CROSCARMELLOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ05DW1A)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	ZYRTEC;D
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50580-719-12	2 in 1 CARTON	02/28/2023	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:50580-719-24	4 in 1 CARTON	02/28/2023	
2		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
NDA	NDA021150	02/28/2023	

---

**Labeler** - Kenvue Brands LLC (118772437)

Revised: 6/2025

Kenvue Brands LLC