MECLIZINE- meclizine hcl 12.5 mg tablet SDA Laboratories, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredient (in each caplet)

Meclizine HCl 12.5mg

Purpose

Antiemetic

Uses

prevents and treats nausea, vomiting or dizziness associated with motion sickness

Warnings

Do not use for children under 12 years of age unless directed by a doctor.

Do not take unless directed by a doctor if you have

- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis

Do not take if you are taking sedatives or tranquilizers, without first consulting your doctor.

When using this product

- do not exceed recommended dosage
- drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

• doage should be taken 1 hour before travel starts

Adults and children 12	take 2 or 4 caplets once daily or as
years and over	directed by doctor

Other information

- **Tamper Evident:** do not use if safety seal under cap is broken or missing
- store at room temperature 20°-25°C (68°-77°F)

Inactive ingredients

Croscarmellose sodium, dicalcium phosphate, magnesium stearate, microcrystalline cellulose, silicon dioxide, stearic acid

Questions?

Adverse drug event call (800) 687-0176 (M - F, 8AM - 4PM EST).

NDC 66424-0386-10

MECLIZINE

Meclizine HCl 12.5mg

Anti-Emetic

• Motion Sickness • Nausea • Vomiting

1000 CAPLETS

SDA

LABORATORIES



-	
Drug Facts	
Active ingredient (in each caplet) Meclizine HCl 12.5mg	Purpose Antiemetic
Uses prevents and treats nausea, vor with motion sickness	nausea, vomiting or dizziness associated
Warnings Do not use for children under 12 years of age unless directed by a doctor. It you have be not take unless directed by a doctor if you have endersoms - incube univation due ha an endersord mosteria.	2 years of age unless directed radoctor if you have
gadoonia a trouve unitating doe to an entanged prostate by a breathing problem such as emphysema or chronic bronchitis. Do not take if you are taking sedatives or tranquilizers, without	te to an enanged prostate the as emphysema or chronic affixes or tranquilizers, without
first consulting your doctor. When using this product a do not exceed recommended by the using this product a do not exceed recommended becage a drowsiness may occur a elochol, sedatives, and tranquities may increase drowsiness a and adonloic of this a becareful when driving a motor vehicle or operating machinery.	not exceed recommended = alcohol, sedatives, and siness = avoid alcoholic drinks vehicle or operating machinery
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children, in case of overdose, get medical help or contact a Poison Control Center right away.	ask a health professional before Iren. In case of overdose, get ontrol Center right away.
Directions ■ dosage should be taken 1 hour before travel starts	ir before travel starts
Adults and children tak	take 2 or 4 caplets once daily or as directed by doctor
Other information I amper Evident: do not use if safety seal under cap is broken or missing store at room temperature 20°-28°C (68°-77°F)	safety seal under cap is -25°C (68°-77°F)
Inactive ingredients Croscamellose sodium, dicale phosphate, magnesium stearate, microcrystalline cellulose, silicon dioxide, stearic acid	scarmellose sodium, dicalcium , microcrystalline cellulose,
drug	event call (800) 687-0176
Distributed by: SDA Laboratories, Inc., Greenwich, CT 06830 MADE IN USA Org 02/15	enwich, CT 06830
	86410

MECLIZINE

meclizine hcl 12.5 mg tablet

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:66424-386
Route of Administration	ORAL		

Active Ingredient/Active Moiety Ingredient Name Basis of Strength MECLIZINE HYDRO CHLO RIDE (UNII: HDP7W44CIO) (MECLIZINE - UNII:3L5TQ84570) MECLIZINE HYDRO CHLO RIDE 12.5 mg

Inactive Ingredients			
Ingredient Name	Strength		
CROSCARMELLOSE SODIUM (UNII: M28 OL1HH48)			
CALCIUM PHO SPHATE, DIBASIC, ANHYDRO US (UNII: L11K75P92J)			
MAGNESIUM STEARATE (UNII: 70097M6I30)			
CELLULOSE, MICRO CRYSTALLINE (UNII: OP1R32D61U)			
SILICON DIO XIDE (UNII: ETJ7Z6XBU4)			
STEARIC ACID (UNII: 4ELV7Z65AP)			

Product Characteristics				
Color	white (WHITE COLOR)	Score	no score	
Shape	CAPSULE (CAPSULE SHAPED TABLET)	Size	13mm	
Flavor		Imprint Code	PH0 49	
Contains				

Packaging			
# Item Code	Package Description	Marketing Start Date	Marketing End Date
NDC:66424-386- 01	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	0 2/0 1/20 18	
NDC:66424-386- 10	1000 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	0 2/0 1/20 18	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part336	0 2/0 1/20 18	

Labeler - SDA Laboratories, Inc. (948067889)

Registrant - Pharbest Pharmaceuticals, Inc. (557054835)

Establishment			
Name	Address	ID/FEI	Business Operations
Pharbest Pharmaceuticals, Inc.		557054835	manufacture(66424-386), pack(66424-386), analysis(66424-386)

Revised: 2/2018 SDA Laboratories, Inc.