

**COLD AND FLU DAYTIME- acetaminophen, dextromethorphan hbr,
phenylephrine hcl solution
Bi-Mart**

Bi-Mart 44-012

Active ingredients (in each 15 mL)

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer
Cough suppressant
Nasal decongestant

Uses

- temporarily relieves common cold and flu symptoms:
 - sore throat
 - nasal congestion
 - fever
 - headache
 - minor aches and pains
 - cough due to minor throat and bronchial irritation

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a

doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if the user has

- difficulty in urination due to enlargement of the prostate gland
- cough that occurs with too much phlegm (mucus)
- liver disease
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- heart disease
- diabetes
- thyroid disease
- high blood pressure

Ask a doctor or pharmacist before use if the user is

taking the blood thinning drug warfarin.

When using this product

do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**
- mL = milliliter; FL OZ = fluid ounce
- only use the dose cup provided
- do not take more than 5 doses per 24 hours

adults and children 12 years and over	30 mL every 4 hours
children 6 to under 12 years	15 mL every 4 hours
children under 6 years	do not use

Other information

- **each 15 mL contains:** sodium 13 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

anhydrous citric acid, FD&C yellow #6, flavors, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sodium metabisulfite, sodium saccharin, sorbitol, sucralose

Questions or comments?

1-800-935-6737

Principal display panel**BI-MART**

Compare to the active ingredients
in Vicks® DayQuil® Cold & Flu*

**DAYTIME
COLD & FLU**

Acetaminophen (Pain reliever/fever reducer)
Dextromethorphan HBr (Cough suppressant)
Phenylephrine HCl (Nasal decongestant)

Headache, Fever, Sore Throat,
Minor Aches & Pains,
Nasal Congestion,
Cough

Alcohol Free

Menthol Flavored

12 FL OZ (355 mL)

TAMPER EVIDENT: DO NOT USE IF PRINTED

NECK WRAP IS BROKEN OR MISSING

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

BI-MART UNCONDITIONAL MONEY BACK GUARANTEE

If you are not completely satisfied with your BI-MART QUALITY PRODUCT, regardless of the reason, return the unused portion and your entire purchase price will be cheerfully refunded.

*This product is not manufactured or distributed by The Procter & Gamble Company, distributor of Vicks® DayQuil® Cold & Flu.

50844 REV0722A01202

Distributed by:

BI-MART

Eugene, OR 97402

TAMPER EVIDENT: DO NOT USE IF PRINTED
NECK WRAP IS BROKEN OR MISSING



PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

BI-MART UNCONDITIONAL MONEY BACK GUARANTEE
If you are not completely satisfied with your BI-MART
QUALITY PRODUCT, regardless of the reason, return the
unused portion and your entire purchase price will be
cheerfully refunded.

*This product is not manufactured or
distributed by The Procter & Gamble
Company, distributor of Vicks®
DayQuil® Cold & Flu.

50844 REV0722A01202

Distributed by:
BI-MART
Eugene, OR 97402

3001000549

G9526-012-11-1

No Print / No Varnish Area
Lot # and Exp. Info

BI-MART

Compare to the active ingredients
in Vicks® DayQuil® Cold & Flu*

DAYTIME

COLD & FLU

Acetaminophen (Pain reliever/fever reducer)
Dextromethorphan HBr (Cough suppressant)
Phenylephrine HCl (Nasal decongestant)

Headache, Fever, Sore Throat,
Minor Aches & Pains,
Nasal Congestion,
Cough



Menthol Flavored

Alcohol Free

12 FL OZ (355 mL)

Drug Facts

Active ingredients (in each 15 mL)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses ■ temporarily relieves common cold and flu symptoms:
■ sore throat ■ nasal congestion ■ fever
■ headache ■ minor aches and pains
■ cough due to minor throat and bronchial irritation

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if
■ adult takes more than 4,000 mg of acetaminophen in 24 hours
■ child takes more than 5 doses in 24 hours
■ taken with other drugs containing acetaminophen
■ adult has 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if the user has

■ difficulty in urination due to enlargement of the prostate gland
■ cough that occurs with too much phlegm (mucus) ■ liver disease
■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema ■ heart disease ■ diabetes ■ thyroid disease
■ high blood pressure

PEEL HERE FOR MORE DRUG FACTS

Questions or comments? 1-800-935-6737
Inactive ingredients anhydrous citric acid, FD&C yellow #6, sodium benzoate, sodium citrate dihydrate, sodium metabisulfite, sodium saccharin, sorbitol, sucralose
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ use by expiration date on package
Other information ■ each 15 mL contains: sodium 13 mg
adults and children 12 years and over 30 mL every 4 hours
children 6 to under 12 years 15 mL every 4 hours
children under 6 years do not use
Directions ■ do not take more than directed ■ mL = milliliter; FL OZ = fluid ounce ■ only use the dose cup provided ■ do not take more than 5 doses per 24 hours
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
■ cough comes back or occurs with rash or headache that lasts. These (children) or 7 days (adults) ■ pain, nasal congestion, or cough gets worse or lasts more than 5 days ■ redness or swelling is present ■ new symptoms occur ■ fever gets worse or lasts more than 3 days ■ nervousness, dizziness, or sleeplessness occur
Stop use and ask a doctor if When using this product do not exceed recommended dosage. ■ when using this product do not exceed recommended dosage. ■ thinning drug warfarin.
Drug Facts (continued) Ask a doctor or pharmacist before use if the user is taking the blood

Bi-Mart 44-012

COLD AND FLU DAYTIME

acetaminophen, dextromethorphan hbr, phenylephrine hcl solution

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:37835-526

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
SODIUM METABISULFITE (UNII: 4VON5FNS3C)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color	orange	Score	
Shape		Size	
Flavor	MENTHOL	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37835-526-02	355 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/06/2023	
2	NDC:37835-526-30	354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/18/2018	01/06/2023

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/18/2018	

Labeler - Bi-Mart (027630078)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	manufacture(37835-526) , pack(37835-526)

Revised: 1/2025

Bi-Mart