

**COLD AND FLU DAYTIME- acetaminophen, dextromethorphan hbr,  
phenylephrine hcl solution  
Bi-Mart**

-----

**Bi-Mart 44-012**

***Active ingredients (in each 15 mL)***

Acetaminophen 325 mg  
Dextromethorphan HBr 10 mg  
Phenylephrine HCl 5 mg

***Purpose***

Pain reliever/fever reducer  
Cough suppressant  
Nasal decongestant

***Uses***

- temporarily relieves common cold and flu symptoms:
  - sore throat
  - nasal congestion
  - fever
  - headache
  - minor aches and pains
  - cough due to minor throat and bronchial irritation

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a

doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if the user has**

- difficulty in urination due to enlargement of the prostate gland
- cough that occurs with too much phlegm (mucus)
- liver disease
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- heart disease
- diabetes
- thyroid disease
- high blood pressure

### **Ask a doctor or pharmacist before use if the user is**

taking the blood thinning drug warfarin.

### **When using this product**

**do not exceed recommended dosage.**

### **Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- **do not take more than directed**
- mL = milliliter; FL OZ = fluid ounce
- only use the dose cup provided
- do not take more than 5 doses per 24 hours

adults and children 12 years and over	30 mL every 4 hours
children 6 to under 12 years	15 mL every 4 hours
children under 6 years	do not use

**Other information**

- **each 15 mL contains:** sodium 13 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

**Inactive ingredients**

anhydrous citric acid, FD&C yellow #6, flavors, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sodium metabisulfite, sodium saccharin, sorbitol, sucralose

**Questions or comments?**

**1-800-935-6737**

**Principal display panel****BI-MART**

Compare to the active ingredients  
in Vicks® DayQuil® Cold & Flu\*

**DAYTIME  
COLD & FLU**

**Acetaminophen** (Pain reliever/fever reducer)  
**Dextromethorphan HBr** (Cough suppressant)  
**Phenylephrine HCl** (Nasal decongestant)

Headache, Fever, Sore Throat,  
Minor Aches & Pains,  
Nasal Congestion,  
Cough

Alcohol Free

*Menthol Flavored*

**12 FL OZ (355 mL)**

**TAMPER EVIDENT: DO NOT USE IF PRINTED**

## **NECK WRAP IS BROKEN OR MISSING**

### **PARENTS:**

Learn about teen medicine abuse

**[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)**

### **BI-MART UNCONDITIONAL MONEY BACK GUARANTEE**

If you are not completely satisfied with your BI-MART QUALITY PRODUCT, regardless of the reason, return the unused portion and your entire purchase price will be cheerfully refunded.

\*This product is not manufactured or distributed by The Procter & Gamble Company, distributor of Vicks® DayQuil® Cold & Flu.

50844 REV0722A01202

Distributed by:

**BI-MART**

Eugene, OR 97402

TAMPER EVIDENT: DO NOT USE IF PRINTED  
NECK WRAP IS BROKEN OR MISSING



**PARENTS:**

Learn about teen medicine abuse

[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

**BI-MART UNCONDITIONAL MONEY BACK GUARANTEE**  
If you are not completely satisfied with your BI-MART  
QUALITY PRODUCT, regardless of the reason, return the  
unused portion and your entire purchase price will be  
cheerfully refunded.

\*This product is not manufactured or  
distributed by The Procter & Gamble  
Company, distributor of Vicks®  
DayQuil® Cold & Flu.

50844 REV0722A01202

Distributed by:  
**BI-MART**  
Eugene, OR 97402

3001000549

**G9526-012-11-1**

No Print / No Varnish Area  
Lot # and Exp. Info

**BI-MART**

Compare to the active ingredients  
in Vicks® DayQuil® Cold & Flu\*

# DAYTIME

## COLD & FLU

**Acetaminophen** (Pain reliever/fever reducer)  
**Dextromethorphan HBr** (Cough suppressant)  
**Phenylephrine HCl** (Nasal decongestant)

Headache, Fever, Sore Throat,  
Minor Aches & Pains,  
Nasal Congestion,  
Cough



Menthol Flavored

Alcohol Free

**12 FL OZ (355 mL)**

### Drug Facts

Active ingredients (in each 15 mL)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

**Uses** ■ temporarily relieves common cold and flu symptoms:  
■ sore throat ■ nasal congestion ■ fever  
■ headache ■ minor aches and pains  
■ cough due to minor throat and bronchial irritation

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if  
■ adult takes more than 4,000 mg of acetaminophen in 24 hours  
■ child takes more than 5 doses in 24 hours  
■ taken with other drugs containing acetaminophen  
■ adult has 3 or more alcoholic drinks every day while using this product  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash  
If a skin reaction occurs, stop use and seek medical help right away.  
**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
■ if you have ever had an allergic reaction to this product or any of its ingredients

### Ask a doctor before use if the user has

■ difficulty in urination due to enlargement of the prostate gland  
■ cough that occurs with too much phlegm (mucus) ■ liver disease  
■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema ■ heart disease ■ diabetes ■ thyroid disease  
■ high blood pressure

**PEEL HERE FOR MORE DRUG FACTS**

**Drug Facts (continued)**

Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin.

When using this product do not exceed recommended dosage.

**Stop use and ask a doctor if**  
■ nervousness, dizziness, or sleeplessness occur  
■ fever gets worse or lasts more than 3 days  
■ redness or swelling is present ■ new symptoms occur  
■ pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults)  
■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**  
■ do not take more than directed  
■ mL = milliliter; FL OZ = fluid ounce  
■ only use the dose cup provided  
■ do not take more than 5 doses per 24 hours

adults and children 12 years and over	30 mL every 4 hours
children 6 to under 12 years	15 mL every 4 hours
children under 6 years	do not use

**Other information** ■ each 15 mL contains: sodium 13 mg  
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ use by expiration date on package

**Inactive ingredients** anhydrous citric acid, FD&C yellow #6, sodium benzoate, sodium citrate dihydrate, sodium metabisulfite, sodium saccharin, sorbitol, sucralose

**Questions or comments? 1-800-935-6737**

**Bi-Mart 44-012**

## COLD AND FLU DAYTIME

acetaminophen, dextromethorphan hbr, phenylephrine hcl solution

### Product Information

**Product Type**

HUMAN OTC DRUG

**Item Code (Source)**

NDC:37835-526

Route of Administration ORAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>TRISODIUM CITRATE DIHYDRATE</b> (UNII: B22547B95K)	
<b>SODIUM METABISULFITE</b> (UNII: 4VON5FNS3C)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	

### Product Characteristics

<b>Color</b>	orange	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	MENTHOL	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37835-526-02	355 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/06/2023	
2	NDC:37835-526-30	354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/18/2018	01/06/2023

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/18/2018	

**Labeler** - Bi-Mart (027630078)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	manufacture(37835-526) , pack(37835-526)

Revised: 1/2026

Bi-Mart