

NAPROXEN- naproxen tablet
Safeway, Inc

Pain Relief

Naproxen Sodium Tablets USP, 220 mg (NSAID)
Pain reliever/Fever reducer

contains no ingredient made from a gluten-containing grain (wheat, barley, or rye)

Strength to last 12 hours

Active Ingredient (in each caplet)

Naproxen sodium 220mg (NSAID)

Purposes

Pain Reliever/Fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- headache
- toothache
- the common cold
- temporarily reduces fever

Allergy Alert

Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives

■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

Adults and children 12 years and older ■ take 1 caplet every 8 to 12 hours while symptoms last

- for the first dose you may take 2 caplets within the first hour
- do not exceed 2 caplets in any 8- to 12-hour period
- do not exceed 3 caplets in a 24-hour period

Children under 12 years ■ ask a doctor

Other Information

- **each caplet contains:** sodium 20 mg
- store at 20 - 25°C (68 - 77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive Ingredients

FD&C blue#2 aluminum lake, hypromellose 2910, maize starch, microcrystalline cellulose, polyethylene glycol, povidone k-30, sodium starch glycolate, stearic acid, titanium dioxide

Questions or Comments?

call 1 -877-770-3183 Mon – Fri 8:00 AM EST to 5:00 PM PST

Principal Display Panel

COATING
FREE AREA

700000004585
200000003480

DISTRIBUTED BY:
BAYER HEALTHCARE
21000
PISCATAWAY, NJ 08854-4000
1-877-770-3183

Scan here
for more
information

MADE IN INDIA

PAPER BOX PLASTIC BOTTLE

size	: 1-3/4 X 1-3/4 X 3-3/8
ref #	: PPI180604B
material	: .016 SBS

COATING
FREE AREA

200000004585
700000003480

COATING
FREE AREA

Drug Facts (continued)

- Have 3 or more alcoholic drinks every day while using this product
- Take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- If you have ever had an allergic reaction to any other pain reliever/fever reducer
- Right before or after heart surgery

Ask a doctor before use if

- The stomach bleeding warning applies to you
- You have a history of stomach problems, such as heartburn
- You have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- You are taking a diuretic
- You have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- Under a doctor's care for any serious condition
- Taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- Taking any other drug

When using this product

- Take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- You experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- Have stomach pain that does not get better
- You have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- Pain gets worse or lasts more than 10 days
- Fever gets worse or lasts more than 3 days
- You have difficulty swallowing

Drug Facts (continued)

- If tablets (like the pill) is stuck in your throat
- Redness or swelling is present in the painful area
- Any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions or do not take more than directed

- The smallest effective dose should be used
- Drink a full glass of water with each dose

Adults and children 12 years and older

- Take 1 caplet every 8 to 12 hours while symptoms last
- For the first dose you may take 2 caplets within the first hour
- Do not exceed 2 caplets in any 8- to 12-hour period
- Do not exceed 3 caplets in a 24-hour period

Children under 12 years

- Ask a doctor

Other information

- Each caplet contains sodium 20 mg
- Store at 20° - 25°C (68° - 77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients

FD&C Blue #2 aluminum lake, hydroxyethylcellulose 2910, maize starch, microcrystalline cellulose, polyethylene glycol, povidone I-30, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?

call 1-877-770-3183 Mon – Fri 8:00 AM EST to 5:00 PM PST

COATING
FREE AREA

POVIDONE K30 (UNII: U725QWY32X)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
STARCH, CORN (UNII: O8232NY3SJ)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	

Product Characteristics

Color	blue (light blue)	Score	no score
Shape	OVAL ((Caplet -Shaped))	Size	12mm
Flavor		Imprint Code	220
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-062-20	200 in 1 BOTTLE; Type 0: Not a Combination Product	09/30/2023	
2	NDC:21130-062-09	90 in 1 BOTTLE; Type 0: Not a Combination Product	08/30/2023	
3	NDC:21130-062-05	50 in 1 BOTTLE; Type 0: Not a Combination Product	08/30/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091353	08/30/2023	

Labeler - Safeway, Inc (009137209)

Revised: 12/2025

Safeway, Inc