

ACETAMINOPHEN, ASPIRIN AND CAFFEINE- acetaminophen, aspirin and caffeine tablet
Safeway, Inc

Migraine Relief

Acetaminophen, Aspirin and Caffeine Tablets, USP 250 mg, 250 mg and 65 mg

Pain Reliever / Pain Reliever Aid

Active ingredients (in each caplet)

Acetaminophen 250mg
Aspirin 250 mg (NSAID)**
Caffeine 65 mg

**nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/Pain reliever aid

Use

- treats migraine

Reye's syndrome:

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert:

Acetaminophen may cause severe skin reactions.
Symptoms may include: ■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert:

Allergy alert: Aspirin may cause a severe allergic reaction which may include:
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 2 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Medication overuse headache warning:

Headaches may worsen if this product is used for 10 or more days per month

Caffeine warning:

The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine containing medications, foods or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness and occasionally rapid heart beat.

Do not use

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if

- you have never had migraines diagnosed by a health professional
- you have a headache that is different from your usual migraines
- you have the worst headache of your life
- you have fever and stiff neck
- you have headaches beginning after or caused by head injury, exertion, coughing or bending
- you experienced your first headache after the age of 50
- you have daily headaches

- you have a migraine so severe as to require bed rest
- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have problems or serious side effects from taking pain relievers or fever reducers

- you have vomiting with your migraine headache

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for: • diabetes • gout • arthritis
- under a doctor's care for any serious condition
- taking any other drug
- taking any other product that contains aspirin, acetaminophen or any other pain reliever/fever reducer

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint • vomit blood • have bloody or black stools
 - have stomach pain that does not get better
- your migraine is not relieved or worsens after first dose
- new or unexpected symptoms occur
- ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep Out of Reach of Children

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed

adults take 2 caplets with a glass of water

- if symptoms persist or worsen, ask your doctor
- do not take more than 2 caplets in 24 hours, unless directed by a doctor

COATING
FREE AREA

200000004489
700000003411

Drug Facts (continued)

- Take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- Have 2 or more alcoholic drinks every day while using this product
- Take more or for a longer time than directed
- Medication overuse headache warning:** Headaches may worsen if this product is used for 10 or more days per month
- Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness and occasionally rapid heart beat.

Do not use

- If you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- With any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if

- You have never had migraines diagnosed by a health professional
- You have a headache that is different from your usual migraines
- You have the worst headache of your life
- You have fever and stiff neck
- You have headaches beginning after or caused by head injury, exertion, coughing or bending
- You have headaches beginning after the age of 50
- You have sinus headaches
- You have a migraine so severe as to require bed rest
- You have liver disease
- Stomach bleeding warning applies to you
- You have a history of stomach problems, such as heartburn
- You have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- You are taking a diuretic
- You have asthma
- You have problems or serious side effects from taking pain

Drug Facts (continued)

- Relieves or fever reducers
- You have vomiting with your migraine headache
- Ask a doctor or pharmacist before use if you are
 - taking a prescription drug for: • diabetes • gout • arthritis
 - under a doctor's care for any serious condition
 - taking any other drug
 - taking any other product that contains aspirin, acetaminophen or any other pain reliever/fever reducer

Stop use and ask a doctor if

- An allergic reaction occurs. Seek medical help right away.
- You experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- Have stomach pain that does not get better
- Your migraine is not relieved or worsens after first dose
- New or unexpected symptoms occur
- Ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 30 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep Out of Reach of Children

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions • do not use more than directed

Adults: Take 2 caplets with a glass of water

- If symptoms persist or worsen, ask your doctor
- Do not take more than 2 caplets in 24 hours, unless directed by a doctor

Under 18 years of age: ask a doctor

Other information

- Store at controlled room temperature 20° to 25°C (68° to 77°F)
- Close cap tightly after use
- Read all product information

Drug Facts (continued)

Keep this caplet for important information.

Inactive ingredients: croscarmellose sodium, hydroxypropyl methylcellulose, hydroxypropyl methylcellulose, polyethylene glycol, sodium lauryl sulfate and titanium dioxide

Questions or comments?

For more information, visit us at www.signaturecare.com or call 1-800-333-3333. ©2018 B&B LDC, Inc. All rights reserved.

This product is manufactured and controlled by B&B LDC, Inc. (Signature Care), a subsidiary of Crocker/Magnate HealthCare, a division of Crocker/Magnate HealthCare.

Scan here for more information



size : 1+3/4 X 1+3/4 X 3+3/8
ref # : PP180604B
material : .016 SBS



Quality Guaranteed

Compare to
Excedrin® Migraine
active ingredient*
NDC 21130-681-05

IMPORTANT: READ ALL PRODUCT INFORMATION BEFORE USING. KEEP CARTON FOR IMPORTANT INFORMATION.

TAMPER EVIDENT: DO NOT USE IF POUCH INNER SEAL UNDER THE CAPS IS OPENED OR MISSING.

Drug Facts

Active ingredients (in each caplet) **Purpose**

Acetaminophen 250 mg Pain reliever

Aspirin 250 mg (NSAID)* Pain reliever

Caffeine 65 mg Pain reliever aid

*nonsteroidal anti-inflammatory drug

Use • treats migraine

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: skin redness, hives, rash. If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 2 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

Migraine Relief

ACETAMINOPHEN, ASPIRIN AND CAFFEINE TABLETS, USP

250 mg, 250 mg and 65 mg

Pain Reliever/Pain Reliever Aid

Actual size



50 Caplets*
*Capsule-Shaped Tablets



MADE IN INDIA

DISTRIBUTED BY:
BETTER LIVING BRANDS LLC
P.O. BOX 99,
PLEASANTON, CA 94566-0099
1-888-732-0069 RD 23126

3 21130 78774 7

Lot
Exp.

COATING
FREE AREA

200000004489
700000003411

COATING
FREE AREA



size : 1+3/4 X 1+3/4 X 3+3/8
 ref # : PP1806048
 material : .016 SBS

Drug Facts (continued)

- Take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- Take more or for a longer time than directed
- Medication overdose headache warning: Headaches may worsen if this product is used for 10 or more days per month
- Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods or beverages while taking this product because too much caffeine may cause nervousness, irritability, dizziness and occasionally rapid heart beat.
- Do not use
 - if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
 - with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- Ask a doctor before use if
 - you have never had migraines diagnosed by a health professional
 - you have a headache that is different from your usual migraines
 - you have the worst headache of your life
 - you have fever and stiff neck
 - you have headaches beginning after or caused by head injury, exertion, coughing or bending
 - you experienced your first headache after the age of 50
 - you have daily headaches
 - you have a migraine so severe as to require bed rest
 - you have liver disease
 - stomach bleeding warning applies to you
 - you have a history of stomach problems, such as heartburn
 - you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
 - you are taking a diuretic
 - you have asthma
 - you have problems or serious side effects from taking pain

Drug Facts (continued)

- relieves or lessens
- you have vomiting with your migraine headache
- Ask a doctor or pharmacist before use if you are
 - taking a prescription drug for: • diabetes • gout • arthritis
 - under a doctor's care for any serious condition
 - taking any other drug
 - taking any other product that contains aspirin, acetaminophen or any other pain reliever/fever reducer
- Stop use and ask a doctor if
 - an allergic reaction occurs. Seek medical help right away.
 - you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - your migraine is not relieved or worsens after first dose
 - new or unexpected symptoms occur
 - ringing in the ears or loss of hearing occurs
- If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
- Keep Out of Reach of Children
- In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
- Directions
 - do not use more than directed.
 - adults take 2 caplets with a glass of water
 - if symptoms persist or worsen, ask your doctor
 - do not take more than 2 caplets in 24 hours, unless directed by a doctor
 - under 18 years of age ask a doctor
- Other information
 - store at controlled room temperature 20° to 25°C (68° to 77°F)
 - close cap tightly after use
 - read product information

Signature Care.
 Quality Guaranteed

Compare to Excedrin[®] Migraine active ingredient!
 NDC 21130-681-24

Migraine Relief
 ACETAMINOPHEN, ASPIRIN AND CAFFEINE TABLETS, USP
 250 mg, 250 mg and 65 mg
 Pain Reliever/Pain Reliever Aid

Actual size

24 Caplets[†]
 Caplet-shaped Tablets

MADE IN INDIA
 BETTER LIVING BRANDS LLC
 210, 3024 99
 PLEASANTON, CA 94566-0099
 1-800-729-1929 RD 22128

LOT
 Exp.

IMPORTANT: READ ALL PRODUCT INFORMATION BEFORE USING. KEEP CARTON FOR IMPORTANT INFORMATION.

TAMPER EVIDENT: DO NOT USE IF FOIL INNER SEAL UNDER THE CAP IS OPENED OR MISSING

Drug Facts

Active ingredients (in each caplet)	Purposes
Acetaminophen 250 mg	Pain reliever
Aspirin 250 mg (NSAID) ^{**}	Pain reliever
Caffeine 65 mg	Pain reliever aid

Use • treats migraine

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms can be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: skin redness, hives, rash

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- facial swelling
- asthma (wheezing)
- shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 2 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

ACETAMINOPHEN, ASPIRIN AND CAFFEINE

acetaminophen, aspirin and caffeine tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:21130-681
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

Inactive Ingredients

Ingredient Name	Strength
HYDROXYPROPYL CELLULOSE (110000 WAMW) (UNII: 5Y0974F5PW)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
HYPROMELLOSE 2910 (6 MPA.S) (UNII: 0WZ8WG20P6)	
SODIUM STEARYL FUMARATE (UNII: 7CV7WJK4UI)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	

Product Characteristics

Color	white	Score	no score
Shape	CAPSULE (Capsule shaped Tablet)	Size	17mm
Flavor		Imprint Code	l1
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-681-05	50 in 1 BOTTLE; Type 0: Not a Combination Product	08/04/2023	
2	NDC:21130-681-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	08/04/2023	
3	NDC:21130-681-24	24 in 1 BOTTLE; Type 0: Not a Combination Product	08/04/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA214039	08/04/2023	

Labeler - Safeway, Inc (009137209)

Revised: 12/2023

Safeway, Inc