

**ACETAMINOPHEN, ASPIRIN AND CAFFEINE- acetaminophen, aspirin and caffeine tablet**  
**Safeway, Inc**

-----  
**Migraine Relief**

**Acetaminophen, Aspirin and Caffeine Tablets, USP 250 mg, 250 mg and 65 mg**

**Pain Reliever / Pain Reliever Aid**

**Active ingredients (in each caplet)**

Acetaminophen 250mg  
Aspirin 250 mg (NSAID)\*\*  
Caffeine 65 mg

\*\*nonsteroidal anti-inflammatory drug

**Purposes**

Pain reliever/Pain reliever aid

**Use**

- treats migraine

**Reye's syndrome:**

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:**

Acetaminophen may cause severe skin reactions.  
Symptoms may include: ■ skin reddening ■ blisters ■ rash  
If a skin reaction occurs, stop use and seek medical help right away.

**Allergy alert:**

Allergy alert: Aspirin may cause a severe allergic reaction which may include:  
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

**Liver warning:**

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 2 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

### **Stomach bleeding warning:**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### **Medication overuse headache warning:**

Headaches may worsen if this product is used for 10 or more days per month

### **Caffeine warning:**

The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine containing medications, foods or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness and occasionally rapid heart beat.

### **Do not use**

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

### **Ask a doctor before use if**

- you have never had migraines diagnosed by a health professional
- you have a headache that is different from your usual migraines
- you have the worst headache of your life
- you have fever and stiff neck
- you have headaches beginning after or caused by head injury, exertion, coughing or bending
- you experienced your first headache after the age of 50
- you have daily headaches

- you have a migraine so severe as to require bed rest
- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have vomiting with your migraine headache

### **Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for: • diabetes • gout • arthritis
- under a doctor's care for any serious condition
- taking any other drug
- taking any other product that contains aspirin, acetaminophen or any other pain reliever/fever reducer

### **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint • vomit blood • have bloody or black stools
  - have stomach pain that does not get better
- your migraine is not relieved or worsens after first dose
- new or unexpected symptoms occur
- ringing in the ears or loss of hearing occurs

### **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep Out of Reach of Children**

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- do not use more than directed

adults take 2 caplets with a glass of water

- if symptoms persist or worsen, ask your doctor
- do not take more than 2 caplets in 24 hours, unless directed by a doctor

under 18 years of age ask a doctor

### **Other information**

- store at controlled room temperature 20° to 25°C (68° to 77°F)
- close cap tightly after use
- read all product information before using. Keep this box for important information.

### **Inactive ingredients**

carnauba wax, hydroxypropyl cellulose, hypromellose, microcrystalline cellulose, polyethylene glycol, sodium stearyl fumarate and titanium dioxide

### **Questions or comments?**

contact **1-877-770-3183** Mon-Fri 8:00 AM EST to 5:00 PM PST

### **Principal Display Panel**



SIZE 11\*3/4 A 1\*3/4 A 3\*3/8  
 ref # :PP180604B  
 material :.016 SBS

COATING FREE AREA

2000000489  
 70000003411

**Drug Facts (continued)**  
 Before using, keep this box for important information.  
**Inactive ingredients** include wax, hydroxypropyl methylcellulose, hydroxypropyl methylcellulose, microcrystalline cellulose, polyethylene glycol, croscarmellose sodium, and titanium dioxide.

**Questions or comments?**  
 This product is not manufactured or distributed by USBC (www.usbc.com).  
 For more information, contact 1-877-706-8188, Mon-Fri 8:00 AM EST to 5:00 PM PST.  
 Healthcare professionals: contact 1-877-706-8188.

Signature Care  
 Quality Guaranteed

Compare to Excedrin<sup>®</sup> Migraine active ingredient\*  
 NDC 21130-681-05

Scan here for more information

**Drug Facts (continued)**

relieves or fever reduces

you have vomiting with your migraine headache

**Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for: • diabetes • gout • arthritis
- under a doctor's care for any serious condition
- taking any other drug
- taking any other product that contains aspirin, acetaminophen or any other pain reliever/fever reducer

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- your migraine is not relieved or worsens after first dose
- new or unexpected symptoms occur
- ringing in the ears or loss of hearing occurs

**If pregnant or breast-feeding, ask a health professional before use.** It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep Out of Reach of Children**  
 In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** **do not use more than directed**

adults

- take 2 caplets with a glass of water
- if symptoms persist or worsen, ask your doctor
- do not take more than 2 caplets in 24 hours, unless directed by a doctor

under 18 years of age ask a doctor

**Other information**

- store at controlled room temperature 20° to 25°C (68° to 77°F)
- close cap tightly after use
- read all product information

**Drug Facts (continued)**

relieves or fever reduces

you have vomiting with your migraine headache

**Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for: • diabetes • gout • arthritis
- under a doctor's care for any serious condition
- taking any other drug
- taking any other product that contains aspirin, acetaminophen or any other pain reliever/fever reducer

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- your migraine is not relieved or worsens after first dose
- new or unexpected symptoms occur
- ringing in the ears or loss of hearing occurs

**If pregnant or breast-feeding, ask a health professional before use.** It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep Out of Reach of Children**  
 In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** **do not use more than directed**

adults

- take 2 caplets with a glass of water
- if symptoms persist or worsen, ask your doctor
- do not take more than 2 caplets in 24 hours, unless directed by a doctor

under 18 years of age ask a doctor

**Other information**

- store at controlled room temperature 20° to 25°C (68° to 77°F)
- close cap tightly after use
- read all product information

**Signature Care**  
 Quality Guaranteed

**Migraine Relief**  
 ACETAMINOPHEN, ASPIRIN AND CAFFEINE TABLETS, USP  
 250 mg, 250 mg and 65 mg  
 Pain Reliever/Pain Reliever Aid

Actual size

50 Caplets<sup>1</sup>  
<sup>1</sup>Capulet-Shaped Tablets

LOT 2113078774

MADE IN INDIA

DISTRIBUTED BY:  
 BETTER LIVING BRANDS LLC  
 P.O. BOX 99  
 PLEASANTON, CA 94566-0099  
 415-884-7538

RD 23128

3 21130 78774 7

**IMPORTANT: READ ALL PRODUCT INFORMATION BEFORE USING. KEEP CARTON FOR IMPORTANT INFORMATION.**

**TAMPER EVIDENT: DO NOT USE IF FOIL INNER SEAL UNDER THE CAP IS OPENED OR MISSING**

**Drug Facts**

**Active ingredients (in each caplet)**

Active ingredients (in each caplet)	Purposes
Acetaminophen 250 mg	Pain reliever
Aspirin 250 mg (NSAID) <sup>**</sup>	Pain reliever
Caffeine 65 mg	Pain reliever aid

<sup>\*\*</sup>nonsteroidal anti-inflammatory drug

**Use** • treats migraine

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: • skin redness • hives • rash

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 2 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

COATING FREE AREA

2000000489  
 70000003411

COATING FREE AREA

Lot

Exp.

Principal Display Panel



## ACETAMINOPHEN, ASPIRIN AND CAFFEINE

acetaminophen, aspirin and caffeine tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:21130-681
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
<b>CAFFEINE</b> (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>HYDROXYPROPYL CELLULOSE (110000 WAMW)</b> (UNII: 5Y0974F5PW)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>HYPROMELLOSE 2910 (6 MPA.S)</b> (UNII: 0WZ8WG20P6)	
<b>SODIUM STEARYL FUMARATE</b> (UNII: 7CV7WJK4UI)	
<b>POLYETHYLENE GLYCOL 400</b> (UNII: B697894SGQ)	
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)	

## Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	CAPSULE (Capsule shaped Tablet)	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	l1
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-681-05	50 in 1 BOTTLE; Type 0: Not a Combination Product	08/04/2023	
2	NDC:21130-681-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	08/04/2023	
3	NDC:21130-681-24	24 in 1 BOTTLE; Type 0: Not a Combination Product	08/04/2023	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA214039	08/04/2023	

**Labeler** - Safeway, Inc (009137209)

Revised: 12/2025

Safeway, Inc