

TOPCARE NASAL- oxymetazoline hydrochloride spray
Topco Associates LLC

Topco Associates LLC. Nasal Spray Drug Facts

Active ingredient

Oxymetazoline hydrochloride 0.05%

Purpose

Nasal decongestant

Uses

- temporarily relieves nasal congestion due to:
- common cold
- hay fever
- upper respiratory allergies
- temporarily relieves sinus congestion and pressure
- shrinks swollen nasal membranes so you can breathe more freely

Warnings

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland

When using this product

- **do not use more than directed**
- do not use for more than 3 days. Use only as directed. Frequent or prolonged use may cause nasal congestion to recur or worsen.
- temporary discomfort such as burning, stinging, sneezing or an increase in nasal discharge may occur
- use of this container by more than one person may spread infection

Stop use and ask a doctor if

symptoms persist

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

If swallowed get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- adults and children 6 to under 12 years of age (with adult supervision): 2 or 3 sprays in each nostril not more often than every 10 to 12 hours. Do not exceed 2 doses in any 24-hour period.
- children under 6 years of age: ask a doctor

To Use: Push firmly down on cap and turn counter clockwise. To spray, squeeze bottle quickly and firmly. Do not tilt head backward while spraying. Wipe nozzle clean after use. Secure cap after use.

Other information

- store at 20-25°C (68-77°F)
- retain carton for future reference on full labeling

Inactive ingredients

benzalkonium chloride solution, benzyl alcohol, dibasic sodium phosphate, edetate disodium, monobasic sodium phosphate, polyethylene glycol, povidone, propylene glycol, purified water

Questions or comments?

1-888-423-0139

Principal Display Panel

TopCare® health

COMPARE TO AFRIN® NASAL SPRAY ACTIVE INGREDIENT

Nasal Spray

NASAL DECONGESTANT

OXYMETAZOLINE HCl 0.05%

ORIGINAL


12 HOUR RELIEF

- Fast, Powerful Congestion Relief
- For Colds & Allergies

- Maximum Strength

1 FL OZ (30 mL)

NDC 36800-304-10



+
TopCare
health

COMPARE TO
 AFRIN® NASAL SPRAY
 ACTIVE INGREDIENT*

Nasal Spray

NASAL DECONGESTANT
 OXYMETAZOLINE HCl 0.05%

ORIGINAL

12
HOUR
RELIEF

DISTRIBUTED BY
 TOPCO ASSOCIATES LLC
 ELK GROVE VILLAGE, IL 60007
 ©TOPCO PERA0521
 QUESTIONS? 1-888-423-0139
 topcare@topco.com
 www.topcarebrand.com



Scan here for more
 information or
 call 1-888-423-0139

GF


**GLUTEN
FREE**

✓

**QUALITY
GUARANTEED**

This TopCare® product is laboratory
 tested to guarantee its highest quality.
 Your total satisfaction is guaranteed.

1 FL OZ (30 mL)





+
TopCare
health

Nasal Spray

NASAL DECONGESTANT
 OXYMETAZOLINE HCl 0.05%

ORIGINAL

12
HOUR
RELIEF

DISTRIBUTED BY
 TOPCO ASSOCIATES LLC
 ELK GROVE VILLAGE, IL 60007
 ©TOPCO PERA0521
 QUESTIONS? 1-888-423-0139
 topcare@topco.com
 www.topcarebrand.com



Scan here for more
 information or
 call 1-888-423-0139

GF

**GLUTEN
FREE**

✓

**QUALITY
GUARANTEED**

This TopCare® product is laboratory
 tested to guarantee its highest quality.
 Your total satisfaction is guaranteed.

DO NOT USE IF PRINTED NEG. READING
 IS BROKEN OR MISSING

Drug Facts	
Active ingredient Oxymetazoline hydrochloride 0.05%.....Nasal decongestant	Purpose
Uses <ul style="list-style-type: none"> ■ temporarily relieves nasal congestion due to: <ul style="list-style-type: none"> ■ common cold ■ hay fever ■ upper respiratory allergies ■ temporarily relieves sinus congestion and pressure ■ shrinks swollen nasal membranes so you can breathe more freely 	
Warnings Ask a doctor before use if you have <ul style="list-style-type: none"> ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ trouble urinating due to an enlarged prostate gland 	
When using this product <ul style="list-style-type: none"> ■ do not use more than directed ■ do not use for more than 3 days. Use only as directed. Frequent or prolonged use may cause nasal congestion to recur or worsen. ■ temporary discomfort such as burning, stinging, sneezing or an increase in nasal discharge may occur ■ use of this container by more than one person may spread infection 	
Stop use and ask a doctor if symptoms persist If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away (1-800-222-1222).	
Directions <ul style="list-style-type: none"> ■ adults and children 6 to under 12 years of age (with adult supervision): 2 or 3 sprays in each nostril not more often than every 10 to 12 hours. Do not exceed 2 doses in any 24-hour period. ■ children under 6 years of age: ask a doctor To Use: Push firmly down on cap and turn counter clockwise. To spray, squeeze bottle quickly and firmly. Do not tilt head backward while spraying. Wipe nozzle clean after use. Secure cap after use.	
Other information <ul style="list-style-type: none"> ■ store at 20-25°C (68-77°F) ■ retain carton for future reference on full labeling 	

Drug Facts (continued)

Inactive ingredients


benzalkonium chloride solution, benzyl alcohol, dibasic sodium phosphate, edetate disodium, monobasic sodium phosphate, polyethylene glycol, povidone, propylene glycol, purified water

Questions or comments?
 1-888-423-0139

*This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Afrin® Nasal Spray.

How to use:

TO USE: Push firmly down on cap and turn counter clockwise. Secure cap after use.

0 36800 13005 0

30410 88 C8

TOPCARE NASAL

oxymetazoline hydrochloride spray

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:36800-304
Route of Administration	NASAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OXYMETAZOLINE HYDROCHLORIDE (UNII: K89MJ0S5VY) (OXYMETAZOLINE - UNII:8VLN5B44ZY)	OXYMETAZOLINE HYDROCHLORIDE	0.05 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7)	
BENZYL ALCOHOL (UNII: LKG8494WBH)	
SODIUM PHOSPHATE, DIBASIC, UNSPECIFIED FORM (UNII: GR686LBA74)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
SODIUM PHOSPHATE, MONOBASIC, UNSPECIFIED FORM (UNII: 3980JIH2SW)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	

Product Characteristics

Color	WHITE (Translucent)	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-304-10	1 in 1 CARTON	12/15/1989	
1		30 mL in 1 BOTTLE, SPRAY; Type 0: Not a Combination Product		
2	NDC:36800-304-72	1 in 1 CARTON	03/17/2009	11/15/2017
2		37 mL in 1 BOTTLE, SPRAY; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/15/1989	

Revised: 12/2025

Topco Associates LLC