ACID REDUCER- ranitidine tablet
McKesson

Drug Facts

ACTIVE INGREDIENT (IN EACH TABLET)

Ranitidine 150 mg (as ranitidine hydrochloride, USP 168 mg)

PURPOSE

Acid reducer

USES

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by certain foods and beverages

WARNINGS

Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor
- with other acid reducers
- if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding,

Ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

DIRECTIONS

- adults and children 12 years and over:
  - to **relieve** symptoms, swallow 1 tablet with a glass of water
  - to **prevent** symptoms, swallow 1 tablet with a glass of water **30 to 60 minutes before** eating food or drinking beverages that cause heartburn
  - can be used up to twice daily (do not take more than 2 tablets in 24 hours)
- children under 12 years: ask a doctor

OTHER INFORMATION

- **TAMPER EVIDENT: DO NOT USE IF THE CARTON OR PRINTED FOIL UNDER CAP IS OPEN OR TORN.**
- store at 20° - 25° C (68° - 77° F)
- avoid excessive heat or humidity
- this product is sodium and sugar free

INACTIVE INGREDIENTS

Colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, talc, titanium dioxide

QUESTIONS?

call **1-800-406-7984**

PRINCIPAL DISPLAY PANEL

sunmark®

***COMPARE TO ZANTAC 150® ACTIVE INGREDIENT**

NDC 49348-026-04

Ranitidine Tablets, USP 150 mg

acid reducer

Prevents and Relieves Heartburn Associated with Acid Indigestion & Sour Stomach

Maximum Strength

24 TABLETS

Distributed By McKesson

5104606 / 0813


sunmark®
Ranitidine Tablets, USP 150 mg
acid reducer
sunmark®
*COMPARE TO ZANTAC 150® ACTIVE INGREDIENT
NDC 49348-026-04
Ranitidine Tablets, USP 150 mg
acid reducer
Prevents and Relieves
Heartburn Associated with
Acid Indigestion & Sour Stomach
Maximum Strength
24 TABLETS
sunmark®
Ranitidine Tablets, USP 150 mg
acid reducer
MADE IN INDIA
MCKESSON
Another Quality Product
Distributed By McKesson
One Post Street, San Francisco, CA 94104
Questions or comments?
Please visit us at www.commonwealbrand.com
0813
*All trademarks are property of their respective owners. This product is not affiliated with the makers/owners of Zantac 150®.
Read the directions, consumer information leaflet and warnings before use.
Keep the carton. It contains important information.
Expiration Date:
Batch No.
NON VARNISH
Uses
relieves heartburn associated with acid indigestion and sour stomach
prevents heartburn associated with acid indigestion and sour stomach brought on by certain foods and beverages
Warnings
Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers
Do not use
if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor
with other acid reducers
if you have kidney disease, except under the advice and supervision of a doctor
Ask a doctor before use if you have
frequent chest pain
frequent heartburn, particularly with heartburn
stomach pain
nausea or vomiting
unexplained weight loss
had heartburn over 3 months. This may be a sign of a more serious condition
heartburn with lightheadedness, sweating or dizziness
chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
Directions
adults and children 12 years and over:
to relieve symptoms, swallow 1 tablet with a glass of water
to prevent symptoms, swallow 1 tablet with a glass of water 30 to 60 minutes before eating food or drinking beverages that cause heartburn
can be used up to twice daily (do not take more than 2 tablets in 24 hours)
children under 12 years: ask a doctor
Other information
TAMPER EVIDENT: DO NOT USE IF THE CARTON OR PRINTED FOIL UNDER CAP IS OPEN OR TORN.
store at 20° - 25° C (68° - 77° F)
avoid excessive heat or humidity
this product is sodium and sugar free
Inactive ingredients
colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, talc, titanium dioxide
Questions? call 1-800-406-7984
5104606
5104606

ACID REDUCER

ranitidine tablet

Product Information			
Product Type	HUMAN OTC DRUG	**Item Code (Source)**	NDC:49348-026
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	**Basis of Strength**	**Strength**
RANITIDINE HYDROCHLORIDE (UNII: BK76465IHM) (RANITIDINE - UNII:884KT10YB7)	RANITIDINE	150 mg

Inactive Ingredients	
Ingredient Name	**Strength**
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics			
Color	pink	**Score**	no score
Shape	OVAL (Oval Shaped)	**Size**	12mm
Flavor		**Imprint Code**	9R
Contains			

Packaging				
#	**Item Code**	**Package Description**	**Marketing Start Date**	**Marketing End Date**
1	NDC:49348-026-04	1 in 1 CARTON		
1		24 in 1 BOTTLE		
2	NDC:49348-026-54	1 in 1 CARTON		
2		65 in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date

ANDA	ANDA200536	01/07/2014	

Labeler - McKesson (177667227)

Registrant - Ranbaxy Pharmaceuticals Inc. (937890044)

Establishment

Name	Address	ID/FEI	Business Operations
Shasun Pharmaceuticals Limited		915786829	manufacture(49348-026)

Revised: 1/2014 McKesson