

URINARY PAIN RELIEF- phenazopyridine hydrochloride tablet

YYBA CORP

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Drug Facts

Active ingredient (in each tablet)

Phenazopyridine Hydrochloride 99.5 mg

Purpose

Urinary tract analgesic

Uses

Relief from urinary pain, burning, urgency and frequency associated with urinary tract infections.

Warnings

Do not exceed recommended dosage.

Ask a doctor before use if you have

- kidney disease
- allergies to foods, preservatives or dyes
- had a hypersensitive reaction to Phenazopyridine Hydrochloride

Do not use

if you have Glucose-6-Phosphate Dehydrogenase (G6PD) deficiency unless approved by your physician

When using this product

- stomach upset may occur, taking this product with or after meals may reduce stomach upset
- your urine will become reddish-orange in color. This is not harmful, but care should be taken to avoid staining clothing or other items.

Stop use and ask a doctor if

- your symptoms last for more than 2 days
- you suspect you are having an adverse reaction to the medication
- long-term administration of phenazopyridine hydrochloride has induced neoplasia in rats (large intestine) and mice (liver). Although no association between

phenazopyridine hydrochloride and human neoplasia has been reported, adequate epidemiological studies along these lines have not been conducted.

If pregnant or breastfeeding,

ask a health professional before use. **A pregnancy test and consultation with a health professional if pregnancy is confirmed is recommended prior to use.**

Keep out of reach of children.

In case of an overdose, get medical help or contact a Poison Control Center call (1-800-222-1222) right away.

Directions

Adults and children 12 years and older: Take 2 tablets 3 times daily with or after meals as needed for up to two days. Take with a full glass of water. Do not use for more than 2 days (12 tablets) without consulting a doctor

Children under 12: Do not use without consulting a doctor

Other information

- This product can interfere with laboratory tests including urine, glucose (sugar), and ketones tests
- This product may stain contact lenses and other items if handled after touching tablets.
- Store at room temperature (59°-86°F) in a dry place and protect from light

Inactive ingredients

colloidal silicone dioxide, croscarmellose sodium, hydroxypropyl methylcellulose, magnesium stearate, maize (corn) starch, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch

Questions?

call toll-free **1-866-933-6337**

PRINCIPAL DISPLAY PANEL

3 73581 00063

Welmate

100% SATISFACTION GUARANTEED

URINARY PAIN RELIEF

MAX STRENGTH OTC

Phenazopyridine HCl 99.5 mg

FAST ACTING

- ✓ Urinary Analgesic
- ✓ Targets Pain Source
- ✓ Relieves Urinary Pain, Burning & Urgency

IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED

Facts

Ingredient Phenazopyridine Hydrochloride 99.5 mg

Purpose Urinary tract analgesic

Do not exceed recommended dosage. Do not use if you have kidney allergies to foods, preservatives, or dyes hypersensitive reaction to Phenazopyridine or Glucose-6-Phosphate deficiency unless by your physician

Peel for more information



WARNING: This product contains phenazopyridine hydrochloride, which is known to cause cancer. For more information go to www.P65Wt.com.

welmate®

URINARY PAIN RELIEF

MAX STRENGTH OTC*

100% SATISFACTION GUARANTEED

FAST ACTING



72 TABLETS

Drug Facts (continued)

- Stop use and ask a doctor if**
- your symptoms last for more than 2 days
 - you suspect you are having an adverse reaction to the medication
- Long-term administration of phenazopyridine hydrochloride has induced neoplasia in rats (large intestine) and mice (liver). Although no association between phenazopyridine hydrochloride and human neoplasia has been reported, adequate epidemiological studies along these lines have not been conducted.
- If pregnant or breast-feeding,** ask a healthcare professional before use. **A pregnancy test and consultation with a healthcare professional if pregnancy is confirmed is recommended prior to use.**
- Keep out of reach of children.** In case of an overdose, get medical help or contact a Poison Control Center call (1-800-222-1222) right away.
- Directions**
- Adults and children 12 years and older: Take 2 tablets 3 times daily with or after meals as needed for up to two days. Take with a full glass of water. Do not use for more than 2 days (12 tablets) without consulting a doctor.
 - Children under 12: Do not use without consulting a doctor.

Drug Facts

- Purpose**
- Phenazopyridine Hydrochloride 99.5 mg.....Urinary tract analgesic
- Use** Relief from urinary pain, burning, urgency, and frequency associated with urinary tract infections.
- Warnings**
- Do not exceed recommended dosage.**
- Ask a doctor before use if you have** ■ kidney disease ■ allergies to foods, preservatives, or dyes ■ had a hypersensitive reaction to Phenazopyridine Hydrochloride
- Do not use if you have** Glucose-6-Phosphate Dehydrogenase (G6PD) deficiency unless approved by your physician
- When using this product**
- stomach upset may occur, taking this product with or after meals may reduce stomach upset
 - your urine will become reddish-orange in color. This is not harmful, but care should be taken to avoid staining clothing or other items.

*Maximum strength without a prescription

Distributed by: Wellspring, Airmont, NY 10952 U.S.A.

PEEL FOR MORE INFORMATION



WARNING: This product contains phenazopyridine hydrochloride, which is known to cause cancer. For more information go to www.P65Wt.com.

welmate®

URINARY PAIN RELIEF

MAX STRENGTH OTC*

100% SATISFACTION GUARANTEED

FAST ACTING



72 TABLETS

Drug Facts (continued)

- Stop use and ask a doctor if**
- your symptoms last for more than 2 days
 - you suspect you are having an adverse reaction to the medication
- Long-term administration of phenazopyridine hydrochloride has induced neoplasia in rats (large intestine) and mice (liver). Although no association between phenazopyridine hydrochloride and human neoplasia has been reported, adequate epidemiological studies along these lines have not been conducted.
- If pregnant or breast-feeding,** ask a healthcare professional before use. **A pregnancy test and consultation with a healthcare professional if pregnancy is confirmed is recommended prior to use.**
- Keep out of reach of children.** In case of an overdose, get medical help or contact a Poison Control Center call (1-800-222-1222) right away.
- Directions**
- Adults and children 12 years and older: Take 2 tablets 3 times daily with or after meals as needed for up to two days. Take with a full glass of water. Do not use for more than 2 days (12 tablets) without consulting a doctor.
 - Children under 12: Do not use without consulting a doctor.

Drug Facts

- Purpose**
- Phenazopyridine Hydrochloride 99.5 mg.....Urinary tract analgesic
- Use** Relief from urinary pain, burning, urgency, and frequency associated with urinary tract infections.
- Warnings**
- Do not exceed recommended dosage.**
- Ask a doctor before use if you have** ■ kidney disease ■ allergies to foods, preservatives, or dyes ■ had a hypersensitive reaction to Phenazopyridine Hydrochloride
- Do not use if you have** Glucose-6-Phosphate Dehydrogenase (G6PD) deficiency unless approved by your physician
- When using this product**
- stomach upset may occur, taking this product with or after meals may reduce stomach upset
 - your urine will become reddish-orange in color. This is not harmful, but care should be taken to avoid staining clothing or other items.

*Maximum strength without a prescription

Distributed by: Wellspring, Airmont, NY 10952 U.S.A.

PEEL FOR MORE INFORMATION

chemicals
is known
to cause
cancer

36 TABLETS



DO NOT USE IF

Drug

Active (in each Phenazopyridine Hydrochloride)

Use Relief from urinary pain, burning, urgency, and frequency associated with urinary tract infections.

Warnings

Do not exceed recommended dosage.

Ask a doctor before use if you have ■ kidney disease ■ allergies to foods, preservatives, or dyes ■ had a hypersensitive reaction to Phenazopyridine Hydrochloride

Do not use if you have Glucose-6-Phosphate Dehydrogenase (G6PD) deficiency unless approved by your physician

When using this product

- stomach upset may occur, taking this product with or after meals may reduce stomach upset
- your urine will become reddish-orange in color. This is not harmful, but care should be taken to avoid staining clothing or other items.

PEEL HERE

7 0004

100% Satisfaction Guarantee
 Return for a full refund if you are not completely satisfied with your purchase.

Phenazopyridine HCl 99.5 mg

- ✓ Urinary Analgesic
- ✓ Targets Pain Source
- ✓ Relieves Urinary Pain, Burning & Urgency

VALUE SIZE
120
 TABLETS

RECOMMENDED BY
 HEALTHCARE PROFESSIONALS

DO NOT USE IF PRINTED SEAL IS BROKEN

Drug Facts
Active ingredient (in each tablet)
 Phenazopyridine Hydrochloride

Use
 Relief from urinary pain and frequency associated with urinary tract infections.

Warnings
 Do not exceed recommended dose. Ask a doctor before use if you have a history of liver disease, kidney disease, or allergies to food or drugs. Do not use if you are pregnant or breastfeeding. Do not use if you have a history of urinary tract infections.

Do not use if you have a history of urinary tract infections.

When using this product
 • stomach upset may occur with or after meals may occur
 • your urine will become orange
 • This is not harmful, but to avoid staining clothing or toilet paper, avoid staining clothing or toilet paper.

PEEL FOIL HERE

Drug Facts (continued)

Stop use and ask a doctor if

- your symptoms last for more than 2 days
- you suspect you are having an adverse reaction to the medication

Long-term administration of phenazopyridine hydrochloride has induced neoplasia in rats (large intestine) and mice (liver). Although no association between phenazopyridine hydrochloride and human neoplasia has been reported, adequate epidemiological studies along these lines have not been conducted.

If pregnant or breast-feeding, ask a healthcare professional before use. A pregnancy test and consultation with a healthcare professional if pregnancy is confirmed is recommended prior to use.

Keep out of reach of children. In case of an overdose, get medical help or contact a Poison Control Center call (1-800-222-1222) right away.

Directions

- Adults and children 12 years and older:** Take 2 tablets 3 times daily with or after meals as needed for up to two days. Take with a full glass of water. Do not use for more than 2 days (12 tablets) without consulting a doctor.
- Children under 12:** Do not use without consulting a doctor.

Other information

- This product can interfere with laboratory tests including urine, glucose (sugar), and ketones tests
- This product may stain soft contact lenses and other items if handled after touching tablets
- Store at room temperature (59°-86°F) in a dry place and protect from light

Inactive ingredients
 colloidal silicon dioxide, croscarmellose sodium, hydroxypropyl methylcellulose, magnesium stearate, maize (corn) starch, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch

Questions? call toll-free 1-866-933-6337

*Maximum strength without a prescription
 Distributed by: WellSpring, Airmont, NY 10952 U.S.A

URINARY PAIN RELIEF

phenazopyridine hydrochloride tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:73581-919
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PHENAZOPYRIDINE HYDROCHLORIDE (UNII: 0EWG668W17) (PHENAZOPYRIDINE - UNII:K2J09EMJ52)	PHENAZOPYRIDINE HYDROCHLORIDE	99.5 mg

Inactive Ingredients

Ingredient Name	Strength
MAGNESIUM STEARATE (UNII: 70097M6I30)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
STARCH, CORN (UNII: O8232NY3SJ)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POVIDONE K30 (UNII: U725QWY32X)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	

Product Characteristics

Color	brown	Score	no score
Shape	ROUND	Size	7mm
Flavor		Imprint Code	WM7
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73581-919-72	72 in 1 BOTTLE; Type 0: Not a Combination Product	04/01/2026	
2	NDC:73581-919-36	36 in 1 BOTTLE; Type 0: Not a Combination Product	04/01/2026	
3	NDC:73581-919-12	120 in 1 BOTTLE; Type 0: Not a Combination Product	04/01/2026	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		04/01/2026	

Labeler - YYBA CORP (006339772)

Registrant - YYBA CORP (006339772)

Revised: 4/2026

YYBA CORP