

NOCOLD-S- acetaminophen, chlorpheniramine maleate liquid
Lydia Co., Ltd.

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Acetaminophen, Chlorpheniramine Maleate

temporarily relieves thee symptoms due to cold or flu

minor aches and pains headache, nasal congestion, runny nose, cough, sneezing, sore throat

Keep out of reach of children

■ Adult and children 16 years and over: take 1 bottle (20ml) per session 3 times per day after meal

see product label

sodium citrate dihydrate, L-sodium glutamate, sugar, d-sorbitol solution, distilled water for oral use

Comprehensive cold medicine

Han Shin

NOCOLD-S LIQUID

runny nose /sneezing /body aches

Drug Facts

Active Ingredient(in each bottle - 20ml / 0.676oz) **Purpose**

Acetaminophen 200 mg -----Pain reliever/ fever reducer

Chlorpheniramine maleate 2.5 mg -----Antihistamine

Uses

temporarily relieves these symptoms due to a cold or flu · minor aches and pains · headache · nasal congestion · runny nose · cough · sneezing · sore throat

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if · adult takes more than 3 doses in 24 hours, which is the maximum daily amount · child takes more than 1 dose in 24 hours, which is the maximum daily amount · taken with other drugs containing acetaminophen · adult has 3 or more alcoholic drinks every day while using this product

Do not use

· with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

· exceed recommended dosage and administration.

· with the following medication: Antitussive expectorants, other cold medication, antipyretic and analgesics, antihistamine drugs, sedatives, etc.

Ask a doctor before use if the user has liver disease

Ask a doctor or pharmacist before use if the user have

· liver disease · thyroid disease · diabetes · hypertension · weak constitution or high fever · heart disease or the elderly · glaucoma (e.g., eye pain or dimmed vision, etc.) · parents or siblings who are susceptible to urticaria, contact dermatitis, bronchial asthma, allergic rhinitis, migraines, food allergies, etc.

· previously experienced drug allergies (e.g., fever, rash, joint pain, asthma, skin irritation, etc.) · varicella or influenza or infants and children under 14 years of age who are suspected of varicella or influenza · difficulty in urination · edema

Ask a doctor or pharmacist before use (for children and infants)

· Children should take this medication under the guidance of their parents.

· Infants under 2 years old should be given treatment by a doctor and should not take this medication unless it is absolutely necessary; infants under 2 years old should take this medication with the guardian's awareness and should be monitored closely.

Stop use and ask a doctor if

· occurrence of reduced urine output, swelling of the face or limbs, heavy eyelids, stiffness of hands, high blood pressure or headache, and other symptoms after taking this product.

· asthma

· Pseudo aldosterone disease: When individuals are in long-term use of the preparation of which the daily dose of licorice is over 1g, symptoms of pseudoaldosteronism, such as hypokalemia, increased blood pressure, sodium and water retention, edema, and weight gain, may occur. After a full observaion (e.g., determination of serum potassium concentration),

Overdose warning

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away.

Store at cool temperature and dry place with a closed container.

Avoid direct sunlight.- Store in a container other than its original container is equivalent to misuse.

Directions

· do not take more than the recommended dose

· adults and children 16 years and over: take 1 bottle (20ml) per session 3 times per day after meal.

Inactive Ingredients

Sodium Citrate Dihydrate, L-Sodium Glutamate, Sugar, D-Sorbitol Solution, Distilled Water

Distributed by

Bio Mission Group Inc.

9925 Painter Ave #0,Whittier, CA90605, 562-273-0692

Manufactured by

HanKook ShinYak Pharmaceutical Co.,Ltd.

39-83, Jungsan-gil, Yangcheon-myeon, Nonsan, Chungcheongnam-do,320850, South Korea

Question or Comments

Call +1-82-41-740-8900

**If you have purchased an expired, altered, deteriorated, or contaminated product, it can be exchanged ate the place of purchase.*

NOCOLD-S					
acetaminophen, chlorpheniramine maleate liquid					
Product Information					
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72988-0033		
Route of Administration	ORAL				
Active Ingredient/Active Moiety					
Ingredient Name		Basis of Strength	Strength		
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE -		CHLORPHENIRAMINE	2.5 mg		

UNII:3U6IO1965U)	MALEATE	in 20 mL
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	200 mg in 20 mL

Inactive Ingredients

Ingredient Name	Strength
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72988-0033-1	20 mL in 1 BOTTLE; Type 0: Not a Combination Product	01/23/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		01/23/2023	

Labeler - Lydia Co., Ltd. (695735569)

Registrant - Lydia Co., Ltd. (695735569)

Establishment

Name	Address	ID/FEI	Business Operations
Lydia Co., Ltd.		695735569	manufacture(72988-0033)