

**PAIN RELIEF- acetaminophen tablet
BETTER LIVING BRANDS LLC.**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

341R_Albertsons_21130-431_ACETAMINOPHEN CAPLETS 500 mg

Drug Facts

Active ingredient (in each caplet)

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Uses

temporarily relieves minor aches and pains due to:

- the common cold
- headache
- backache
- minor pain of arthritis
- toothache
- muscular aches
- premenstrual and menstrual cramps

temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Do not take more than directed.

See overdose warning

adults and children 12 years and over	<ul style="list-style-type: none">• take 2 caplets every 6 hours while symptoms last• do not take more than 6 caplets in 24 hours, unless directed by a doctor• do not take for more than 10 days unless directed by a doctor
children under 12 years	ask a doctor

Other information

- SODIUM FREE
- store between 20-25°C (68-77°F) excursions permitted between 15-30°C (59-86°F)
- use by expiration date on package

Inactive ingredients

hypromellose, mineral oil, povidone, pregelatinized starch, sodium starch glycolate*, stearic acid, titanium dioxide

*may contain this ingredient

Questions or comments? 1-877-290-4008

Adhesive Area

Drug Facts (continued)

Keep out of reach of children.
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions Do not take more than directed. See overdose warning **adults and children 12 years and over:**

- take 2 caplets every 6 hours while symptoms last
- do not take more than 6 caplets in 24 hours, unless directed by a doctor
- do not take for more than 10 days unless directed by a doctor

children under 12 years: ■ ask a doctor unless directed by a doctor

Other information ■ **SODIUM-FREE**

- store between 20-25°C (68-77°F)
- excursions permitted between 15-30°C (59-86°F)
- use by expiration date on package

Questions or comments?

1-877-290-4008
 DIS TRIBUTED BY:
 BETTER LIVING BRANDS, LLC
 P.O. BOX 99
 PLEASANTON, CA 94566-0099
 41-888-723-3929
 341R 0223
 RD 23055
 35354

howZreycycleInfo

PLASTIC BOTTLE

LOVE IT OR IT'S ON US.®



NDC 21130-431-42

Quality Guaranteed

Extra Strength
Pain Relief
ACETAMINOPHEN
 CAPLETS 500 mg
 Pain Reliever/Fever Reducer
 For Adults **24 CAPLETS**

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Drug Facts

Active ingredient (in each caplet)

Acetaminophen 500 mg.....Pain reliever/fever reducer

Uses ■ temporarily relieves minor aches and pains due to: the common cold, headache, backache, minor pain of arthritis, toothache, muscular aches, premenstrual and menstrual cramps

■ temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product. **Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

If pregnant or breast-feeding, ask a health professional before use.

Drug Facts (continued under label)

Varnish Omit Area

Lot No.:
Exp Date:



Exp Date: _____
Lot No.: _____
Varnish Omit



Print/Varnish Omit Area

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts

Active ingredient (in each caplet) Purpose
Acetaminophen 500 mg... Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - the common cold
 - headache
 - backache
 - minor pain of arthritis
 - toothache
 - muscular aches
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - new symptoms occur
 - redness or swelling is present
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Do not take more than directed. See overdose warning

Drug Facts (continued)

- | | |
|--|---|
| adults and children 12 years and over | <ul style="list-style-type: none">take 2 caplets every 6 hours while symptoms lastdo not take more than 6 caplets in 24 hours, unless directed by a doctordo not take for more than 10 days unless directed by a doctor |
| children under 12 years | ask a doctor |

Other information

- SODIUM FREE
- store between 20-25°C (68-77°F) excursions permitted between 15-30°C (59-86°F)
- use by expiration date on package

Inactive ingredients hypromellose, mineral oil, povidone, pregelatinized starch, sodium starch glycolate*, stearic acid, titanium dioxide *may contain this ingredient

Questions or comments? 1-877-290-4008

This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division, owner of the registered trademark Tylenol®.
DISTRIBUTED BY: BETTER LIVING BRANDS LLC
P.O. BOX 99, PLEASANTON, CA 94566-0099
†1-888-723-3929
341R 0223

Compare to Extra Strength Tylenol® active ingredient NDC 27030-481-42

Actual Size

24 CAPLETS

Extra Strength Pain Relief

ACETAMINOPHEN CAPLETS 500 mg

Pain Reliever/Fever Reducer

For Adults



826 E Varnish/Print Omit Area

Print/Varnish Omit Area



Varnish Omit

Lot No:
Exp Date:



RD 23054 5S954



READ AND KEEP CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION
TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts

Active ingredient (in each caplet) **Purpose**
Acetaminophen 500 mg Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - the common cold
 - headache
 - backache
 - minor pain of arthritis
 - toothache
 - muscular aches
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

 If a skin reaction occurs, stop use and seek medical help right away.

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Drug Facts (continued)

■ if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - new symptoms occur
 - redness or swelling is present
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Do not take more than directed.
See overdose warning

- | | |
|--|---|
| adults and children 12 years and over | <ul style="list-style-type: none"> take 2 caplets every 6 hours while symptoms last do not take more than 6 caplets in 24 hours, unless directed by a doctor do not take for more than 10 days unless directed by a doctor |
| children under 12 years | ask a doctor |

Drug Facts (continued)

Other information

- SODIUM FREE**
- store between 20-25°C (68-77°F) excursions permitted between 15-30°C (59-86°F)
- use by expiration date on package

Inactive ingredients hypromellose, mineral oil, povidone, pregelatinized starch, sodium starch glycolate*, stearic acid, titanium dioxide *may contain this ingredient

Questions or comments? 1-877-290-4000

DISTRIBUTED BY:
BETTER LIVING BRANDS LLC
P.O. BOX 99
PLEASANTON, CA 94566-0099
11-888-723-3929
341R 0223

Varnish Omit

Compare to Extra Strength Tylenol[®] active ingredients

NDC 211301-431-05

Extra Strength Pain Relief
ACETAMINOPHEN CAPLETS
500 mg
Pain Reliever/Fever Reducer

Actual Size



50 CAPLETS

• For adults



Quality Guaranteed

Varnish Omit

E827R

Varnish/Print Omit Area





NDC 21130-431-05

Quality
Guaranteed

Extra Strength Pain Relief ACETAMINOPHEN CAPLETS 500 mg

Pain Reliever/Fever Reducer

- For adults

50 CAPLETS

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION
TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER
THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING**

Drug Facts

Active ingredient (in each caplet)

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Uses ■ temporarily relieves minor aches and pains due to: the common cold, headache, backache, minor pain of arthritis, toothache, muscular aches, menstrual and menstrual cramps
■ temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product. **Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash if a skin reaction occurs, stop use and seek medical help right away. If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.**

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions: Do not take more than directed. See overdose warning adults and children 12 years and over. ■ take 2 caplets every 6 hours while symptoms last ■ do not take more than 6 caplets in 24 hours, unless directed by a doctor ■ do not take for more than 10 days unless directed by a doctor

Other information ■ SODIUM FREE ■ store between 20-25°C (68-77°F) excursions permitted between 15-30°C (59-86°F) ■ use by expiration date on package

Questions or comments? 1-877-290-4008

**LOVE IT
OR IT'S ON US:**

DISTRIBUTED BY: BETTER LIVING BRANDS LLC
P.O. BOX 99, PLEASANTON, CA 94566-0099
1-888-723-3929

RD 23054
SS954

341R 0223

Lot No.:

Exp. Date:



UNVARNISHED AREA

Coating/Print Omit Area



READ AND KEEP CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION
TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING.

Drug Facts

Active ingredient *Purpose*
(in each caplet)
Acetaminophen 500 mg Pain reliever/fever reducer

- Uses**
- temporarily relieves minor aches and pains due to:
 - the common cold
 - headache
 - backache
 - minor pain of arthritis
 - toothache
 - muscular aches
 - premenstrual and menstrual cramps
 - temporarily reduces fever

Warnings
Lower warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Do not use

- with any other drug containing acetaminophen

Drug Facts (continued)

(prescription or nonprescription) If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
If you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 5 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions Do not take more than directed.
See overdose warning.
adults and children 12 years and over:

- take 2 caplets every 6 hours while symptoms last
- do not take more than 6 caplets in 24 hours, unless directed by a doctor
- do not take for more than

Drug Facts (continued)

10 days unless directed by a doctor
children under 12 years ask a doctor

Other information

- SODIUM FREE
- store between 20-25°C (68-77°F) excursions permitted between 15-30°C (59-86°F)
- use by expiration date on package

Inactive ingredients
hydroxypropylcellulose, mineral oil, polydioxane, pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide
*may contain this ingredient

Questions or comments? 1-877-260-4006

*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division, owner of the registered trademark Tylenol®.
DISTRIBUTED BY: BETTER LIVING BRANDS LLC
P.O. BOX 96, PLEASANTON, CA 94566-0099
1-888-723-3829

341R 0223

Coating Omit

Compare to Extra Strength Tylenol® active ingredient*
NDC 2180-431-01
Aspirin

Extra Strength Pain Relief
ACETAMINOPHEN
CAPLETS 500 mg
Pain Reliever/Fever Reducer
• For Adults



100 CAPLETS

E1323R

Coating/Print Omit Area

Coating Omit





NDC 21190-431-01

Quality
Guaranteed

Extra Strength
Pain Relief
ACETAMINOPHEN
CAPLETS 500 mg

Pain Reliever/Fever Reducer

• For Adults

100 CAPLETS

THIS PRODUCT DOES NOT CONTAIN ANY DRUGS THAT ARE KNOWN TO BE HARMSFUL TO SEAL OR OPEN THE RESIN OR PLASTIC BOTTLE OR CAP, OR TO CAUSE IRRITATION TO THE SKIN OR EYES.
NEED AND KEEP CAPLETS FROM COMPLETE WARMINGS AND PRODUCT INFORMATION

Drug Facts

Active ingredient (in each caplet): Acetaminophen 500 mg
Purpose: Pain reliever/fever reducer
Uses: Temporary relief of minor aches and pains due to: common cold, headache, backache, minor pain of arthritis, toothache, muscular aches, premenstrual and menstrual cramps
Temporarily reduces fever.

Warnings:
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours or with other drugs containing acetaminophen. 3 or more alcoholic drinks every day while using this product. Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening, blisters, rash.

If a skin reaction occurs, stop use and seek medical help right away. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Avoid use if you are taking other acetaminophen products. For more information, call 1-800-222-1222.

Directions: Do not take more than directed. See overdose warning. adults and children 12 years and over: Take 2 caplets every 6 hours while symptoms last. Do not take more than 6 caplets in 24 hours, unless directed by a doctor. Do not take for more than 10 days unless directed by a doctor.

Other information: SODIUM FREE. Store between 20°-25°C (68°-77°F); excursions permitted between 15°-30°C (59°-86°F). Use by expiration date on package. **Questions or comments? 1-877-236-1088**

DISTRIBUTED BY: **BEAUMONT PHARMAS LLC**
P.O. BOX 69
11-388723-0003
3418 0223
89664

LOVE IT
OR IT'S ON US
RD 22955
api-pharm.com

Lot No.: _____
Exp. Date: _____



Varnish Omit Area



Quality Guaranteed

Compare to Extra Strength Tylenol[®] Active Ingredient
NDC 2130-431-15

Extra Strength Pain Relief ACETAMINOPHEN CAPLETS 500 mg
Pain Reliever/
Fever Reducer
• For Adults
Actual Size
150 CAPLETS

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts

Active ingredient (in each caplet) Acetaminophen 500 mg.....Pain reliever/fever reducer

Uses
temporarily relieves minor aches and pains due to:
■ headache ■ fever ■ cold ■ sore throat ■ toothache
■ minor aches ■ backache ■ menstrual cramps
■ temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
■ more than 4,000 mg of acetaminophen in 24 hours
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product
Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin redness ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

Drug Facts (continued under label)
*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., Medical Consumer Healthcare Division, owner of the registered trademark "Tylenol".
DISTRIBUTED BY: BETTERLIVING BRANDS, LLC
P.O. BOX 39, PLEASANTON, CA 94658-0039
1-888-727-3828

how2recycle.info
PLASTIC BOTTLE
Scan here for more information
RD 230656
LOT 1818101111
34FR10223
Lot No.:
Exp. Date:

LOVE IT OR IT'S ON US!
Varnish Omit Area

Adhesive Area

Drug Facts (continued)
Do not use
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if you are allergic to acetaminophen or any of the inactive ingredients in this product
Ask a doctor before use if you have liver disease
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin
Stop use and ask a doctor if
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ new symptoms occur
■ redness or swelling is present
These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Directions: Do not take more than directed.
See overdose warning
adults and children 12 years and over
■ take 2 caplets every 6 hours while symptoms last
■ do not take more than 6 caplets in 24 hours, unless directed by a doctor
■ do not take for more than 10 days unless directed by a doctor
children under 12 years
ask a doctor
Other information
■ SO DIUM FREE
■ store between 20°-25°C (68°-77°F); excursions permitted between 15°-30°C (59°-86°F)
■ use by expiration date on package
Inactive ingredients: hypromellose, mineral oil, polydioxane, pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide, may contain this ingredient
Questions or comments? 1-877-290-4008



Compare to
Extra Strength
Tylenol[®]
active ingredient*
NDC 21130-431-50

Quality
Guaranteed

Extra Strength Pain Relief

**ACETAMINOPHEN
CAPLETS 500 mg**

Pain Reliever/Fever Reducer
• For Adults

Actual Size



500 CAPLETS

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts

Active ingredient (in each caplet) Purpose
Acetaminophen 500 mg.....Pain reliever/
fever reducer

Uses ■ temporarily relieves minor aches and pains due to: ■ the common cold ■ headache
■ backache ■ minor pain of arthritis
■ toothache ■ muscular aches
■ premenstrual and menstrual cramps
■ temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

Do not use ■ with any other drug containing

Drug Facts (continued)

acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ new symptoms occur
■ redness or swelling is present
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions Do not take more than directed.

See overdose warning

adults and children 12 years and over

■ take 2 caplets every 6 hours while symptoms last
■ do not take more than 6 caplets in 24 hours, unless directed by a doctor
■ do not take for more than 10 days unless directed by a doctor

children under 12 years

ask a doctor

Other information

■ **SODIUM FREE** ■ store between 20-25°C (68-77°F) excursions permitted between 15-30°C (59-86°F) ■ use by expiration date on package

Inactive ingredients

hypromellose, mineral oil, povidone, pregelatinized starch, sodium starch glycolate*, stearic acid, titanium dioxide
*may contain this ingredient

Questions or comments? 1-877-290-4008

LOVE IT OR IT'S ON US!

This product is not
distributed by:
DISTRIBUTED BY: BETTER LIVING BRANDS LLC
14000 BUCKINGHAM PLACE, SIMONSON, CA 94596
1-800-729-3829
© 2011
Scan the info



how2recycle.info

PLASTIC BOTTLE

50554

FD 20852

Lot No.:

Exp Date:



21130177743 4

UNWARRANTED AREA

PAIN RELIEF

acetaminophen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:21130-431
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
POVIDONE (UNII: FZ989GH94E)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
STARCH, CORN (UNII: O8232NY3SJ)	
TITANIUM DIOXIDE (UNII: 15F1X9V2JP)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MINERAL OIL (UNII: T5L8T28FGP)	

Product Characteristics

Color	white	Score	no score
Shape	CAPSULE (Capsule shaped tablet)	Size	18mm
Flavor		Imprint Code	TCL341
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-431-42	1 in 1 CARTON	06/01/2023	
1		24 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:21130-431-50	500 in 1 BOTTLE; Type 0: Not a Combination Product	06/01/2023	
3	NDC:21130-431-15	150 in 1 BOTTLE; Type 0: Not a Combination Product	06/01/2023	
4	NDC:21130-431-25	1 in 1 CARTON	06/01/2023	
4		250 in 1 BOTTLE; Type 0: Not a Combination Product		
5	NDC:21130-431-05	1 in 1 CARTON	06/22/2023	
5		50 in 1 BOTTLE; Type 0: Not a Combination Product		
6	NDC:21130-431-01	1 in 1 CARTON	06/22/2023	
6		100 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	06/01/2023	

Labeler - BETTER LIVING BRANDS LLC. (009137209)

Registrant - TIME CAP LABORATORIES, INC. (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
TIME CAP LABORATORIES, INC.		037052099	manufacture(21130-431)