

PAIN RELIEF- acetaminophen tablet
BETTER LIVING BRANDS LLC.

340R_Albertsons_21130-043_Pain Relief Acetaminophen Tablets 325mg

Drug Facts

Active ingredient (in each tablet)

Acetaminophen 325 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:

- headache
- muscular aches
- backache
- minor pain of arthritis
- the common cold
- toothache
- premenstrual and menstrual cramps

- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 10 tablets (3,250 mg) in 24 hours for adults or 5 tablets (1,625 mg) in 24 hours for children. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours, which is the maximum daily amount
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if the user has liver disease

Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days in adults
- pain gets worse or lasts more than 5 days in children under 12 years
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Do not take more than directed.

See overdose warning

adults and children 12 years and over	<ul style="list-style-type: none">• take 2 tablets every 4 to 6 hours while symptoms last• do not take more than 10 tablets in 24 hours, unless directed by a doctor• do not use for more than 10 days unless directed by a doctor
children 6 years to under 12 years	<ul style="list-style-type: none">• take 1 tablet every 4 to 6 hours while symptoms last• do not take more than 5 tablets in 24 hours• do not use for more than 5 days unless directed by a doctor
children under 6 years	ask a doctor

Other information

- SODIUM FREE
- store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients povidone, pregelatinized starch, sodium starch glycolate*, stearic acid

*may contain this ingredient

Questions or comments? 1-877-290-4008

Side of Package

Coating/Print Omit Area



Lot No.: 211301160333
 Exp Date: 3/2024

-Bottom of Package-

-Back of Package-

-Top of Package-

READ AND KEEP CAREFUL FOR COMPLETE WARNINGS AND PRECAUTION INFORMATION. TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, OR OPEN ON INSPECTION.

Drug Facts

Active Ingredient (in each tablet) Purpose
Acetaminophen 325 mg.....Pain reliever/fever reducer

- Uses**
- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - backache
 - minor pain of arthritis
 - the common cold
 - toothache
 - premenstrual and menstrual cramps
 - temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 10 tablets (3,250 mg) in 24 hours for adults or 5 tablets (1,625 mg) in 24 hours for children. Severe liver damage may occur if:
 • adult takes more than 4,000 mg of acetaminophen in 24 hours
 • child takes more than 5 doses in 24 hours, which is the maximum daily amount
 • taken with other drugs containing acetaminophen
 • adult has 3 or more alcoholic drinks every day while using this product
Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:
 • skin redness • hives • rash
 If a skin reaction occurs, stop use and seek medical help right away.

Do not use
 • with any other drug containing acetaminophen

Drug Facts (continued)

(prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

If you are allergic to acetaminophen or any of the inactive ingredients in this product, ask a doctor before use. If the user has their disease, ask a doctor or pharmacist before use. If the user is taking the blood thinning drug warfarin, stop use and ask a doctor if:
• pain gets worse or lasts more than 10 days in adults
• pain gets worse or lasts more than 5 days in children under 12 years
• fever gets worse or lasts more than 3 days
• new symptoms occur • redness or swelling to present These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.
Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). **Get medical attention or call for adults as well as for children even if you do not notice any signs or symptoms.**

Directions
• Do not take more than directed. See overdose warning.
adults and children 12 years and over
 • take 2 tablets every 4 to 6 hours while symptoms last
 • do not take more than 10 tablets in 24 hours, unless directed by a doctor
 • do not use for more than 10 days unless directed by a doctor

Drug Facts (continued)

children 6 years to under 12 years
 • take 1 tablet every 4 to 6 hours while symptoms last
 • do not take more than 5 tablets in 24 hours
 • do not use for more than 5 days unless directed by a doctor
 • ask a doctor

Other Information
• SODIUM FREE
 • Store at 25°C (77°F) excursions permitted between 15°-30°C (59°-86°F)
 • Use by a date other than on package

Inactive ingredients: polydiox, pregelatinized starch, sodium starch glycolate*, stearic acid *may contain this ingredient

Questions or comments? 1-877-296-4008

*This product is not manufactured or distributed by Johnson & Johnson Consumer, Inc., McNeil Consumer Healthcare Division, owner of the registered trademark Tylenol® Regular Strength.

DISTRIBUTED BY: BETTER LIVING BRANDS LLC
P.O. BOX 90, PLEASANTON, CA 94566-0090
†1-888-723-3029 340R 0823

Coating Omit

Compare to Regular Strength Tylenol® active ingredients! NDC 3750-040-01

Regular Strength | Aspirin Free
Pain Relief

Actual Size
ACETAMINOPHEN 325 mg
 Pain Reliever/Fever Reducer
 • Contains no aspirin

100 TABLETS



E1323R

Coating/Print Omit Area

Coating Omit



1/8" Print Bleed

Side of Package



NDC 21130-043-01

Quality
GuaranteedRegular Strength | Aspirin Free
Pain Relief**ACETAMINOPHEN**
325 mg

Pain Reliever/Fever Reducer

- Contains no aspirin

100 TABLETS

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION
TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL
OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts**Active ingredient (in each tablet)**

Acetaminophen 325 mg.....Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to: headache, muscular aches, backache, minor pain of arthritis, the common cold, toothache, premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

- **Liver warning:** This product contains acetaminophen. The maximum daily dose of this product is 10 tablets (3,250 mg) in 24 hours for adults or 5 tablets (1,625 mg) in 24 hours for children. Severe liver damage may occur if ■ adult takes more than 4,000 mg of acetaminophen in 24 hours ■ child takes more than 5 doses in 24 hours, which is the maximum daily amount ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks every day while using this product. **Allergy alert:** acetaminophen may cause severe skin reactions.

Symptoms may include: ■ skin redness ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions Do not take more than directed. See overdose warning.

adults and children 12 years and over: ■ take 2 tablets every 4 to 6 hours while symptoms last ■ do not take more than 10 tablets in 24 hours, unless directed by a doctor ■ do not use for more than 10 days unless directed by a doctor

children 6 years to under 12 years: ■ take 1 tablet every 4 to 6 hours while symptoms last ■ do not take more than 5 tablets in 24 hours ■ do not use for more than 5 days unless directed by a doctor

children under 6 years ■ ask a doctor

Other information ■ SODIUM FREE ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ use by expiration date on package

Questions or comments? 1-877-290-4008

LOVE IT

OR IT'S ON US!

DISTRIBUTED BY: BETTER LIVING BRANDS LLC
P.O. BOX 99, PLEASANTON, CA 94566-0099
1-888-723-3929

RD 23159 S5954

34UR 0623

Lot No.:

Exp. Date:

Varnish
Omit Area**PAIN RELIEF**

acetaminophen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:21130-043
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z 65AP)	
POVIDONE (UNII: FZ989GH94E)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	TCL340
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-043-01	1 in 1 CARTON	08/08/2023	
1		100 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	08/08/2023	

Labeler - BETTER LIVING BRANDS LLC. (009137209)**Registrant** - TIME CAP LABORATORIES, INC. (037052099)**Establishment**

Name	Address	ID/FEI	Business Operations
TIME CAP LABORATORIES, INC.		037052099	manufacture(21130-043)

Revised: 2/2025

BETTER LIVING BRANDS LLC.