

**LOW DOSE ASPIRIN 81MG- aspirin tablet**  
**BETTER LIVING BRANDS LLC.**

-----  
**481R\_Albertson\_21130-481\_Aspirin 81mg EC Yellow**

***Drug Facts***

***Active ingredient (in each tablet)***

Aspirin 81 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever

***Uses***

- for the temporary relief of minor aches and pains or as recommended by your doctor. **Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.**
- ask your doctor about other uses for enteric coated 81 mg Aspirin.

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if you are** taking a prescription drug for

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  1. feel faint
  2. vomit blood
  3. have bloody or black stools
  4. have stomach pain that does not get better
  - pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - redness or swelling is present
  - new symptoms occur
  - ringing in the ears or a loss of hearing occurs

**If pregnant or breast-feeding, ask a health professional before use . It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

**Other information**

- store at room temperature. Avoid excessive heat above 40°C (104°F)

**Inactive ingredients** anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10 aluminum lake, iron oxide ochre, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate

**Questions or comments? 1-877-290-4008**

Coating/Print  
Omit Area



Lot No.: 2113048199/513  
Exp Date: 3/21/2025  
81023190  
5584  
4818023  
1-888-722-3899  
P.O. BOX 100  
ELIZABETH, NJ 07208  
PAIN RELIEVER  
DISTRIBUTORS  
Bayer HealthCare LLC  
Scan here for  
more product information  
\* Take your doctor's  
advice before using  
the product.

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION**

**TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING**

**Drug Facts**

**Active ingredient (in each tablet)** Aspirin 81 mg (NSAID)\*  
**Purpose** Pain reliever  
\*nonsteroidal anti-inflammatory drug

**Uses** ■ for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief. ■ ask your doctor about other uses for enteric coated 81 mg Aspirin.

**Warnings**  
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
Allergy alert: Aspirin may cause a severe allergic reaction which may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock  
Stomach bleeding warning: This product contains

**Drug Facts (continued)**

an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you are taking a diuretic ■ you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for ■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if ■ an allergic reaction occurs. Seek medical help right away. ■ you experience any of the following signs of stomach bleeding: ● feel faint ● vomit blood ● have bloody or black stools ● have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more

**Drug Facts (continued)**

than 3 days ■ redness or swelling is present ■ new symptoms occur ■ ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions** ■ drink a full glass of water with each dose ■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor ■ children under 12 years: consult a doctor

**Other information** ■ Store at room temperature. Avoid excessive heat above 40°C (104°F)

**Inactive ingredients** anhydrous lactose, carnauba wax, colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10 aluminum lake, iron oxide ochre, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polyorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate

**Questions or comments?** 1-877-290-4008

Compare to Bayer® Low Dose Aspirin active ingredient†

NDC 21130-481-32



Quality Guaranteed

Low Dose  
**Aspirin**  
**81 mg**

Pain reliever (NSAID)

- Enteric coated
- Aspirin regimen\*\*

Actual Size



32 TABLETS

ENTERIC COATING (OATF)

(This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer®.)



100% Recycled  
Paper and Plastic



Coating/Print Omit Area

Lot No.:  
Exp Date:



RD 23159  
52954  
\*\*This type of box is not for sale in the United States.

**READ AND KEEP CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION**

**TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING**

**Drug Facts**

**Active ingredient (in each tablet)** Purpose  
Aspirin 81 mg (NSAID)\* Pain reliever  
hydrocodone bitartrate and ibuprofen

**Uses**  
is for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will provide fast relief of headaches or other symptoms needing immediate relief. In all your doctor visit your doctor about each 81 mg Aspirin.

**Warnings**  
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, it changes in behavior with nausea and vomiting occur, contact a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include: hives, rash, swelling, asthma (wheezing), or shock.  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you are age 65 or older, if you have had stomach ulcers or bleeding problems, if you take a blood thinner (anticoagulant) or stomach drug, if you take other drugs containing prescription or nonprescription NSAIDs (aspirin).

**Drug Facts (continued)**

ibuprofen, naproxen, or others) or have 3 or more alcoholic drinks every day while using this product. Do not use if you are allergic to aspirin or any other pain reliever/fever reducer or if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have a history of stomach problems, such as heartburn, or you have high blood pressure, heart disease, liver disease, or kidney disease. Ask a doctor or pharmacist before use if you are taking a prescription drug for a great or diabetes or arthritis.

Stop use and ask a doctor if an allergic reaction occurs. Seek medical help right away if you experience any of the following signs of stomach bleeding:  
• last faint  
• vomit blood  
• have bloody or black stools  
• have stomach pain that does not get better  
• pain gets worse or lasts more than 3 days  
• lower gas worse or lasts more than 3 days  
• nausea or sweating is present  
• new symptoms occur  
• ringing in the ears or a loss of hearing occurs  
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Drug Facts (continued)**

**Directions**  
Drink a full glass of water with each dose. Adults and children 12 years and over: take 4 to 6 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor. Children under 12 years: consult a doctor.

**Other information** Store at room temperature. Avoid excessive heat above 40°C (104°F).

**Inactive ingredients** anhydrous lactose, croscarmellose sodium, croscarmellose sodium, D10 yellow #10 aluminum lake, iron oxide ochre, methacrylic acid and ethyl acrylate copolymer, microcrystalline cellulose, polyvinyl alcohol, stearic acid, sodium hydroxide, sodium lauryl sulfate, starch, talc, titanium dioxide, triethyl citrate.

**Questions or comments?** 1-877-266-4088

This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer.

**DISTRIBUTED BY:**  
BETTER LIVING BRANDS LLC  
P.O. BOX 90  
PLEASANTON, CA 94566-0090  
1-888-723-3320  
481R 0623



Coating Omit

Compare to Bayer® Low Dose Aspirin active ingredient\*

NDC 211301-481-20

Actual Size



200 TABLETS

Low Dose Aspirin 81 mg

Pain reliever (NSAID)

- Enteric coated
- Aspirin regimen\*\*



Coating/Print Omit Area

Low Dose Aspirin 81 mg  
Pain reliever (NSAID)  
200 TABLETS



NDC 211301-481-20

Coating Omit



<b>SODIUM HYDROXIDE</b> (UNII: 55X04QC32I)
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)
<b>ANHYDROUS LACTOSE</b> (UNII: 3SY5LH9PMK)
<b>METHACRYLIC ACID AND ETHYL ACRYLATE COPOLYMER</b> (UNII: NX76LV5T8J)
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)
<b>TALC</b> (UNII: 7SEV7J4R1U)
<b>TRIETHYL CITRATE</b> (UNII: 8Z96QXD6UM)
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)

### Product Characteristics

<b>Color</b>	yellow	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	7mm
<b>Flavor</b>		<b>Imprint Code</b>	HEART
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-481-32	1 in 1 CARTON	07/14/2023	
1		32 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:21130-481-12	1 in 1 CARTON	07/14/2023	
2		120 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:21130-481-20	1 in 1 CARTON	07/14/2023	
3		200 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:21130-481-50	500 in 1 BOTTLE; Type 0: Not a Combination Product	07/14/2023	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	07/14/2023	

**Labeler** - BETTER LIVING BRANDS LLC. (009137209)

**Registrant** - TIME CAP LABORATORIES, INC. (037052099)

# Establishment

Name	Address	ID/FEI	Business Operations
TIME CAP LABORATORIES, INC.		037052099	manufacture(21130-481)

Revised: 2/2025

BETTER LIVING BRANDS LLC.