

**ALLERGY MULTI-SYMPTOM- acetaminophen, chlorpheniramine maleate and phenylephrine hcl tablet, film coated**

**ARMY AND AIR FORCE EXCHANGE SERVICE**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Exchange Select 44-455C**

***Active ingredients (in each caplet)***

Acetaminophen 325 mg

Chlorpheniramine maleate 2 mg

Phenylephrine HCl 5 mg

***Purpose***

Pain reliever

Antihistamine

Nasal decongestant

***Uses***

- temporarily relieves these symptoms of hay fever or other upper respiratory allergies:
  - headache
  - nasal congestion
  - runny nose and sneezing
  - minor aches and pains
  - sinus congestion and pressure
- temporarily relieves these additional symptoms of hay fever:
  - itchy, watery eyes
  - itching of the nose or throat
- helps clear nasal passages
- helps decongest sinus openings and passages

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters
- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**

- difficulty in urination due to enlargement of the prostate gland
- glaucoma
- high blood pressure
- a breathing problem such as emphysema or chronic bronchitis
- heart disease
- thyroid disease
- diabetes
- liver disease

**Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

**When using this product**

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic beverages
- use caution when driving a motor vehicle or operating machinery
- may cause drowsiness

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain or nasal congestion gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

***Directions***

- **do not take more than directed**
- adults and children 12 years and over
  - take 2 caplets every 4 hours
  - swallow whole - do not crush, chew, or dissolve

- do not take more than 10 caplets in 24 hours
- children under 12 years: ask a doctor

**Other information**

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

**Inactive ingredients**

corn starch, crospovidone, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, sucralose, talc, titanium dioxide

**Questions or comments?**

**1-800-426-9391**

**Principal Display Panel**

exchange √ **select™**

Pseudoephedrine Free

**ALLERGY**

**MULTI-SYMPTOM**

**Acetaminophen,**

Chlorpheniramine maleate, Phenylephrine HCl  
Pain Reliever, Antihistamine, Nasal Decongestant

- Headache
- Runny Nose
- Sinus Pressure
- Sneezing
- Nasal Congestion
- Itchy, Watery Eyes

**Actual Size**

**24 Cool Caplets**

√ **quality value**

50844 REV0818K45508

**"SATISFACTION GUARANTEED OR YOUR MONEY BACK"**


Manufactured For Your Military Exchanges

Distributed by: LNK International, Inc.

Hauppauge, NY 11788

1-800-426-9391

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING**



exchange **select**<sup>™</sup>

Pseudoephedrine Free

# ALLERGY

## MULTI-SYMPTOM

**Acetaminophen,**  
Chlorpheniramine maleate, Phenylephrine HCl  
Pain Reliever, Antihistamine, Nasal Decongestant

- Headache
- Sinus Pressure
- Nasal Congestion
- Runny Nose
- Sneezing
- Itchy, Watery Eyes



Actual Size

24 Cool  
Caplets

quality  
value

50844 REV0818K45508

No print area  
Lot no. & Exp. Date

\*SATISFACTION GUARANTEED OR YOUR MONEY BACK<sup>™</sup>  
Manufactured For Your Military Exchanges  
Distributed by: LNK International, Inc.  
Hempstead, NY 11788  
1-800-426-9391

### Drug Facts

KEEP OUTER PACKAGE FOR COMPLETE  
PRODUCT INFORMATION

#### Active ingredients

(in each caplet)

Acetaminophen 325 mg  
Chlorpheniramine maleate 2 mg  
Phenylephrine HCl 5 mg  
Nasal decongestant

#### Purpose

Pain reliever

### Drug Facts (continued)

#### Uses

temporarily relieves these symptoms of  
hay fever or other upper respiratory allergies:  
■ headache ■ nasal congestion  
■ runny nose and sneezing  
■ minor aches and pains  
■ sinus congestion and pressure

### Drug Facts (continued)

Questions or comments? 1-800-426-9391

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS  
OPENED OR IF BLISTER UNIT IS TORN, BROKEN  
OR SHOWS ANY SIGNS OF TAMPERING



### Drug Facts (continued)

#### Other information

■ TAMPER EVIDENT:  
DO NOT USE IF OUTER PACKAGE IS OPENED OR  
BLISTER IS TORN OR BROKEN  
■ store at 25°C (77°F); excursions permitted  
between 15°-30°C (59°-86°F)  
■ see end flap for expiration date and lot number

### Drug Facts (continued)

#### Inactive ingredients

corn starch, croscollon,  
flavor, magnesium stearate, microcrystalline cellulose,  
polyethylene glycol, polyvinyl alcohol, povidone, silicon  
dioxide, sodium starch glycolate, stearic acid, sucralose,  
talc, titanium dioxide

#### Do not use

■ with any other drug containing acetaminophen  
(prescription or nonprescription). If you are not sure  
whether a drug contains acetaminophen, ask a  
doctor or pharmacist.  
■ If you are now taking a prescription monoamine  
oxidase inhibitor (MAOI) (certain drugs for  
depression, psychiatric or emotional conditions, or  
Parkinson's disease), or for 2 weeks after stopping  
the MAOI drug. If you do not know if your  
prescription drug contains an MAOI, ask a doctor or  
pharmacist before taking this product.  
■ If you have ever had an allergic reaction to this  
product or any of its ingredients

#### Ask a doctor before use if you have

■ difficulty in  
urination due to enlargement of the prostate gland  
■ glaucoma ■ high blood pressure

#### Directions

■ do not take more than directed  
■ adults and children 12 years and over  
take 2 caplets every 4 hours  
■ do not take more than 10 caplets in 24 hours  
■ children under 12 years: ask a doctor

#### Keep out of reach of children. In case of accidental

overdose, get medical help or contact a Poison  
Control Center (1-800-222-1222) right away. Prompt  
medical attention is critical for adults as well as for  
children even if you do not notice any signs or  
symptoms.  
■ If pregnant or breast-feeding, ask a health  
professional before use.  
■ pain or nasal congestion gets worse or lasts more  
than 7 days ■ new symptoms occur  
■ fever gets worse or lasts more than 3 days  
■ redness or swelling is present  
These could be signs of a serious condition.  
If pregnant or breast-feeding, ask a health  
professional before use.

8-0066E-455008AMSR  
REV0818K45508



Exchange Select 44-455

## ALLERGY MULTI-SYMP TOM

acetaminophen, chlorpheniramine maleate and phenylephrine hcl tablet, film coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:5530 1-455
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

### Inactive Ingredients

Ingredient Name	Strength
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TALC (UNII: 7SEV7J4R1U)	
STARCH, CORN (UNII: O8232NY3SJ)	
CROSPVIDONE (UNII: 2S7830E561)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	

### Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	17mm
<b>Flavor</b>	MINT	<b>Imprint Code</b>	44;455
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55301-455-08	2 in 1 CARTON	06/28/2005	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	06/28/2005	

**Labeler** - ARMY AND AIR FORCE EXCHANGE SERVICE (001695568)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	PACK(55301-455)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	MANUFACTURE(55301-455)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	PACK(55301-455)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	PACK(55301-455)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	PACK(55301-455)

Revised: 7/2019

ARMY AND AIR FORCE EXCHANGE SERVICE