

**IPANA 20% BENZOCAINE TOPICAL- benzocaine gel**  
**Maxill Inc.**

-----  
**Ipana 20% Benzocaine Topical Strawberry**

**Active Ingredients (in each gram)**

Benzocaine 200mg

**Purpose**

Oral Anesthetic

**Use**

For oral mucosal use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures.

**Warnings**

**Allergy alert:** Do not use on patients with a history of allergies to local anesthetics such as benzocaine or other "caine" anesthetics.

**Do not use** for more than 7 days unless directed by a dentist or doctor. If sore mouth symptoms do not improve in 7 days; irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your dentist or doctor promptly.

**When using this product** avoid contact with eyes. If contact occurs, flush with water.

**Stop use and consult a health care practitioner** if the following symptoms appear: weakness, confusion, headache, difficulty breathing, and/or pale, grey or blue colored skin, as these may be signs of methemoglobinemia, a rare disorder, which may appear up to 2 hours after use.

**Do not exceed recommended dosage.** If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center immediately.

**If pregnant or breastfeeding,** ask a health professional before use.

**Keep out of reach of children.**

**Directions**

Apply only amount needed to the oral mucosa to prevent or relieve pain.

**Other Information**

Store at 59°-86°F (15°-30°C). Protect from freezing.

**Inactive Ingredients**

flavoring, PEG 3350, PEG 400, sodium saccharin. May contain yellow #5 (tartrazine), yellow #6, red #3, red #40, blue #1, green #3 as a color additive.

**Questions?**

1-855-462-9455 or 1-519-631-7370

## Drug Facts

<b>Active ingredient (in each gram)</b>	<b>Purpose</b>
---	----------------

Benzocaine 200 mg .....	Oral Anesthetic
----------------------------	--------------------

### Use

For oral mucosal use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures.

### Warnings

**Allergy Alert:** Do not use on patients with a history of allergies to local anesthetics such as benzocaine or other "caine" anesthetics.

**Do not use** for more than 7 days unless directed by a dentist or doctor. If sore mouth symptoms do not improve in 7 days; irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your dentist or doctor promptly.

**When using this product** avoid contact with eyes. If contact occurs, flush with water.

**Stop use and consult a health care practitioner** if the following symptoms appear: weakness, confusion, headache, difficulty breathing, and/or pale, grey or blue colored skin, as these may be signs of methemoglobinemia a ▼

## Drug Facts (continued)

rare disorder, which may appear up to 2 hours after use.

### Do not exceed recommended dosage.

If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center immediately.

**If pregnant or breast feeding,** ask a health professional before use.

**Keep out of the reach of children.**

### Directions

Apply only amount needed to the oral mucosa to prevent or relieve pain.

### Other information

Store at 59° - 86° F (15° - 30° C). Protect from freezing.

### Inactive ingredients

flavoring, PEG 3350, PEG 400, sodium saccharine. May contain yellow #5 (tartrazine), yellow #6, red #3, red #40, blue #1, green #3 as a color additive.

### Questions?

1-855-462-9455 or  
1-519-631-7370

Manufactured for:

**maxill inc.**

Cortland, OH, USA 44410  
www.maxill.com

99093750 REV.2016/08

**PEEL BACK**  
For Drug Facts

TO REORDER,  
CALL OR VISIT  
**1-855-462-9455**  
**www.maxill.com**  
MADE IN USA  
REV.2016/08

**ipana**  
**20% BENZOCAINE**  
**TOPICAL GEL**  
Net Wt. 1 oz. (28 g)

**IPANA 20% BENZOCAINE TOPICAL**

benzocaine gel

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:69510-358
<b>Route of Administration</b>	DENTAL		

### Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>BENZOCAINE</b> (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	200 mg in 1 g

### Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>POLYETHYLENE GLYCOL 3350</b> (UNII: G2M7P15E5P)	
<b>POLYETHYLENE GLYCOL 400</b> (UNII: B697894SGQ)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>FD&amp;C YELLOW NO. 5</b> (UNII: I753WB2F1M)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>FD&amp;C RED NO. 3</b> (UNII: PN2ZH5LOQY)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C GREEN NO. 3</b> (UNII: 3P3ONR6O1S)	

### Product Characteristics

<b>Color</b>	red	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	STRAWBERRY	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:69510-358-30	28 g in 1 JAR; Type 0: Not a Combination Product	08/01/2016	

### Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M022	08/01/2016	

**Labeler** - Maxill Inc. (079343581)

