

PEROXYGUARD 102 CONCENTRATE- hydrogen peroxide teat dip solution
AgroChem Inc

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

PEROXYGUARD 102 CONCENTRATE (1:9)

PEROXYGUARD 102 CONCENTRATE

TEAT DIP

WARNING

CAUSES SERIOUS EYE IRRITATION

FOR FARM AND INDUSTRIAL USE ONLY.

KEEP OUT OF REACH OF CHILDREN.

KEEP OUT OF DIRECT SUNLIGHT.

AVOID FREEZING

PREVENTION: Wash skin thoroughly after handling. Wear eye protection/ face protection.

FIRST AID / RESPONSE:

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/ attention.

STORAGE: Store locked up.

DISPOSAL: Dispose of contents/ container to an approved waste disposal plant

EMERGENCY TELEPHONE NUMBER (Chemtrec) 800-424-9300 CCN205494

ACTIVE INGREDIENTS WHEN MIXED:

Hydrogen Peroxide 1.0% (Stabilized)

Lactic Acid - 0.5%

Glycerine - 2.0%

PeroxyGuard is available as a dip, a spray, or in a concentrate.

Use PeroxyGuard effectively to reduce the spread of mastitis causing organisms. Recommended for pre milking applications.

Directions:

General Use

IMPORTANT: THIS IS A CONCENTRATE AND MUST BE DILUTED WITH WATER PRIOR TO USE. Mix (1) part concentrate to (9) parts of clean soft water (mix by volume).

Use a teat dip cup to apply. Clean dip cup just prior to use.

If cow's teats are sore or chapped, consult a veterinarian before starting or continuing to use teat dip.

If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product.

If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container.

Teat Sprays can also be foamed or sprayed using recommended equipment.

Pre Dipping Use

Perform forestripping and inspect teats for clinical mastitis.

Pre-wash teats thoroughly just prior milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels

Just before each cow is milked, dip, foam, or spray, each teat full-length into the teat dip or foam cup containing product.

Allow product to stand for 30 seconds.

Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

Post Dipping Use

Immediately after each milking, use product.

Dip each teat full-length into the teat dip cup containing product. Allow to air dry. Do not wipe.

If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

Extended Use

When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking.

Dipping should be resumed at least twice daily one week before calving.

PEROXYGUARD 102 CONCENTRATE (1:9)

TEAT DIP

WARNING

AVOID FREEZING.
KEEP OUT OF DIRECT SUNLIGHT.
CAUSES SERIOUS EYE IRRITATION.
KEEP OUT OF REACH OF CHILDREN.
FOR FARM AND INDUSTRIAL USE ONLY.



PREVENTION: Wash skin thoroughly after handling. Wear eye protection/ face protection.

FIRST AID / RESPONSE:

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/ attention.

STORAGE: Store locked up.

DISPOSAL: Dispose of contents/ container to an approved waste disposal plant

EMERGENCY TELEPHONE NUMBER (Chemtrec) 800-424-9300 CON05494

ACTIVE INGREDIENTS WHEN MIXED:

Hydrogen Peroxide 1.0% (Stabilized)

Lactic Acid - 0.5%

Glycerine - 2.0%

PeroxyGuard is available as a dip, a spray, or in a concentrate.

Use PeroxyGuard effectively to reduce the spread of mastitis causing organisms. Recommended for pre-milking applications.

Directions:

General Use:

IMPORTANT: THIS IS A CONCENTRATE AND MUST BE DILUTED WITH WATER PRIOR TO USE. Mix (1) part concentrate to (9) parts of clean soft water (mix by volume).

Use a test dip cup to apply. Clean dip cup just prior to use.

If cow's teats are sore or chapped, consult a veterinarian before starting or continuing to use test dip.

If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product.

If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container.

Test Sprays can also be foamed or sprayed using recommended equipment.

Pre Dipping Use:

Perform forestripping and inspect teats for clinical mastitis.

Pre-wash teats thoroughly just prior milking with appropriate udder wash solution or pre-milking test dip. Teats should then be dried with single-service towels.

Just before each cow is milked, dip, foam, or spray, each teat full-length into the test dip or foam cup containing product.

Allow product to stand for 30 seconds.

Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

Post Dipping Use:

Immediately after each milking, use product.

Dip each teat full-length into the test dip cup containing product. Allow to air dry. Do not wipe.

If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

Extended Use:

When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking.

Dipping should be resumed at least twice daily one week before calving.



AgroChem

26 Freedom Way - Saratoga Springs, NY 12866 - (518)226-4850

LOT#:

EXP:

PEROXYGUARD 102 CONCENTRATE

hydrogen peroxide teat dip solution

Product Information

| | | | |
|-------------------------|-----------------|--------------------|----------------|
| Product Type | OTC ANIMAL DRUG | Item Code (Source) | NDC:17307-8820 |
| Route of Administration | topical | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------|------------------|
| HYDROGEN PEROXIDE (UNII: BBX060AN9V) (HYDROGEN PEROXIDE - UNII:BBX060AN9V) | HYDROGEN PEROXIDE | 0.18 kg in 1 kg |
| LACTIC ACID (UNII: 33X04XA5AT) (LACTIC ACID - UNII:33X04XA5AT) | LACTIC ACID | 0.045 kg in 1 kg |

Inactive Ingredients

| Ingredient Name | Strength |
|-----------------------------|-----------------|
| GLYCERIN (UNII: PDC6A3C0OX) | 0.18 kg in 1 kg |
| WATER (UNII: 059QF0K00R) | 0.38 kg in 1 kg |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|-------------------------|----------------------|--------------------|
| 1 | NDC:17307-8820-1 | 19 kg in 1 CONTAINER | | |
| 2 | NDC:17307-8820-2 | 57 kg in 1 DRUM | | |
| 3 | NDC:17307-8820-3 | 208 kg in 1 DRUM | | |
| 4 | NDC:17307-8820-4 | 1040 kg in 1 CELLO PACK | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|-----------------------|--|----------------------|--------------------|
| unapproved drug other | | 03/16/2020 | |

Labeler - AgroChem Inc (805427700)

Registrant - AgroChem Inc (805427700)

Establishment

| Name | Address | ID/FEI | Business Operations |
|---------------|---------|-----------|------------------------------|
| Agro Chem Inc | | 805427700 | api manufacture, manufacture |

Revised: 3/2020

AgroChem Inc