BOVA PRO 55- iodine spray AgroChem Inc

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

BovaPro 55 Teat Spray

BOVAPRO 55 TEAT SPRAY

IODINE

TEAT SPRAY

WARNING

CAUSES SERIOUS EYE IRRITATION

FOR FARM AND INDUSTRIAL USE ONLY.

KEEP OUT OF REACH OF CHILDREN.

KEEP OUT OF DIRECT SUNLIGHT.

AVOID FREEZING

PREVENTION: Wash skin thoroughly after handling. Wear eye protection/ face protection.

FIRST AID / RESPONSE:

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention.

STORAGE: Store locked up.

DISPOSAL: Dispose of contents/container to an approved waste disposal plant

EMERGENCY TELEPHONE NUMBER (Chemtrec) 800-424-9300 CCN205494

INGREDIENTS:

Active Ingredient - Iodine 0.5 % (5.0 % Polyethoxy Polypropoxy Polyethoxy Ethanol - Iodine Complex) Titration via USP33 Reissue, Topical Solution

Emollients: Glycerine 5.0%

BovaPro is made with IoPlex technology and is NPE free. BovaPro is available as a dip, a spray, or in a concentrate.

Use BovaPro effectively to reduce the spread of mastitis causing organisms. Recommended for pre and post milking applications.

Directions:

General Use

Ready to use. Do not dilute. Do not mix product with water, any other teat dips, or other product.

Use a teat dip cup to apply. Clean dip cup just prior to use.

If cow's teats are sore or chapped, consult a veterinarian before starting or continuing to use teat dip.

If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product.

If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container.

Teat Sprays can also be foamed or sprayed using recommended equipment.

Pre Dipping Use

Perform forestripping and inspect teats for clinical mastitis.

Pre-wash teats thoroughly just prior milking with appropriate udder wash solution or premilking teat dip. Teats should then be dried with single-service towels

Just before each cow is milked, dip, foam, or spray, each teat full-length into the teat dip or foam cup containing product.

Allow product to stand for 30 seconds.

Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

Post Dipping Use

Immediately after each milking, use product.

Dip each teat full-length into the teat dip cup containing product. Allow to air dry. Do not wipe.

If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

Extended Use

When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking.

Dipping should be resumed at least twice daily one week before calving.

BOVAPRO 55 TEAT SPRAY

IODINE TEAT SPRAY

WARNING



CAUSES SERIOUS EYE IRRITATION

FOR FARM AND INDUSTRIAL USE ONLY.
KEEP OUT OF REACH OF CHILDREN.
KEEP OUT OF DIRECT SUNLIGHT.
AVOID FREEZING

PREVENTION: Wash skin thoroughly after handling. Wear eye protection/ face protection.

FIRST AID / RESPONSE:

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/ attention.

STORAGE: Store locked up.

DISPOSAL: Dispose of contents/ container to an approved waste disposal plant

EMERGENCY TELEPHONE NUMBER (Chemtrec) 800-424-9300 CCN205494

INGREDIENTS:

Active Ingredient - Iodine 0.5 % (5.0 % Polyethoxy Polypropoxy Polyethoxy Ethanol - Iodine Complex) Titration via USP33 Reissue, Topical Solution

Emollients: Glycerine 5.0%

BovaPro is made with IoPlex technology and is NPE free. BovaPro is available as a dip, a spray, or in a concentrate.

NET CONTENTS:



Use BovaPro effectively to reduce the spread of mastitis causing organisms. Recommended for pre and post milking applications.

Directions:

General Use

Ready to use. Do not dilute. Do not mix product with water, any other teat dips, or other product.

Use a teat dip cup to apply. Clean dip cup just prior to use.

If cow's teats are sore or chapped, consult a veterinarian before starting or continuing to use teat dip.

If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product.

If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container.

Teat Sprays can also be foamed or sprayed using recommended equipment.

Pre Dipping Use

Perform forestripping and inspect teats for clinical mastitis.

Pre-wash teats thoroughly just prior milking with appropriate udder wash solution or pre-milking teat dip.

Teats should then be dried with single-service towels

Just before each cow is milked, dip, foam, or spray, each teat full-length into the teat dip or foam cup containing product.

Allow product to stand for 30 seconds.

Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

Post Dipping Use

Immediately after each milking, use product.

Dip each teat full-length into the teat dip cup containing product. Allow to air dry. Do not wipe. If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

Extended Use

When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking.

Dipping should be resumed at least twice daily one week before calving.





BOVAPRO 55 TEAT SPRAY



WARNING



CAUSES SERIOUS EYE IRRITATION

FOR FARM AND INDUSTRIAL USE ONLY.
KEEP OUT OF REACH OF CHILDREN.
KEEP OUT OF DIRECT SUNLIGHT.
AVOID FREEZING

PREVENTION: Wash skin thoroughly after handling. Wear eye protection/ face protection.

FIRST AID / RESPONSE:

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention.

STORAGE: Store locked up.

DISPOSAL: Dispose of contents/ container to an approved waste disposal plant

EMERGENCY TELEPHONE NUMBER (Chemtrec) 800-424-9300 CCN205494

INGREDIENTS:

Active Ingredient - Iodine 0.5 % (5.0 % Polyethoxy Polypropoxy Polyethoxy Ethanol - Iodine Complex) Titration via USP33 Reissue, Topical Solution

Emollients: Glycerine 5.0%

BovaPro is made with IoPlex technology and is NPE free. BovaPro is available as a dip, a spray, or in a concentrate.

NET CONTENTS:



Use BovaPro effectively to reduce the spread of mastitis causing organisms. Recommended for pre and post milking applications.

Directions:

General Use

 $Ready \ to \ use. \ Do \ not \ dilute. \ Do \ not \ mix \ product \ with \ water, any \ other \ teat \ dips, or \ other \ product.$

Use a teat dip cup to apply. Clean dip cup just prior to use.

If cow's teats are sore or chapped, consult a veterinarian before starting or continuing to use teat dip.

If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product.

If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container.

Teat Sprays can also be foamed or sprayed using recommended equipment.

Pre Dipping Use

Perform forestripping and inspect teats for clinical mastitis.

Pre-wash teats thoroughly just prior milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels

Just before each cow is milked, dip, foam, or spray, each teat full-length into the teat dip or foam cup containing product.

Allow product to stand for 30 seconds.

Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

Post Dipping Use

Immediately after each milking, use product.

Dip each teat full-length into the teat dip cup containing product. Allow to air dry. Do not wipe. If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

Extended Use

When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking.

Dipping should be resumed at least twice daily one week before calving.





BOVA PRO 55

iodine spray

{

Product Information			
Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:17307-0550
Pouts of Administration	TOPICAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
IODINE (UNII: 9679TC07X4) (IODINE - UNII:9679TC07X4)	IODINE	.05 kg in 1 kg	

Inactive Ingredients			
Ingredient Name	Strength		
WATER (UNII: 059QF0KO0R)	.90 kg in 1 kg		
GLYCERIN (UNII: PDC6A3C0OX)	.05 kg in 1 kg		

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:17307-0550-1	19 kg in 1 CONTAINER		
2	NDC:17307-0550-2	57 kg in 1 DRUM		
3	NDC:17307-0550-3	208 kg in 1 DRUM		
4	NDC:17307-0550-4	1040 kg in 1 CONTAINER		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		08/01/2013	

Labeler - AgroChem Inc (805427700)

Establishment				
Name	Address	ID/FEI	Business Operations	
Agro Chem Inc		805427700	api manufacture, manufacture	

Revised: 1/2018 AgroChem Inc