# H-E-B SOLUTIONS SUNSCREEN COCONUT SPF 50 CONTINUOUS- avobenzone, homosalate, octisalate, octocrylene spray H.E.B

-----

#### H-E-B SOLUTIONS SUNSCREEN COCONUT SPF 50 CONTINUOUS SPRAY

#### **Active Ingredients**

Avobenzone 3.0%, Homosalate 10.0%, Octisalate 5.0%, Octocrylene 4.0%

#### **Purpose**

Sunscreen

#### Uses

- helps prevent sunburn
- if used as directed with other sun protetion measures (see **Directions**), decreases the risk of skin cancer and early skin aging caused by the sun

#### Warnings

#### For external use only

#### FLAMMABLE:

- do not use near heat, flame or while smoking
- avoid long term stoage above 104°F (40°C)

#### Do not use

• on damaged or broken skin.

#### When using this product

• keep out of eyes. Rinse with water to remove. • do not puncture or incinerate. Contents under pressure. • do not store at temperatures above 120°F.

### Stop use and ask a doctor if

• rash occurs.

### Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

#### **Directions**

- apply liberally 15 minutes before sun exposure
- hold can 4-6 inches away from body, spray evenly to ensure complete coverage
- do not spray into face. Spray into hand and apply to face.
- use in well ventilated, but not windy areas
- reapply:
- after 80 minutes of swimming or sweating
- immediately after towel drying
- at least every 2 hours
- children under 6 months of age: Ask a doctor
- Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:
- limit time in the sun, especially from 10 a.m. 2 p.m.
- wearl long-sleeved shirts, pants, hats, and sunglasses

#### Other Information


- protect the product in this container from excessive heat and direct sun
- may stain or damage some fabrics, materials or surfaces

## □Inactive Ingredients

SD alcohol 40-B, butyloctyl salicylate, acrylates/octylacrylamide copolymer, glycerin, saccharomyces/podophyllum peltatum ferment filtrate, caprylic/capric triglyceride, stearoxytrimethylsilane, tocopherol, panthenol, butylene glycol, fragrance

#### Label





#### PABA, Retinyl Palmitate & Paraben Free\* No Oxybenzone or Octinoxate • Dermatologist Tested

## Drug Facts

Active ingredients

Avobenzone 3%, Homosalate 10% Octisalate 5%, Octocrylene 4% .....

\*\*Uses\*\* • helps prevent sunburn\*\* • if used as directed with other sun protection measures (see \*\*Directions\*\*), decreases the risk of skin cancer and early skin aging caused by the sun

*Warnings* For external use only

Flammable:

• do not use near heat, flame or while smoking
• avoid long term storage above 104°F (40°C)

Do not use . on damaged or broken skin.

When using this product • keep out of eyes. Rinse with water to remove. • Keep away from face to avoid breathing it. • do not puncture or incinerate. Contents under pressure. • do not store at temperatures above 120° F.

Stop use and ask a doctor if . rash occurs.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions • Spray Iberally and spread evenly by hand 15 minutes before sun exposure • Hold can 4-6 inches away from the skin to apply • Do not spray directly into face. Spray into hands then apply to face • Do not apply in windy conditions • Use in a well-ventilated area • reapply: • after 80 minutes of swimming or • sweating • immediately after towel drying • at least every 2 hours • children under 6 months of age: Ask a doctor • Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including: • limit time in the sun, especially from 10 a.m. - 2 o.m. • wear long-sleeved shirts, anats, hats, and sunclasses a.m. - 2 p.m. • wear long-sleeved shirts, pants, hats, and sunglasses

Other information • protect the product in this container from excessive heat and direct sun • may stain or damage some fabrics, materials or surfaces

Inactive ingredients acrylates/octylacrylamide copolymer, butylene dycol, butyloctyl salcylate, caprylic/capric triglycerde, fragrance, glycerin, parthenol, saccharomyces/podochyllum petatum ferment filtrate, SD alcohol 40-B, stearoxytrimethylsitane, tocopherol.

Questions or comments? Call toll free 1-800-527-7731

IO PARABENS SEPARATELY ADDED TO PRESERVE THIS PRODUC







**Dermatologist Tested** Hypoallergenic · Oil Free · PABA Free Retinyl Palmitate Free · Paraben Free Contains Vitamin E

#### **Drug Facts**

Active Ingredients
Avokenzone 3%
Nomosalsie 10%
Octosinte 5%
Octocrylene 4% Purvose Sunscreen

Uses - helps prevent sunburn - if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

## Warnings For external use only

- FLAMMABLE:
- do not use near heat, flame or while smoking
   avoid long term storage above 104°F (40°C)
- Do not use + on damaged or broken skin.

When using this product \* keep out of eyes, Rinse with water to remove, \* do not puncture or incinerate, Contents under pressure, \* do not store at temperatures above 120° F.

Stop use and ask a doctor if + rash occurs,

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

#### Directions

- Directions

   apply liberally 15 minutes before sun exposure

   apply liberally 15 minutes before sun exposure

   boll can 4-6 inches away from body, spray avenly to ensure complete coverage

   do not spray into face, Spray into hand and apply to face,

   use in well ventilated, but not windy areas

   reapply:

   after 30 minutes of swimming or sweating

   immediately after towel drying

   at least every 2 hours

- at least every 2 hours
- at least every 2 hours
   children under 6 morths of age; ask a doctor
   Sun Protection Measures, Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad-apactum SPF of 15 or higher and other sun protection measures including:
- Iimit time in the sun, especially from 10 a.m. 2 p.m.
   wear long-sleeve shirts, pants, hats, and sunglasses

Other Information

• protect this product from excessive heat and direct sun

• may stain or damage some fabrics, materials or surfaces

Inactive Ingredients
SD alcohol 40-8, butyloctyl salicylate, acrylates/octylacrylamide
copolymer, fragrance, glycerin, caprylic/capric triglyceride,
stearoxytrimethylaliane, butylene glycol, tocopherol,
saccharomyces/podophyllum peltatum ferment filtrate, panthenol

## Questions or Comments? Call toll free 1-800-527-7731

MADE WITH PRIDE AND CARE FOR H-E-B°,

SAN ANTONIO, TEXAS 78204

























Sunscreen **UVA/UVB Protection** Water Resistant

(80 minutes) Coconut Scented Paraben Free

**BROAD SPECTRUM SPF 50** 





NET WT. 5.5 OZ (156 g)





## H-E-B SOLUTIONS SUNSCREEN COCONUT SPF 50 CONTINUOUS

avobenzone, homosalate, octisalate, octocrylene spray

<b>Product Information</b>			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:37808-905
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety				
Ingredient Name	<b>Basis of Strength</b>	Strength		
AVOBENZONE (UNII: G63QQF2NOX) (AVOBENZONE - UNII:G63QQF2NOX)	AVOBENZONE	30 mg in 1 g		
HOMOSALATE (UNII: V06SV4M95S) (HOMOSALATE - UNII:V06SV4M95S)	HOMOSALATE	100 mg in 1 g		
OCTISALATE (UNII: 4X49Y0596W) (OCTISALATE - UNII:4X49Y0596W)	OCTISALATE	50 mg in 1 g		
OCTOCRYLENE (UNII: 5A68WGF6WM) (OCTOCRYLENE - UNII:5A68WGF6WM)	OCTOCRYLENE	40 mg in 1 g		

Inactive Ingredients			
Ingredient Name	Strength		
ALCOHOL (UNII: 3K9958V90M)			
ACRYLATE/ISOBUTYL METHACRYLATE/N-TERT-OCTYLACRYLAMIDE COPOLYMER (75000 MW) (UNII: JU3XHR8VWK)			
TOCOPHEROL (UNII: R0ZB2556P8)			
GLYCERIN (UNII: PDC6A3C0OX)			
STEAROXYTRIMETHYLSILANE (UNII: 9862TW94B2)			
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)			
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)			
BUTYLOCTYL SALICYLATE (UNII: 2EH13UN8D3)			
PANTHENOL (UNII: W/9CM0067Z)			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37808-905- 37	156 g in 1 CAN; Type 0: Not a Combination Product	03/07/2019	
2	NDC:37808-905- 48	258 g in 1 CAN; Type 0: Not a Combination Product	01/25/2022	
3	NDC:37808-905- 74	76.5 g in 1 CAN; Type 0: Not a Combination Product	01/04/2024	
4	NDC:37808-905- 63	2 in 1 PACKAGE	01/04/2024	
4		156 g in 1 CAN; Type 0: Not a Combination Product		

Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
OTC Monograph Drug	M020	03/07/2019			

## **Labeler -** H.E.B (007924756)

Revised: 9/2024 H.E.B