

MAGNESIUM CITRATE- magnesium citrate liquid
Meijer Distribution Inc

Meijer Magnesium Citrate Oral Solution - Cherry

Active Ingredient

Magnesium citrate 1.745 g (in each fl oz)

Purpose

Saline laxative

Uses

- For relief of occasional constipation (irregularity)
- Generally produces bowel movement in 1/2 to 6 hours

Warnings

Ask a doctor before use if you have

- Kidney disease
- a magnesium restricted diet
- Abdominal pain, nausea, or vomiting
- Noticed a sudden change in bowel habits that persists over a period of 2 weeks
- Already used a laxative for a period longer than 1 week

Ask a doctor or pharmacist before use if you are

- Taking any other drug.
- Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if

you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Store at temperatures between 46° and 86°F (8° and 30° C)

Directions

- shake well before using
- drink a full glass (8 ounces) of liquid with each dose
- may be taken as a single daily dose or in divided doses

| | |
|--|--|
| adults and children 12 years of age and over | 6.5 to 10 fl oz maximum 10 fl oz in 24 hours |
| children 6 to under 12 years of age | 3 to 7 fl oz maximum 7 fl oz in 24 hours |
| children 2 to under 6 years of age | 2 to 3 fl oz in 24 hours maximum 3 fl oz in 24 hours |
| children under 2 years of age | ask a doctor |

Other Information

each fl oz contains: magnesium 290 mg

each fl oz contains: sodium 1 mg

Inactive Ingredients

benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water

Questions?

1-888-235-2466

DIST. MEIJER DISTRIBUTION, INC.

GRAND RAPIDS, MI 48455

www.meijer.com L0000700

Do not use if tamper evident twist-off cap is missing, broken, or separated from neckring

NDC 79481-0033-9

meijer®

CHERRY FLAVOR

magnesium

citrate

Saline Laxative/ Oral Solution

• DYE-FREE • SUGAR-FREE

Alcohol content 0.005% (from flavouring)

10 FL OZ (296 mL)

Drug Facts**Active ingredient (in each fl oz) Purpose**
Magnesium citrate 1.745 gSaline laxative**Uses:** ■ For relief of occasional constipation (irregularity).
■ Generally produces bowel movement in ½ to 6 hours**Warnings:****Ask a doctor before use if you have**

- Kidney disease ■ a magnesium restricted diet
- Abdominal pain, nausea, or vomiting
- Noticed a sudden change in bowel habits that persists over a period of 2 weeks
- Already used a laxative for a period longer than 1 week

Ask a doctor or pharmacist before use if you are

- Taking any other drug.
- Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.**If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.****Do not use if tamper evident twist-off cap is missing, broken, or separated from neckring**COPY FREE AREA LEFT OPEN FOR
LOT AND EXPIRATION DATE

NDC 79481-0033-9

meijer**CHERRY FLAVOR**
magnesium citrate**Saline Laxative / Oral Solution**

• DYE-FREE • SUGAR-FREE

Alcohol content 0.005% (from flavoring)

10 FL OZ (296 mL)**Drug Facts (continued)**

Store at temperatures between 46° and 86°F (8° and 30°C)

Directions: ■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses

| | |
|--|---|
| adults and children 12 years of age and over | 6.5 to 10 fl oz maximum 10 fl oz in 24 hours |
| children 6 to under 12 years of age | 3 to 7 fl oz maximum 7 fl oz in 24 hours |
| children 2 to under 6 years of age | 2 to 3 fl oz maximum 3 fl oz in 24 hours |
| children under 2 years of age | ask a doctor |

Other Information:

- each fl oz contains: magnesium 290 mg
- each fl oz contains: sodium 1 mg

Inactive ingredients: benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water**Questions?** 1-888-235-2466DIST. MEIJER DISTRIBUTION, INC.
GRAND RAPIDS, MI 49544 L0000700
www.meijer.com PID 5155746 R0223**MAGNESIUM CITRATE**

magnesium citrate liquid

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|----------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:79481-0033 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------|-----------------------|
| MAGNESIUM CITRATE (UNII: RHO26O1T9V) (MAGNESIUM CATION - UNII:T6V3LHY838) | MAGNESIUM CITRATE | 1.745 g in 29.6 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| BENZOIC ACID (UNII: 8SKN0B0MIM) | |
| WATER (UNII: 059QF0KO0R) | |
| CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP) | |
| SUCRALOSE (UNII: 96K6UQ3ZD4) | |
| EDETATE DISODIUM ANHYDROUS (UNII: 8NLQ36F6MM) | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|-----------|---------------------|----------------------|--------------------|
| | | | | |

| | | | | |
|------------------------------|---|--|-----------------------------|---------------------------|
| 1 | NDC:79481-0033-9 | 296 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | 04/24/2023 | |
| Marketing Information | | | | |
| Marketing Category | Application Number or Monograph Citation | | Marketing Start Date | Marketing End Date |
| OTC Monograph Drug | M007 | | 04/24/2023 | |

Labeler - Meijer Distribution Inc (006959555)

Revised: 12/2024

Meijer Distribution Inc