MAGNESIUM CITRATE- magnesium citrate liquid Meijer Distribution Inc

Meijer - Magnesium Citrate Oral Solution - Grape

Active Ingredient

Magnesium citrate 1.745 g (in each fl oz)

Purpose

Saline laxative

Uses

- For relief of occasional constipation (irregularity)
- Generally produces bowel movement in 1/2 to 6 hours

Warnings

Ask a doctor before use if you have

- Kidney disease
- a magnesium restricted diet
- Abdominal pain, nausea, or vomiting
- Noticed a sudden change in bowel habits that persists over a period of 2 weeks
- Already used a laxative for a period longer than 1 week

Ask a doctor or pharmacisy before use if you are

- Taking any other drug.
- Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if

you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Store at temperatures between 46° and 86°F (8° and 30° C)

Directions

- shake well before using
- drink a full glass (8 ounces) of liquid with each dose
- may be taken as a single daily dose or in divided doses

adults and children 12 years of age and over	6.5 to 10 fl oz maximum 10 fl oz in 24 hours	
children 6 to under 12 years of age	3 to 7 fl oz maximum 7 fl oz in 24 hours	
children 2 to under 6 years of age	2 to 3 fl oz in 24 hours maximum 3 fl oz in 24 hours	
children under 2 years of age	ask a doctor	

Other Information

each fl oz contains: magnesium 290 mg

each fl oz contains: sodium 1 mg

Inactive Ingredients

benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water

Questions?

1-888-235-2466

DIST. MEIJER DISTRIBUTION, INC.

GRAND RAPIDS, MI 49544

www.meijer.com L0000750

R0823

Do not use if tamper-evident twist-off cap is

missing, broken or separated from neckring

NDC 79481-0034-9

meijer®

GRAPE FLAVOR

magnesium

citrate

Saline Laxative/ Oral Solution

• DYE-FREE • SUGAR-FREE

Alcohol content 0.0125% (from flavouring)

10 FL OZ (296 mL)

Drug Facts

Active ingredient (in each fl oz) Purpose Magnesium citrate 1.745 gSaline laxative

Uses: For relief of occasional constipation (irregularity).

Generally produces bowel movement in ½ to 6 hours

- Warnings:
 Ask a doctor before use if you have
 Kidney disease a magnesium restricted diet
 Abdominal pain, nausea, or vomiting
 Noticed a sudden change in bowel habits that persists over a period of 2 weeks
- Already used a laxative for a period longer than 1 week

Ask a doctor or pharmacist before use if you are

- Taking any other drug.
 Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Do not use if tamper evident twist-off cap is missing, broken, or separated from neckring

COPY FREE AREA LEFT OPEN FOR LOT AND EXPIRATION DATE

meijer.

Magnesium Citrate

Saline Laxative/Oral Solution

Alcohol content 0.0125% (from flavoring)

Dye-Free

Sugar-Free



NDC 79481-0034-9

(296 mL)

Drug Facts (continued)

Store at temperatures between 46° and 86°F (8° and 30°C)

Directions: ■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses

adults and children 12 6.5 to 10 fl oz maximum years of age and over 10 fl oz in 24 hours children 6 to under 12 3 to 7 fl oz maximum years of age 7 fl oz in 24 hours children 2 to under 6 2 to 3 fl oz maximum years of age 3 fl oz in 24 hours children under 2 years of age

ask a doctor

Other Information:

- each fl oz contains: magnesium 290 mg each fl oz contains: sodium 1 mg

Inactive ingredients: benzoic acid. citric acid, disodium EDTA, flavor, sucralose, water

Questions? 1-888-235-2466

DIST. BY MEIJER DISTRIBUTION, INC. GRAND RAPIDS, MI 49544 www.meijer.com

L0000750



MAGNESIUM CITRATE

magnesium citrate liquid

Product Information

Product Type HUMAN OTC DRUG **Item Code (Source)** NDC:79481-0034

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
	MAGNESIUM CITRATE	1.745 g in 29.6 mL

Inactive Ingredients			
Ingredient Name	Strength		
BENZOIC ACID (UNII: 85KN0B0MIM)			
WATER (UNII: 059QF0KO0R)			
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)			
SUCRALOSE (UNII: 96K6UQ3ZD4)			
EDETATE DISODIUM ANHYDROUS (UNII: 8NLQ36F6MM)			

Packaging

#	Item Code	Package Description	Marketing Start Date	Date
	NDC:79481- 0034-9	296 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	04/24/2023	
M	larketing	Information		
M	larketing Marketing Category	Information Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
	Marketing	Application Number or Monograph Citation	_	

Labeler - Meijer Distribution Inc (006959555)

Establishment							
Name	Address	ID/FEI	Business Operations				
Pharma Nobis, LLC		118564114	manufacture(79481-0034)				

Revised: 12/2025 Meijer Distribution Inc