

DEFENSOL D- acetaminophen, chlorpheniramine maleate, phenylephrine hci tablet

Menper Distributors, Inc.

Defensol D

Active ingredients (in each tablet)

Acetaminophen 325 mg

Chlorpheniramine maleate 2 mg

Phenylephrine HCl 5 mg

Purpose

Pain Reliever/ Fever Reducer

Antihistamine

Nasal Decongestant

Uses

Temporarily relieves these symptoms associated with cold or flu:

- headache
- nasal congestion
- sore throat
- fever
- minor aches and pains

Temporarily relieves minor aches, pains and headache as well as these symptoms of hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- nasal congestion
- itching of the nose or throat
- itchy, watery eyes

Temporarily relieves minor aches, pains, headache and nasal congestion as well as sinus congestion and pressure, and reduces swelling of nasal passages

Warnings

Liver Warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 tablets in any 24-hour period, which is the maximum daily amount
- with other drugs containing acetaminophen

- 3 or more alcoholic drinks every day while using this product

Sore throat warning:if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- To sedate a child or to make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for two weeks after stopping the MAOI, drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- glaucoma
- a breathing problem such as emphysema, asthma, or chronic bronchitis

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking any other oral nasal decongestant or stimulant
- taking any other pain reliever/ fever reducer
- taking sedatives or tranquilizers

When using this product

- **do not use more than directed**
- excitability may occur with children
- drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days

- redness or swelling is present
- new symptoms occur

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions: do not take more than 8 tablets in a 24 hour period.

- do not exceed recommended dosage. Taking more than the recommended dose (overdose) may cause serious liver damage

age	dose
adults and children 12 years and over	2 tablets every 4 hours
children under 12 years	do not use

Inactive ingredients: FD&C blue #2, hypromellose, magnesium stearate, PEG 400, povidone, pregelatinized starch, silicon dioxide, stearic acid, titanium dioxide

Questions? Menper Distributors, Inc. **1-800-560-5223**; M-F 9 AM - 4 PM Eastern

DEFENSOL D

Acetaminophen • Pain Reliever/Fever reducer
Chlorpheniramine maleate • Antihistamine
Phenylephrine HCl • Nasal decongestant

Multi-Symptom
Formula

Relieves:

- Nasal Congestion
- Sinus Headache/Pain
- Sneezing
- Itchy, Watery Eyes
- Itchy Throat
- Runny Nose

30 Tablets

Drug Facts

Active ingredients (in each tablet)
 Acetaminophen 325 mg
 Chlorpheniramine maleate 2 mg
 Phenylephrine HCl 5 mg

Purpose
 Pain Reliever/Fever Reducer
 Antihistamine
 Nasal Decongestant

Uses
 Temporarily relieves these symptoms associated with a cold or flu:
 ■ headache ■ nasal congestion ■ sore throat ■ fever
 ■ minor aches and pains
 ■ sinus pain, pressure, and headache, as well those
 symptoms of hay fever or other upper respiratory allergies:
 ■ runny nose ■ sneezing ■ nasal congestion
 ■ itching of the nose or throat ■ itchy, watery eyes
 ■ itchy throat ■ watery eyes
 ■ as well as sinus congestion and pressure, and reduces swelling of nasal
 passages

Warnings
 Overdose warning: This product contains acetaminophen.
 ■ more than 8 tablets in any 24-hour period, which is the maximum daily amount
 ■ with other drugs containing acetaminophen
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (Parkinson's disease) or for 2 weeks after stopping the MAOI drug.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
Do not use
 ■ to sedate a child or to make a child sleepy
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (Parkinson's disease) or for 2 weeks after stopping the MAOI drug.
 If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
 ■ if you are pregnant, planning to get pregnant, or breastfeeding, or if you are nursing a child.
 ■ nonprescription. If you are not sure whether a drug contains acetaminophen ask a doctor or pharmacist.
Ask a doctor before use if you have
 ■ liver disease ■ heart disease ■ high blood pressure
 ■ thyroid disease ■ diabetes ■ enlarged prostate gland ■ glaucoma
 ■ a breathing problem such as emphysema, asthma, or chronic bronchitis
Ask a doctor or pharmacist before use if you are
 ■ taking the blood thinning drug warfarin
 ■ taking any other oral nasal decongestant or stimulant
 ■ taking any other pain reliever/fever reducer
 ■ taking sedatives or tranquilizers
When using this product
 ■ do not use more than directed ■ drowsiness may occur
 ■ alcohol, sedatives and tranquilizers may increase drowsiness
 ■ be careful when driving a motor vehicle or operating machinery
 ■ excitability may occur especially in children

Distributed by: **Menpar Distributors, Inc.**
 6500 NW 35th Ave., Miami, FL 33147
 www.menpardistributors.com
 Lot# / Exp:
 Rev: 11/11
 PEEL HERE FOR DRUG FACTS INFORMATION

Drug Facts (continued)

Stop use and ask a doctor if ■ you get nervous, dizzy, or sleepless
 ■ pain or nasal congestion gets worse or lasts more than 7 days
 ■ symptoms do not improve after 7 days of use
 ■ redness or swelling is present ■ new symptoms occur

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

Overdose warning
 ■ If you or your child get medical help or contact a Poison Control Center right away. From medical attention, you may need to be hospitalized or hospitalized if you do not notice any signs or symptoms.

Directions ■ do not take more than 8 tablets in a 24 hour period
 ■ take 2 tablets every 4 hours, as directed. Do not use this product if printed seal around neck is torn, broken, or missing.
 ■ dose (overdose) may cause serious liver damage.
 ■ this adult product is not intended for use in children under 12 years of age

age	dose
adults and children 12 years and over	2 tablets every 4 hours
children under 12 years	do not use

Other information
 ■ store at room temperature, 15°-30°C (59°-86°F)
 ■ TAMPER-EVIDENT DISCLOSURE: Do not use this product if printed seal around neck is torn, broken, or missing.

Inactive ingredients FD&C blue #2, hydroxyethylcellulose, magnesium stearate, PEG 400, polydioxane, sodium carboxymethyl starch, starch, silicon dioxide, stearic acid, titanium dioxide

Questions?
 Menpar Distributors, Inc. 1-800-560-9223
 M-F 9 AM - 4 PM Eastern Standard Time

Defensol D is not manufactured by Schering-Plough, S.A. de C.V. (SP-MEXICO), owner of the registered trademark "Defensol D".

DEFENSOL D

acetaminophen, chlorpheniramine maleate, phenylephrine hci tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53145-005
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
HYPROMELLOSE 2208 (100 MPA.S) (UNII: B1QE5P712K)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POVIDONE (UNII: FZ989GH94E)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	blue	Score	2 pieces
Shape	ROUND	Size	10mm
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53145-005-02	50 in 1 CARTON	02/21/2012	
1		2 in 1 PACKAGE; Type 0: Not a Combination Product		
2	NDC:53145-005-30	1 in 1 BOX	02/21/2012	
2		30 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	02/21/2012	

Labeler - Menper Distributors, Inc. (101947166)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc		868734088	manufacture(53145-005)

Revised: 7/2025

Menper Distributors, Inc.