# ONLY FOR FEVER- acetaminophen tablet O4 Global Trading Usa, Llc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

# Only Forth OFEVER

# **Drug Facts**

## Active ingredient (in each caplet)

Acetaminophen 500mg

# **Purpose**

Pain reliever / Fever reducer

#### Uses:

- temporarily relieves minor aches and pains due to:
  - the common cold
  - backache
  - toothache
  - premenstrual and menstrual cramps
  - temporarily reduces fever.
  - headache
  - minor pain of arthritis
  - muscular aches
- temporarily reduces fever.

## Warnings:

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

# **Allergy Alert:**

Acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

#### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had any allergic reaction to this product or any of its ingredients.

**Ask a doctor before use if you have** liver disease

**Ask a doctor or pharmacist before use if are** taking the blood thinning drug warfarin

## Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or last more than 3 days
- new symptoms occur.
- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children. OVERDOSE WARNING:** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions:**

Do not take more than directed. **SEE OVERDOSE WARNING.** 

adults and children 12 years of age and over	<ul> <li>take 2 caplets every 6 hours while symptoms last</li> <li>do not take more than 6 caplets in 24 hours, unless directed by doctor</li> <li>do not take more than 10 days, unless directed by a doctor</li> </ul>
children under 12 years of age	• ask a doctor

## Other Information:

- Store at room temperature between 20-25°C (68-77°F).
- avoid excessive heat, cold and humidity.
- close cap tightly after use.

## **Inactive Ingredients:**

hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide

# **Questions or comments?**

**1-833-665-9444.** (Mon-Fri 9am-5pm)

# **ADULT**

# PAIN RELIEVER FEVER REDUCER

**TAMPER EVIDENT:** Do not use if imprinted safety seal under cap is broken or missing

Distributed By: O4 GLOBAL TRADING USA, LLC.

12222 Merit Drive, Suite 1900

Dallas, TX 75251-3209

# www.onlyfor.com

Packaged By: Strive Pharmaceuticals Inc.

East Brunswick, NJ 08816.

# **Packaging**





(in each caplet, active ingredient

**Drug Facts** 

temporarily relieves minor aches and pains due to:

 premenstrual and menstrual the common coldbackache toothache

minor pain of arthritismuscular aches

iver warning: This product contains acetaminophen. temporarily reduces fever.

 more than 4,000 mg of acetaminophen in 24 hours
 with other drugs containing acetaminophen
 3 or more alcoholic drinks every day while usin Severe liver damage may occur if you take:

this

using

Lot No. 8

# HERE

## Drug Facts

Active ingredient (in each caplet)

Purpose

Acetaminophen 500mg......Pain reliever / Fever reducer

#### Uses:

- temporarily relieves minor aches and pains due to:
- the common cold
- headache
- backache
- minor pain of arthritis
- toothache
- muscular aches
- premenstrual and menstrual cramps
- temporarily reduces fever.

#### Warnings:

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4.000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product 1

# Drug Facts (continued)

#### Allergy Alert:

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

#### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have liver disease Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

### Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present.

These could be signs of serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. OVERDOSE WARNING: In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

## **Drug Facts** (continued)

## Directions:

Do not take more than directed. SEE OVERDOSE WARNING.

adults and children 12 years of age and over

- take 2 caplets every 6 hours while symptoms last
- do not take more than 6 caplets in 24 hours, unless directed by doctor
- do not take more than 10 days. unless directed by a doctor

children under 12 vears

ask a doctor

#### Other Information:

- Store at room temperature between 20-25°C (68-77°F).
- avoid excessive heat, cold and humidity.
- close cap tightly after use.

#### Inactive Ingredients:

hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide

#### Questions or comments?

1-833-665-9444, (Mon-Fri 9am - 5pm)

Distributed By: 04 GLOBAL TRADING USA, LLC.

12222 Merit Drive, Suite 1900 Dallas, TX 75251-3209

www.onlyfor.com

Packaged By: Strive Pharmaceuticals Inc. East Brunswick, NJ 08816.

3

LB7300614395-REV.00-072020

# **ONLY FOR FEVER**

acetaminophen tablet

D 1 .	T C
Product	Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:73006-143

Route of Administration ORAL

# Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINO PHEN (UNII: 36209 ITL9 D) (ACETAMINO PHEN - UNII: 36209 ITL9 D)	ACETAMINOPHEN	500 mg

# **Inactive Ingredients**

Ingredient Name	Strength
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICRO CRYSTALLINE CELLULO SE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PO VIDO NE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIO XIDE (UNII: 15FIX9 V2JP)	

# **Product Characteristics**

Color	white	Score	no score
Shape	CAPSULE	Size	17mm
Flavor		Imprint Code	S500
Contains			

# **Packaging**

	# Item Code	Package Description	Marketing Start Date	Marketing End Date
	1 NDC:73006-143- 95	1 in 1 CARTON	11/15/2020	
l	1	25 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

# **Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	11/15/2020	

# Labeler - O4 Global Trading Usa, Llc (081226861)

Revised: 11/2020 O4 Global Trading Usa, Llc