

SHISEIDO SPORTS BB DARK- octinoxate and titanium dioxide emulsion
SHISEIDO AMERICAS CORPORATION

SHISEIDO SPORTS BB DARK

Drug Facts

<i>Active ingredients</i>	<i>Purpose</i>
OCTINOXATE 4.9%	Sunscreen
TITANIUM DIOXIDE 15%	Sunscreen

Uses

- helps prevent sunburn
- if used as directed with other sun protection measures (see ***Directions***), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs.

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Flammable : Do not use near heat, flame, or while smoking.

Directions

- apply liberally 15 minutes before sun exposure
- reapply
 - after 80 minutes after swimming and sweating
 - immediately after towel drying
 - at least every 2 hours
- **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF of 15 or higher and other sun protection measures including:
 - limit time in the sun, especially from 10 a.m. - 2 p.m.
 - wear long-sleeve shirts, pants, hats, and sunglasses
 - children under 6 months: Ask a doctor

Inactive Ingredients

DIMETHICONE ■ SILICA ■ BIS-PEG-18 METHYL ETHER DIMETHYL

SILANE■ISODODECANE■TRIMETHYLSILOXYSILICATE■SD ALCOHOL 40-B■TRIMETHYLSILOXYSILICATE/DIMETHICONOL CROSSPOLYMER■ISOPROPYL MYRISTATE■ALUMINUM HYDROXIDE■PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE■STEARIC ACID■ISOSTEARIC ACID■DEXTRIN PALMITATE■DISTEARDIMONIUM HECTORITE■BIS-BUTYLDIMETHICONE POLYGLYCERYL-3■LAUROYL LYSINE■PEG/PPG-14/7 DIMETHYL ETHER■PCA DIMETHICONE■PHYTOSTERYL/OCTYLDODECYL LAUROYL GLUTAMATE■PHYTOSTERYL MACADAMIATE■LECITHIN■GLYCYRRHIZA GLABRA (LICORICE) ROOT EXTRACT■SUCROSE TETRASTEARATE TRIACETATE■POLYMETHYLSILSESQUIOXANE■TALC■METHYL METHACRYLATE CROSSPOLYMER■ALUMINA■POLYSILICONE-2■TRIETHOXYCAPRYLYLSILANE■HYDROGEN DIMETHICONE■TOCOPHEROL■BHT■ETHYLPARABEN■FRAGRANCE■TITANIUM DIOXIDE■IRON OXIDES■MICA■

Other information

- Protect this product in this container from excessive heat and direct sun.
- Avoid contact with clothing and other textiles
 - may stain.

Questions or comments?

Call toll free 1-800-906-7503

PRINCIPAL DISPLAY PANEL - 30 mL Bottle Carton

SHISEIDO
GINZA TOKYO

Sports BB

50+
WetForce
Quick Dry

BROAD SPECTRUM
SPF 50+

WATER RESISTANT
(80 MINUTES)

SUNSCREEN

30mL 1 FL. OZ.



Sports BB



BROAD SPECTRUM SPF 50+

SHISEIDO

GINZA TOKYO

Sports BB

50+

WetForce

Quick Dry

BROAD SPECTRUM
SPF 50+

WATER RESISTANT
(80 MINUTES)

SUNSCREEN

30mL 1 FL. OZ.


Drug Facts (continued)

Inactive ingredients

DIMETHICONE • SILICA •
BIS-PEG-18 METHYL ETHER DIMETHYL SILANE •
ISODODECANE • TRIMETHYLSILOXYSILICATE •
SD ALCOHOL 40-8 •
TRIMETHYLSILOXYSILICATE/DIMETHICONOL
CROSSPOLYMER •
ISOPROPYL MYRISTATE • ALUMINUM HYDROXIDE •
PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE •
STEARIC ACID • ISOSTEARIC ACID •
DEXTRIN PALMITATE •
DISTEAROINONIUM Hectorite •
BIS-BUTYLDIMETHICONE POLYGLYCERYL-3 •
LAUROYL LYSINE • PEG/PPG-14/7 DIMETHYL ETHER •
PCA DIMETHICONE •
PHYTOSTERYL OCTYLDODECYL LAUROYL GLUTAMATE •
PHYTOSTERYL MACADAMATE • LECITHIN •
GLYCYRRHIZA GLABRA (LICORICE) ROOT EXTRACT •
SUCROSE TETRASTEARATE TRIACETATE •
POLYMETHYLSILOXODIMANE • TALC •
METHYL METHACRYLATE CROSSPOLYMER • ALUMINA •
POLYSILOXANE-2 • TRIETHOXYCAPRYLYLSILANE •
HYDROGEN DIMETHICONE • TOCOPHEROL-BHT •
ETHYLPARABEN • FRAGRANCE • TITANIUM DIOXIDE •
IRON OXIDES • MICA •

Other Information

- protect this product in this container from excessive heat and direct sun.
- avoid contact with clothing and other textiles-may stain.



SHISEIDO
SPORTS BB
BROAD SPECTRUM SPF 50+

A BB type sunscreen ideal for outdoor scenes such as sports, featuring WetForce technology so the UV protection veil actually becomes stronger when it comes into contact with sweat and water.* Furthermore, Quick Dry technology dries your sweat quickly** to keep you looking fresh, no matter how active you have been.

- Contains contouring powder to utilize the effects of light and shadow for creating a sculpted, three-dimensional look.
- Reducing the quantity of application will lower the level of sunscreen protection significantly.
- Shake well before use.

NON-COMEDOGENIC. DERMATOLOGIST AND OPHTHALMOLOGIST-TESTED.

* To activate WetForce technology, expose the applied area to water or perspiration for 30 minutes.

**in vitro test

Drug Facts

Active ingredients	Purpose
OCTINOXATE 4.9%	Sunscreen
TITANIUM DIOXIDE 15.0%	Sunscreen

Uses

■ helps prevent sunburn ■ if used as directed with other sun protection measures (see *Directions*), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

For external use only

Do not use on damaged or broken skin

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away. ▶

Drug Facts (continued)

Flammable : Do not use near heat, flame, or while smoking.

Directions

For sunscreen use:

■ apply liberally 15 minutes before sun exposure ■ reapply: - after 80 minutes of swimming or sweating - immediately after towel drying - at least every 2 hours ■ Sun Protection Measures. Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF of 15 or higher and other sun protection measures including: - limit time in the sun, especially from 10 a.m. - 2 p.m. - wear long-sleeve shirts, pants, hats, and sunglasses ■ children under 6 months: Ask a doctor ▶




<http://s1872.com/14657AA>
See our website

54.5 X 33 X 82

75 1 2 3 4 5 6 7 8 9 0



SHISEIDO AMERICAS CORPORATION DIST.
NEW YORK, NY 10022
MADE IN U.S.A.



SHISEIDO SPORTS BB DARK

octinoxate and titanium dioxide emulsion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58411-536
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	1.81545 g in 30 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	5.5575 g in 30 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
BIS-PEG-18 METHYL ETHER DIMETHYL SILANE (UNII: OEB4R3VM9C)	
ISODODECANE (UNII: A8289P68Y2)	
TRIMETHYLSILOXYSILICATE (M/Q 0.6-0.8) (UNII: 5041RX63GN)	
ALCOHOL (UNII: 3K9958V90M)	
ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0)	

PEG-9 POLYDIMETHYLSILOXYETHYL DIMETHICONE (UNII: TYP81E471F)
STEARIC ACID (UNII: 4ELV7Z65AP)
ISOSTEARIC ACID (UNII: X33R8U0062)
DEXTRIN PALMITATE (CORN; 20000 MW) (UNII: 89B2BSF9I3)
DISTEARDIMONIUM HECTORITE (UNII: X687XDK09L)
LAUROYL LYSINE (UNII: 113171Q70B)
PEG/PPG-14/7 DIMETHYL ETHER (UNII: 6DNW9T7YT2)
PHYTOSTERYL/OCTYLDODECYL LAUROYL GLUTAMATE (UNII: 65954KGO9Q)
PHYTOSTERYL MACADAMIATE (UNII: 233VSF903M)
GLYCYRRHIZA GLABRA (UNII: 2788Z9758H)
SUCROSE TETRASTEARATE TRIACETATE (UNII: 1K7LBQ045N)
POLYMETHYLSILSESQUIOXANE (4.5 MICRONS) (UNII: 59Z907ZB69)
TALC (UNII: 7SEV7J4R1U)
METHYL METHACRYLATE/GLYCOL DIMETHACRYLATE CROSSPOLYMER (UNII: EG97988M5Q)
ALUMINUM OXIDE (UNII: LMI26O6933)
TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)
HYDROGEN DIMETHICONE (13 CST) (UNII: 4QGR4P2YOI)
.ALPHA.-TOCOPHEROL (UNII: H4N855PNZ1)
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)
ETHYLPARABEN (UNII: 14255EXE39)
FERRIC OXIDE RED (UNII: 1K09F3G675)
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)
FERROSFERRIC OXIDE (UNII: XM0M87F357)
MICA (UNII: V8A1AW0880)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58411-536-10	1 in 1 CARTON	03/01/2019	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M020	03/01/2019	

Labeler - SHISEIDO AMERICAS CORPORATION (193691821)