

CORID- amprolium solution
Huvepharma, Inc

CORID®
(amprolium oral solution)
9.6% Oral Solution Coccidiostat

Net Contents: 1 gal. (128 fl. oz) (3,785 mL)

Active Ingredient: amprolium 9.6%

Approved by FDA under NADA # 013-149

Distributed by: **Huvepharma®**
Huvepharma, Inc.
Peachtree City, GA 30269

INDICATIONS FOR USE: An aid in the treatment and prevention of coccidiosis caused by *Eimeria bovis* and *E. zurnii* in calves. For a satisfactory diagnosis a microscopic examination of the feces should be done before treatment. When treating outbreaks, drug should be administered promptly after diagnosis is determined.

DOSAGE AND ADMINISTRATION:

	5-Day Treatment Protocol	21-Day Prevention
Protocol	Daily Dosage: 10 mg amprolium/kg (10 mg per 2.2 lb body weight)	Daily Dosage: 5 mg amprolium/kg (5 mg per 2.2 lb body weight)

USE DIRECTIONS: There are **96** mg of amprolium in every **1 mL** of CORID 9.6% Solution.

1 fl oz = 29.57 mL	8 fl oz = 1/2 pint (236.56 mL)
1 fl oz = 2 measuring tablespoons	16 fl oz = 1 pint (473.12 mL)

To disperse particles in the product, shake the bottle prior to each use.

IN DRINKING WATER

Mix CORID daily with fresh drinking water.

5-DAY TREATMENT: Add CORID 9.6% Oral Solution to drinking water at the rate of 16 fl oz/100 gal. At the usual rate of water consumption this will provide an intake of approximately 10 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 5 days. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

21-DAY PREVENTION: During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add CORID 9.6% Oral Solution to drinking water at

the

rate of 8 fl oz/100 gal. At usual rates of water consumption this will provide an intake of approximately 5 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 21 days.

AS A DRENCH

CORID drench solutions may be stored in a clean, closed, labeled container for up to 3 days.

5-DAY TREATMENT: Add 3 fl oz CORID 9.6% Oral Solution to 1 pt of water and, with a dose

syringe, give 1 fl oz of this drench solution for each 100 lb (45 kg) body weight. This will provide

a dose of approximately 10 mg amprolium/kg (2.2 lb) body weight. Give daily for 5 days. Use

on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest

of the group has been exposed, and all calves in the group should be treated.

21-DAY PREVENTION: During periods of exposure or when experience indicates that coccidiosis

is likely to be a hazard, add 1 1/2 fl oz of CORID 9.6% Oral Solution to 1 pt of water and, with a

dose syringe, give 1 fl oz of this drench solution for each 100 lb (45 kg) body weight. This will

provide a dose of approximately 5 mg amprolium/kg (2.2 lb) body weight. Give daily for 21

days. Use as the sole source of amprolium.

WITHDRAWAL PERIODS AND RESIDUE WARNINGS: Withdraw 24 hours before slaughter. A withdrawal period has

not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

USER SAFETY WARNINGS: NOT FOR HUMAN USE. Keep this and all drugs out of the reach of children.

Contact Huvepharma Inc. at 1-877-994-4883 or <http://www.huvepharma.us>. For additional information

about reporting side effects for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

PRECAUTIONS: FOR ORAL USE IN ANIMALS ONLY. MAY CAUSE EYE IRRITATION. For irritation, flush with

plenty of water; get medical attention.

STORAGE: Store between 5°-25°C (41°- 77°F) with brief excursions to 40°C.

Note: When product

is sitting for prolonged periods of time, black, brown, or orange to yellow particles may appear on the surface.

Benzoic acid 0.1% added as preservative.

®CORID is a registered trademark of Huvepharma EOOD

Lot Number and Expiration Date can be found on shoulder of bottle.

Restricted Drug (California) - Use Only as Directed

KCP-S00190

Version: 07

Rev: 06-2024

CORID[®]

(amprolium oral solution)

9.6% Oral Solution Coccidiostat

Net Contents: 16 oz (473 ml)

Active Ingredient: amprolium.....9.6%

Approved by FDA under NADA # 013-149

Distributed by:
Huvepharma, Inc.
Peachtree City, GA 30269



INDICATIONS FOR USE : An aid in the treatment and prevention of coccidiosis caused by *Eimeria bovis* and *E. zurnii* in calves. For a satisfactory diagnosis a microscopic examination of the feces should be done before treatment. When treating outbreaks, drug should be administered promptly after diagnosis is determined.

DOSAGE AND ADMINISTRATION:

5-Day Treatment Protocol
Daily Dosage: 10 mg amprolium/kg
(10 mg per 2.2 lb body weight)

21-Day Prevention Protocol
Daily Dosage: 5 mg amprolium/kg
(5 mg per 2.2 lb body weight)

USE DIRECTIONS : There are 96 mg of amprolium in every 1 mL of CORID 9.6% Solution.
1 fl oz = 29.57 mL
1 fl oz = 2 measuring tablespoons

8 fl oz = 1/2 pint (236.56 mL)
16 fl oz = 1 pint (473.12 mL)

IN DRINKING WATER

Mix CORID daily with fresh drinking water.

5-DAY TREATMENT: Add CORID 9.6% Oral Solution to drinking water at the rate of 16 fl oz/100 gal. At the usual rate of water consumption this will provide an intake of approximately 10 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 5 days. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

21-DAY PREVENTION: During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add CORID 9.6% Oral Solution to drinking water at the rate of 8 fl oz/100 gal. At usual rates of water consumption this will provide an intake of approximately 5 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 21 days.

© CORID is a registered trademark of Huvepharma (USA).
Lot Number and Expiration Date are located on shoulder of bottle.

AS A DRENCH

CORID drench solutions may be stored in a clean, closed, labeled container for up to 3 days.

5-DAY TREATMENT: Add 3 fl oz CORID 9.6% Oral Solution to 1 pt of water and, with a dose syringe, give 1 fl oz of this drench solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 10 mg amprolium/kg (2.2 lb) body weight. Give daily for 5 days. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

21-DAY PREVENTION: During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add 1 1/2 fl oz of CORID 9.6% Oral Solution to 1 pt of water and, with a dose syringe, give 1 fl oz of this drench solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 5 mg amprolium/kg (2.2 lb) body weight. Give daily for 21 days.

Use as the sole source of amprolium.

WITHDRAWAL PERIODS AND RESIDUE WARNINGS: Withdraw 24 hours before slaughter. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

USER SAFETY WARNINGS: NOT FOR HUMAN USE. Keep this and all drugs out of the reach of children. Contact Huvepharma Inc. at 1-877-994-4883 or <http://www.huvepharma.us>. For additional information about reporting side effects for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

PRECAUTIONS: FOR ORAL USE IN ANIMALS ONLY. MAY CAUSE EYE IRRITATION. For irritation, flush with plenty of water; get medical attention.

STORAGE: Store between 5°-25°C (41°-77°F) with brief excursions to 40°C.

Benzoic acid 0.1% added as preservative.

Technical Assistance - 1-877-994-4883



KCP-S00012
Version:06
Rev:01-2025

CORID[®]

(amprolium oral solution)

9.6% Oral Solution Coccidiostat

Net Contents: 1 gal. (128 fl. oz) (3,785 mL)

Active Ingredient: amprolium9.6%

Approved by FDA under NADA # 013-149

Distributed by:
Huvepharma, Inc.
Peachtree City, GA 30269



INDICATIONS FOR USE: An aid in the treatment and prevention of coccidiosis caused by *Eimeria bovis* and *E. zurnii* in calves. For a satisfactory diagnosis a microscopic examination of the feces should be done before treatment. When treating outbreaks, drug should be administered promptly after diagnosis is determined.

DOSAGE AND ADMINISTRATION:

5-Day Treatment Protocol
Daily Dosage: 10 mg amprolium/kg
(10 mg per 2.2 lb body weight)

21-Day Prevention Protocol
Daily Dosage: 5 mg amprolium/kg
(5 mg per 2.2 lb body weight)

USE DIRECTIONS: There are 96 mg of amprolium in every 1 mL of CORID 9.6% Solution.

1 fl oz = 29.57 mL 8 fl oz = 1/2 pint (236.56 mL)
1 fl oz = 2 measuring tablespoons 16 fl oz = 1 pint (473.12 mL)

IN DRINKING WATER

Mix CORID daily with fresh drinking water.

5-DAY TREATMENT: Add CORID 9.6% Oral Solution to drinking water at the rate of 16 fl oz/100 gal. At the usual rate of water consumption this will provide an intake of approximately 10 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 5 days. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

21-DAY PREVENTION: During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add CORID 9.6% Oral Solution to drinking water at the rate of 8 fl oz/100 gal. At usual rates of water consumption this will provide an intake of approximately 5 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the the only source of water for 21 days.

AS A DRENCH

CORID drench solutions may be stored in a clean, closed, labeled container for up to 3 days.

5-DAY TREATMENT: Add 3 fl oz CORID 9.6% Oral Solution to 1 pt of water and, with a dose syringe, give 1 fl oz of this drench solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 10 mg amprolium/kg (2.2 lb) body weight. Give daily for 5 days. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

21-DAY PREVENTION: During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add 1 1/2 fl oz of CORID 9.6% Oral Solution to 1 pt of water and, with a dose syringe, give 1 fl oz of this drench solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 5 mg amprolium/kg (2.2 lb) body weight. Give daily for 21 days. Use as the sole source of amprolium.

WITHDRAWAL PERIODS AND RESIDUE WARNINGS: Withdraw 24 hours before slaughter. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

USER SAFETY WARNINGS: NOT FOR HUMAN USE. Keep this and all drugs out of the reach of children. Contact Huvepharma Inc. at 1-877-994-4883 or <http://www.huvepharma.us>. For additional information about reporting side effects for animal drugs, contact FDA at 1-888-FDA-VEIS or <http://www.fda.gov/reportanimalae>.

PRECAUTIONS: FOR ORAL USE IN ANIMALS ONLY. MAY CAUSE EYE IRRITATION. For irritations, flush with plenty of water; get medical attention.

STORAGE: Store between 5°-25°C (41°-77°F) with brief excursions to 40°C.

Benzoic acid 0.1% added as preservative.

© CORID is a registered trademark of Huvepharma EOOD.

Lot Number and Expiration Date can be found on shoulder of bottle.

Restricted Drug (California) - Use Only as Directed



EXP 500010
Version: 05
Rev: 01-2025

CORID[®]

(amprolium oral solution)

9.6% Oral Solution Coccidiostat

Net Contents: 16 oz (473 ml)

Active Ingredient: amprolium.....9.6%

Approved by FDA under NADA # 013-149

Distributed by:
Huvepharma, Inc.
Peachtree City, GA 30269



INDICATIONS FOR USE : An aid in the treatment and prevention of coccidiosis caused by *Eimeria bovis* and *E. zurnii* in calves. For a satisfactory diagnosis a microscopic examination of the feces should be done before treatment. When treating outbreaks, drug should be administered promptly after diagnosis is determined.

DOSAGE AND ADMINISTRATION:

5-Day Treatment Protocol
Daily Dosage: 10 mg amprolium/kg
(10 mg per 2.2 lb body weight)

21-Day Prevention Protocol
Daily Dosage: 5 mg amprolium/kg
(5 mg per 2.2 lb body weight)

USE DIRECTIONS : There are 96 mg of amprolium in every 1 mL of CORID 9.6% Solution.
1 fl oz = 29.57 mL
1 fl oz = 2 measuring tablespoons
8 fl oz = 1/2 pint (236.56 mL)
16 fl oz = 1 pint (473.12 mL)

To disperse particles in the product, shake the bottle prior to each use.

IN DRINKING WATER

Mix CORID daily with fresh drinking water.

5-DAY TREATMENT: Add CORID 9.6% Oral Solution to drinking water at the rate of 16 fl oz/100 gal. At the usual rate of water consumption this will provide an intake of approximately 10 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 5 days. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

21-DAY PREVENTION: During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add CORID 9.6% Oral Solution to drinking water at the rate of 8 fl oz/100 gal. At usual rates of water consumption this will provide an intake of approximately 5 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 21 days.

© CORID is a registered trademark of Huvepharma TCOO.
Lot Number and Expiration Date are located on shoulder of bottle.

AS A DRENCH

CORID drench solutions may be stored in a clean, closed, labeled container for up to 3 days.

5-DAY TREATMENT: Add 3 fl oz CORID 9.6% Oral Solution to 1 pt of water and, with a dose syringe, give 1 fl oz of this drench solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 10 mg amprolium/kg (2.2 lb) body weight. Give daily for 5 days. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

21-DAY PREVENTION: During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add 1 1/2 fl oz of CORID 9.6% Oral Solution to 1 pt of water and, with a dose syringe, give 1 fl oz of this drench solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 5 mg amprolium/kg (2.2 lb) body weight. Give daily for 21 days.

Use as the sole source of amprolium.

WITHDRAWAL PERIODS AND RESIDUE WARNINGS: Withdraw 24 hours before slaughter. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

USER SAFETY WARNINGS: NOT FOR HUMAN USE. Keep this and all drugs out of the reach of children. Contact Huvepharma Inc. at 1-877-994-4883 or <http://www.huvepharma.us>. For additional information about reporting side effects for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

PRECAUTIONS: FOR ORAL USE IN ANIMALS ONLY. MAY CAUSE EYE IRRITATION. For irritation, flush with plenty of water; get medical attention.

STORAGE: Store between 5°-25°C (41°-77°F) with brief excursions to 40°C. Note: When product is sitting for prolonged periods of time, black, brown, or orange to yellow particles may appear on the surface.

Benzoic acid 0.1% added as preservative.

Technical Assistance - 1-877-994-4883



KCP-S00192
Version: 08
Rev: 06-2024

CORID[®]

(amprolium oral solution)

9.6% Oral Solution Coccidiostat

Net Contents: 1 gal. (128 fl. oz) (3,785 mL)

Active Ingredient: amprolium9.6%

Approved by FDA under NADA # 013-149

Distributed by:
Huvepharma, Inc.
Peachtree City, GA 30269



INDICATIONS FOR USE: An aid in the treatment and prevention of coccidiosis caused by *Eimeria bovis* and *E. zurnii* in calves. For a satisfactory diagnosis a microscopic examination of the feces should be done before treatment. When treating outbreaks, drug should be administered promptly after diagnosis is determined.

DOSAGE AND ADMINISTRATION:

5-Day Treatment Protocol
Daily Dosage: 10 mg amprolium/kg
(10 mg per 2.2 lb body weight)

21-Day Prevention Protocol
Daily Dosage: 5 mg amprolium/kg
(5 mg per 2.2 lb body weight)

USE DIRECTIONS : There are 96 mg of amprolium in every 1 mL of CORID 9.6% Solution.

1 fl oz = 29.57 mL
1 fl oz = 2 measuring tablespoons
8 fl oz = 1/2 pint (236.56 mL)
16 fl oz = 1 pint (473.12 mL)

To disperse particles in the product, shake the bottle prior to each use.

IN DRINKING WATER

Mix CORID daily with fresh drinking water.

5-DAY TREATMENT: Add CORID 9.6% Oral Solution to drinking water at the rate of 16 fl oz/100 gal. At the usual rate of water consumption this will provide an intake of approximately 10 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 5 days. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

21-DAY PREVENTION: During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add CORID 9.6% Oral Solution to drinking water at the rate of 8 fl oz/100 gal. At usual rates of water consumption this will provide an intake of approximately 5 mg amprolium/kg (2.2 lb) body weight. Offer this solution as the only source of water for 21 days.

AS A DRENCH

CORID drench solutions may be stored in a clean, closed, labeled container for up to 3 days.

5-DAY TREATMENT: Add 3 fl oz CORID 9.6% Oral Solution to 1 pt of water and, with a dose syringe, give 1 fl oz of this drench solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 10 mg amprolium/kg (2.2 lb) body weight. Give daily for 5 days. Use on a herd basis only; when one or more calves show signs of coccidiosis, it is likely that the rest of the group has been exposed, and all calves in the group should be treated.

21-DAY PREVENTION: During periods of exposure or when experience indicates that coccidiosis is likely to be a hazard, add 1 1/2 fl oz of CORID 9.6% Oral Solution to 1 pt of water and, with a dose syringe, give 1 fl oz of this drench solution for each 100 lb (45 kg) body weight. This will provide a dose of approximately 5 mg amprolium/kg (2.2 lb) body weight. Give daily for 21 days.

Use as the sole source of amprolium.

WITHDRAWAL PERIODS AND RESIDUE WARNINGS: Withdraw 24 hours before slaughter. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

USER SAFETY WARNINGS: NOT FOR HUMAN USE. Keep this and all drugs out of the reach of children. Contact Huvepharma Inc. at 1-877-994-4883 or <http://www.huvepharma.us>. For additional information about reporting side effects for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>.

PRECAUTIONS: FOR ORAL USE IN ANIMALS ONLY. MAY CAUSE EYE IRRITATION. For irritations, flush with plenty of water; get medical attention.

STORAGE: Store between 5°-25°C (41°-77°F) with brief excursions to 40°C. Note: When product is sitting for prolonged periods of time, black, brown, or orange to yellow particles may appear on the surface.

Benzoic acid 0.1% added as preservative.

© CORID is a registered trademark of Huvepharma EOOD.

Lot Number and Expiration Date can be found on shoulder of bottle.

Restricted Drug (California) - Use Only as Directed



KCP-S00192
Version: 07
Rev: 16-2024

CORID

amprolium solution

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:23243-2140
Route of Administration	ORAL		

Active Ingredient/Active Moiety				
Ingredient Name			Basis of Strength	Strength
amprolium (UNII: 95CO6N199Q) (amprolium ion - UNII:H2T307KMZR)			amprolium	96 mg in 1 mL
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:23243-2140-2	473 mL in 1 BOTTLE, PLASTIC		
2	NDC:23243-2140-1	3785 mL in 1 BOTTLE, PLASTIC		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
NADA	NADA013149		12/12/2017	

Labeler - Huvepharma, Inc (619153559)

Registrant - Huvepharma EOOD (552671651)

Establishment			
Name	Address	ID/FEI	Business Operations
Huvepharma, Inc		883128204	medicated animal feed manufacture, pack, label, manufacture, analysis