

# **SPF 30 SUNSCREEN- zinc oxide, titanium dioxide lotion CLEURE**

-----  
**Cleure SPF 30 Sunscreen**

## **ACTIVE INGREDIENT**

Zinc oxide 9%

Titanium Dioxide 4.51%

## **PURPOSE**

Sunscreen

## **USES**

• helps prevent sunburn • If used as directed with other sun protection measures decreases the risk of skin cancer and early skin aging caused by the sun

## **WARNINGS**

• For external use only • Do not use on damaged or broken skin • When using this product keep out of eyes. Rinse with water to remove. • Stop use and ask doctor if rash occurs.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

## **DIRECTIONS**

Apply liberally 15 minutes before sun exposure • reapply: • after 80 minutes of swimming or sweating • immediately after towel drying • at least every 2 hours • Children under 6 months of age: ask a doctor • Sun Protection Measures: Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF of 15 or higher and other sun protection measures including: • Limit time in the sun, especially from 10 a.m. - 2pm • wear long-sleeved shirts, pants, hats and sunglasses.

## **INACTIVE INGREDIENTS**

Water, Caprylic / Capric Triglyceride, Cetearyl Alcohol, STEARETH-2, Glycerin, STEARETH-21, Aloe Vera, Dimethicone, Phenoxyethanol, Sodium Stearoyl Glutamate, Xanthan Gum, Disodium EDTA, Ethylhexylglycerin

protect the product in this container from excessive heat and direct sun.

### Drug Facts

Active ingredient	Purpose
Zinc oxide 9%.....	Sunscreen
Titanium Dioxide 4.51%.....	Sunscreen

**Uses** • helps prevent sunburn • If used as directed with other sun protection measures decreases the risk of skin cancer and early skin aging caused by the sun

**Warnings** • For external use only • Do not use on damaged or broken skin • When using this product keep out of eyes. Rinse with water to remove. • Stop use and ask doctor if rash occurs. • Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

**Directions** • Apply liberally 15 minutes before sun exposure • reapply: • after 80 minutes of swimming or sweating • immediately after towel drying • at least every 2 hours • Children under 6 months of age: ask a doctor • **Sun Protection Measures:** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF of 15 or higher and other sun protection measures including: • Limit time in the sun, especially from 10 a.m. - 2pm • wear long-sleeved shirts, pants, hats and sunglasses.

**Other information** • protect the product in this container from excessive heat and direct sun.

**Inactive Ingredients** • D.I. Water, Caprylic/Capric Triglyceride, Cetearyl Alcohol, Polyhydroxystearic Acid, Glycerin, Stearic Acid, Trihydroxycaprylylsilane, Dimethicone, Phenoxyethanol, Ethylhexylglycerin, Sodium Stearoyl Glutamate

**Questions?** 1-805-981-7600; 1-888-883-4276 or visit [www.cleure.com](http://www.cleure.com)

Distr. By Grace Products, Inc.  
Ventura, California 93003  
[www.cleure.com](http://www.cleure.com)  
1-888-883-4276



7 18415124012

# cleure®

Ingredients with Intention

**DAILY FACE & BODY  
BROAD SPECTRUM**

**SUNSCREEN  
SPF 30**

**MINERAL BASED | LIGHTWEIGHT**

*FREE OF: alcohol, dyes,  
gluten, parabens, fragrance  
formaldehyde, salicylates*

**WITH TITANIUM DIOXIDE  
& ZINC OXIDE**

**DERMATOLOGIST  
RECOMMENDED**

*CRUELTY FREE*

**FOR ALL SKIN TYPES,  
INCLUDING SENSITIVE SKIN**

**4 FL OZ (118 ml)**

## SPF 30 SUNSCREEN

zinc oxide, titanium dioxide lotion

### Product Information

**Product Type**

HUMAN OTC DRUG

**Item Code (Source)**

NDC:83294-101

<b>Route of Administration</b>	TOPICAL
--------------------------------	---------

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>ZINC OXIDE</b> (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)	ZINC OXIDE	9 g in 100 mL
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	4.51 g in 100 mL

**Inactive Ingredients**

Ingredient Name	Strength
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>DIMETHICONE 350</b> (UNII: 2Y53S6ATLU)	
<b>SODIUM STEAROYL GLUTAMATE</b> (UNII: 65A9F4P024)	
<b>STEARETH-21</b> (UNII: 53J3F32P58)	
<b>ALOE VERA LEAF</b> (UNII: ZY81Z83H0X)	
<b>MEDIUM-CHAIN TRIGLYCERIDES</b> (UNII: C9H2L21V7U)	
<b>CETOSTEARYL ALCOHOL</b> (UNII: 2DMT128M1S)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PHENOXYETHANOL</b> (UNII: HIE492ZZ3T)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>ETHYLHEXYLGLYCERIN</b> (UNII: 147D247K3P)	
<b>STEARETH-2</b> (UNII: V56DFE46J5)	

**Product Characteristics**

<b>Color</b>	white	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83294-101-18	118 mL in 1 TUBE; Type 0: Not a Combination Product	08/22/2022	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M020	08/22/2022	

**Labeler - CLEURE (116355659)**

