

**PAIN RELIEF PM- acetaminophen, aspirin, diphenhydramine citrate tablet, film coated**

**Amerisource Bergen**

-----

**Good Neighbor Pharmacy 44-674**

***Active ingredients (in each caplet)***

Acetaminophen 250 mg

Aspirin 250 mg (NSAID)\*

Diphenhydramine citrate 38 mg

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever

Pain reliever

Nighttime sleep-aid

***Uses***

for the temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- hives
- shock
- facial swelling
- asthma (wheezing)

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters
- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours

- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have 3 or more alcoholic drinks every day while using this product
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- take more or for a longer time than directed
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

### **Do not use**

- if you have ever had an allergic reaction to acetaminophen, aspirin, or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other product containing diphenhydramine, even one used on skin
- in children under 12 years of age

### **Ask a doctor before use if**

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have glaucoma
- you have asthma
- you have difficulty in urination due to enlargement of the prostate gland
- you are taking a diuretic
- you have a breathing problem such as emphysema or chronic bronchitis
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

### **Ask a doctor or pharmacist before use if you are taking**

- a prescription drug for:
  - diabetes
  - gout
  - arthritis
- any other drug, or are under a doctor's care for any serious condition
- sedatives or tranquilizers
- any product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

### **When using this product**

- drowsiness may occur
- do not drive a motor vehicle or operate machinery
- avoid alcoholic beverages

### **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts for more than 10 days
- painful area is red or swollen
- new symptoms occur
- ringing in the ears or a loss of hearing occurs
- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness.

These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of reach of children.**

In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### ***Directions***

- **do not take more than directed**
- adults and children 12 years and over: take 2 caplets at bedtime, with a full glass of water
- do not take more than 2 caplets in 24 hours, unless directed by a doctor
- children under 12 years: do not use

### ***Other information***

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

### ***Inactive ingredients***

corn starch, crospovidone, FD&C blue #1 aluminum lake, hypromellose, maltodextrin, medium chain triglycerides, microcrystalline cellulose, polydextrose, povidone, sodium lauryl sulfate, sodium starch glycolate, sodium sulfate, stearic acid, talc, titanium dioxide, yellow iron oxide, zinc stearate

### ***Questions or comments?***

**1-800-426-9391**

***Principal display panel***

**GOOD  
NEIGHBOR  
PHARMACY®**

Compare to Excedrin® PM  
Headache active ingredients†

NDC 46122-456-71

**Pain Relief PM  
acetaminophen**  
aspirin (**NSAID**) & diphenhydramine citrate

*Pain Reliever / Nighttime Sleep Aid*

•**Caffeine free**

**50** Caplets † • Non-habit forming

Actual Size

**TAMPER EVIDENT: DO NOT USE IF  
IMPRINTED SAFETY SEAL UNDER  
CAP IS BROKEN OR MISSING**

†This product is not manufactured or distributed by  
GSK Consumer Healthcare SARL, owner of the  
registered trademark Excedrin® PM Headache.

50844 REV1121A67415

Distributed By  
AmerisourceBergen  
1 West First Avenue  
Conshohocken, PA 19428  
Questions or Concerns?  
[www.mygnp.com](http://www.mygnp.com)

**GOOD NEIGHBOR PHARMACY BRAND PRODUCTS  
SATISFACTION GUARANTEED**

No Print Area  
Lot no. & Exp. Date

ABC#: 10182647  
8770143010  
7

TAMPER EVIDENT: DO NOT USE IF  
IMPRINTED SAFETY SEAL UNDER  
CAP IS BROKEN OR MISSING

**Drug Facts** **KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION**

**Active ingredients (in each caplet)**  
 Acetaminophen 250 mg..... Pain reliever  
 Aspirin 250 mg (NSAID)..... Pain reliever  
 Diphenhydramine citrate 38 mg..... Nighttime sleep-aid  
 • nonsteroidal anti-inflammatory drug

**Uses** for the temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness

**Warnings**  
 Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:  
 • shock • facial swelling • asthma (wheezing)  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:  
 • blisters • rash • skin reddening  
 If a skin reaction occurs, stop use and seek medical help right away.  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:  
 • more than 4,000 mg of acetaminophen in 24 hours  
 • with other drugs containing acetaminophen  
 • 3 or more alcoholic drinks every day while using this product  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you • are age 60 or older • have 3 or more alcoholic drinks every day while using this product • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, ...)

**GOOD NEIGHBOR PHARMACY**

NDC 46122-456-71

**Pain Relief PM** NIGHT TIME  
 acetaminophen  
 aspirin (NSAID) & diphenhydramine citrate  
 Pain Reliever / Nighttime Sleep Aid  
 • Caffeine free  
 50 Caplets | • Non-habit forming Actual Size

Compare to Excedrin<sup>®</sup> PM  
Headache active ingredients!

**AMERICAN PHARMACEUTICAL CORPORATION**

1-800-426-9391  
 Questions or comments?  
 Questions or Concerns?  
 www.mygnp.com

Distributed By  
 AmerisourceBergen  
 1 West First Avenue  
 Conshohocken, PA 19428

95844 REV112167 415  
 registered trademark Excedrin<sup>®</sup> PM Headache,  
 GSK Consumer Healthcare SArL, owner of the  
 This product is not manufactured or distributed by

**Drug Facts (continued)**

B-0045-674-15  
REV112167415

**Drug Facts (continued)**

• ringing in the ears or a loss of hearing occurs  
 • dizziness or lightheadedness  
 • more than 2 weeks of continuous use  
 • serious drowsiness or dizziness  
 • pain relief does not last for more than 10 days  
 • redness or swelling of the mouth  
 • these could be signs of a serious condition.  
**If pregnant or breastfeeding:** Tell your health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
**Keep out of reach of children.** In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**  
 • do not take more than directed  
 • adults and children 12 years and over: take 2 caplets at bedtime, with a full glass of water  
 • do not take more than 2 caplets in 24 hours, unless directed by a doctor  
 • children under 12 years: do not use

**Other information**  
 • store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)  
 • see end flap for expiration date and lot number

**Inactive ingredients** corn starch, croscopolone, FD&C blue #1, aluminum lake, hydroxypropylcellulose, maltodextrin, medium chain triglycerides, microcrystalline cellulose, polydextrose, novolone, sodium lauryl sulfate, sodium starch glycolate, sodium sulfate, stearic acid, talc, titanium dioxide, yellow iron oxide, zinc stearate

**Drug Facts (continued)**

• take more or for a longer time than directed  
 • have had stomach ulcers or bleeding problems  
 • take a blood thinning (anticoagulant) drug  
 • have ever had an allergic reaction to acetaminophen, aspirin, or any other pain reliever/fever reducer  
 • are now taking or have taken any other drug containing acetaminophen, aspirin, or diphenhydramine  
 • are now taking or have taken any other product containing diphenhydramine, even one used on skin  
 • in children under 12 years of age

**Do not use**  
 • if you have liver disease  
 • if you have a history of stomach problems, such as heartburn  
 • if you have glaucoma  
 • if you have asthma  
 • if you have difficulty in urination due to enlargement of the prostate gland  
 • if you are taking a diuretic  
 • if you have a breathing problem such as emphysema or chronic bronchitis  
 • if you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

**Ask a doctor before use if**  
 • you have liver disease  
 • you have a history of stomach problems, such as heartburn  
 • you have glaucoma  
 • you have asthma  
 • you have difficulty in urination due to enlargement of the prostate gland  
 • you are taking a diuretic  
 • you have a breathing problem such as emphysema or chronic bronchitis  
 • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

**Ask a doctor or pharmacist before use if you are taking**  
 • a prescription drug for:  
 • arthritis  
 • diabetes  
 • gout  
 • any other drug, or are under a doctor's care for any serious condition  
 • sedatives or tranquilizers  
 • any product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

**When using this product**  
 • avoid alcoholic beverages  
 • do not drive a motor vehicle or operate machinery  
 • drowsiness may occur  
 • an allergic reaction occurs. Seek medical help right away.  
 • you experience any of the following signs of stomach bleeding:  
 • feel faint  
 • vomit blood  
 • have bloody or black stools  
 • have stomach pain that does not get better

**Good Neighbor Pharmacy 44-674**

<b>PAIN RELIEF PM</b> acetaminophen, aspirin, diphenhydramine citrate tablet, film coated			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:46122-456
<b>Route of Administration</b>	ORAL		
<b>Active Ingredient/Active Moiety</b>			

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
<b>DIPHENHYDRAMINE CITRATE</b> (UNII: 4OD433S209) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE CITRATE	38 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>CROSPVIDONE, UNSPECIFIED</b> (UNII: 2S7830E561)	
<b>FD&amp;C BLUE NO. 1 ALUMINUM LAKE</b> (UNII: J9EQA3S2JM)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)	
<b>MEDIUM-CHAIN TRIGLYCERIDES</b> (UNII: C9H2L21V7U)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYDEXTROSE</b> (UNII: VH2XOU12IE)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>SODIUM SULFATE</b> (UNII: 0YPR65R21J)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)	
<b>ZINC STEARATE</b> (UNII: H92E6QA4FV)	

### Product Characteristics

<b>Color</b>	blue	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	674
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46122-456-71	1 in 1 CARTON	08/31/2016	
1		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	08/31/2016	

**Labeler** - Amerisource Bergen (007914906)**Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		038154464	pack(46122-456)

**Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		832867837	manufacture(46122-456)

**Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		832867894	manufacture(46122-456)

**Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		967626305	pack(46122-456)

**Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		117025878	manufacture(46122-456)

Revised: 1/2024

Amerisource Bergen