

# **CYCLOBENZAPRINE HYDROCHLORIDE- cyclobenzaprine hydrochloride capsule, extended release**

**Quality Care Products, LLC**

-----

## **HIGHLIGHTS OF PRESCRIBING INFORMATION**

**These highlights do not include all the information needed to use CYCLOBENZAPRINE HYDROCHLORIDE EXTENDED-RELEASE CAPSULES safely and effectively. See full prescribing information for CYCLOBENZAPRINE HYDROCHLORIDE EXTENDED-RELEASE CAPSULES.**

### **CYCLOBENZAPRINE HYDROCHLORIDE extended-release capsules, for oral use**

**Initial U.S. Approval: 1977**

#### **----- INDICATIONS AND USAGE -----**

Cyclobenzaprine hydrochloride extended-release capsules are muscle relaxant indicated as an adjunct to rest and physical therapy for relief of muscle spasm associated with acute, painful musculoskeletal conditions. (1)

##### Limitations of Use:

- Cyclobenzaprine hydrochloride extended-release capsules should be used only for short periods (up to 2 or 3 weeks) (1)
- Cyclobenzaprine hydrochloride extended-release capsules have not been found effective in the treatment of spasticity or cerebral palsy (1)

#### **----- DOSAGE AND ADMINISTRATION -----**

- Recommended adult dose for most patients is 15 mg taken once daily. Some patients may require 30 mg taken once daily (2)
- Recommended to take doses at approximately same time each day (2)
- Instruct patients to swallow cyclobenzaprine hydrochloride extended-release capsules intact or to sprinkle capsule contents on a tablespoon of applesauce and swallow immediately without chewing (2)
- Use for periods longer than 2 or 3 weeks is not recommended (2)

#### **----- DOSAGE FORMS AND STRENGTHS -----**

- Extended-release capsules: 15 and 30 mg (3)

#### **----- CONTRAINDICATIONS -----**

- Hypersensitivity to any component of this product (4)
- Concomitant use of monoamine oxidase (MAO) inhibitors or within 14 days after their discontinuation (4)
- During acute recovery phase of myocardial infarction, and in patients with arrhythmias, heart block or conduction disturbances, or congestive heart failure (4)
- Hyperthyroidism (4)

#### **----- WARNINGS AND PRECAUTIONS -----**

- Serotonin syndrome has been reported with cyclobenzaprine when used in combination with other serotonergic drugs (5.1)
- Cyclobenzaprine is structurally related to tricyclic antidepressants which have been reported to produce adverse cardiovascular effects or CNS depressant effects (5.2)
- Use in the elderly is not recommended (5.3)
- Use in patients with hepatic impairment is not recommended (5.4)
- Use with caution in patients with a history of urinary retention, angle-closure glaucoma, increased intraocular pressure and in patients taking anticholinergic medications (5.5)

#### **----- ADVERSE REACTIONS -----**

Most common adverse reactions (incidence  $\geq 3\%$  in any treatment group and greater than placebo): dry mouth, dizziness, fatigue, constipation, nausea, dyspepsia, and somnolence (6)

**To report SUSPECTED ADVERSE REACTIONS, contact TWi Pharmaceuticals at 1-844-518-2989 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

#### **----- DRUG INTERACTIONS -----**

- MAO Inhibitors: Life-threatening interactions may occur (4, 7)
- Serotonergic Drugs: Serotonin syndrome has been reported (5.1, 7)
- CNS Depressants: Effects of alcohol, barbiturates, and other CNS depressants may be enhanced (5.2, 7)

- Tramadol: Seizure risk may be enhanced (7)
- Guanethidine: Antihypertensive effect may be blocked (7)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 4/2020

---

## **FULL PRESCRIBING INFORMATION: CONTENTS\***

### **1 INDICATIONS AND USAGE**

### **2 DOSAGE AND ADMINISTRATION**

### **3 DOSAGE FORMS AND STRENGTHS**

### **4 CONTRAINDICATIONS**

### **5 WARNINGS AND PRECAUTIONS**

- 5.1 Serotonin Syndrome
- 5.2 Tricyclic Antidepressant-like Effects
- 5.3 Use in the Elderly
- 5.4 Use in Patients with Hepatic Impairment
- 5.5 Atropine-like Action

### **6 ADVERSE REACTIONS**

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

### **7 DRUG INTERACTIONS**

### **8 USE IN SPECIFIC POPULATIONS**

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Hepatic Impairment

### **9 DRUG ABUSE AND DEPENDENCE**

- 9.3 Dependence

### **10 OVERDOSAGE**

- 10.1 Manifestations
- 10.2 Management

### **11 DESCRIPTION**

### **12 CLINICAL PHARMACOLOGY**

- 12.1 Mechanism of Action
- 12.3 Pharmacokinetics

### **13 NONCLINICAL TOXICOLOGY**

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
- 13.2 Animal Toxicology and/or Pharmacology

### **14 CLINICAL STUDIES**

### **16 HOW SUPPLIED/STORAGE AND HANDLING**

- 16.1 How Supplied
- 16.2 Storage and Handling

### **17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## **FULL PRESCRIBING INFORMATION**

## 1 INDICATIONS AND USAGE

Cyclobenzaprine hydrochloride extended-release capsules are indicated as an adjunct to rest and physical therapy for relief of muscle spasm associated with acute, painful musculoskeletal conditions. Improvement is manifested by relief of muscle spasm and its associated signs and symptoms, namely, pain, tenderness, and limitation of motion.

### Limitations of Use:

- Cyclobenzaprine hydrochloride extended-release capsules should be used only for short periods (up to two or three weeks) because adequate evidence of effectiveness for more prolonged use is not available and because muscle spasm associated with acute, painful musculoskeletal conditions is generally of short duration and specific therapy for longer periods is seldom warranted.
- Cyclobenzaprine hydrochloride extended-release capsules have not been found effective in the treatment of spasticity associated with cerebral or spinal cord disease or in children with cerebral palsy.

## 2 DOSAGE AND ADMINISTRATION

The recommended adult dose for most patients is one (1) cyclobenzaprine hydrochloride extended-release capsule, 15 mg taken once daily. Some patients may require up to 30 mg/day, given as one (1) cyclobenzaprine hydrochloride extended-release capsule, 30 mg taken once daily or as two (2) cyclobenzaprine hydrochloride extended-release capsules, 15 mg taken once daily.

- It is recommended that doses be taken at approximately the same time each day.
- Use of cyclobenzaprine hydrochloride extended-release capsules for periods longer than two or three weeks is not recommended [*see Indications and Usage (1)*].

Instruct patients to swallow cyclobenzaprine hydrochloride extended-release capsules intact. Alternatively, the contents of the cyclobenzaprine hydrochloride extended-release capsules may be sprinkled over applesauce and then swallowed. This method is appropriate only for patients able to reliably swallow the applesauce without chewing. Other foods have not been tested and should not be substituted for applesauce. Instruct the patient to:

- Sprinkle the contents of the capsule onto a tablespoon of applesauce and consume immediately without chewing.
- Rinse the mouth to ensure all of the contents have been swallowed.
- Discard any unused portion of the cyclobenzaprine hydrochloride extended-release capsules after the contents have been sprinkled on applesauce.

## 3 DOSAGE FORMS AND STRENGTHS

Extended-release capsules USP in the following strengths:

- 15 mg: Capsules are dark orange cap/red opaque body and are imprinted with black ink with “T035” on the body and cap.
- 30 mg: Capsules are orange opaque cap/white opaque body and are imprinted with black ink with “T036” on the body and cap.

## 4 CONTRAINDICATIONS

- Hypersensitivity to any component of this product. These adverse reactions may manifest as an anaphylactic reaction, urticaria, facial and/or tongue swelling, or pruritus. Discontinue cyclobenzaprine hydrochloride extended-release capsules if a hypersensitivity reaction is suspected.
- Concomitant use of monoamine oxidase (MAO) inhibitors or within 14 days after their discontinuation. Hyperpyretic crisis seizures and deaths have occurred in patients receiving

cyclobenzaprine (or structurally similar tricyclic antidepressants) concomitantly with MAO inhibitor drugs.

- During the acute recovery phase of myocardial infarction, and in patients with arrhythmias, heart block or conduction disturbances, or congestive heart failure.
- Hyperthyroidism.

## **5 WARNINGS AND PRECAUTIONS**

### **5.1 Serotonin Syndrome**

The development of a potentially life-threatening serotonin syndrome has been reported with cyclobenzaprine when used in combination with other drugs, such as selective serotonin reuptake inhibitors (SSRIs), serotonin norepinephrine reuptake inhibitors (SNRIs), tricyclic antidepressants (TCAs), tramadol, bupropion, meperidine, verapamil, or MAO inhibitors. The concomitant use of cyclobenzaprine hydrochloride extended-release capsules with MAO inhibitors is contraindicated [*see Contraindications (4)*]. Serotonin syndrome symptoms may include mental status changes (e.g., confusion, agitation, hallucinations), autonomic instability (e.g., diaphoresis, tachycardia, labile blood pressure, hyperthermia), neuromuscular abnormalities (e.g., tremor, ataxia, hyperreflexia, clonus, muscle rigidity), and/or gastrointestinal symptoms (e.g., nausea, vomiting, diarrhea). Treatment with cyclobenzaprine hydrochloride extended-release capsules and any concomitant serotonergic agents should be discontinued immediately if the above reactions occur and supportive symptomatic treatment should be initiated. If concomitant treatment with cyclobenzaprine hydrochloride extended-release capsules and other serotonergic drugs is clinically warranted, careful observation is advised, particularly during treatment initiation or dose increases.

### **5.2 Tricyclic Antidepressant-like Effects**

Cyclobenzaprine is structurally related to the tricyclic antidepressants, e.g., amitriptyline and imipramine. Tricyclic antidepressants have been reported to produce arrhythmias, sinus tachycardia, prolongation of the conduction time leading to myocardial infarction and stroke [*see Contraindications (4)*]. Cyclobenzaprine hydrochloride extended-release capsules may enhance the effects of alcohol, barbiturates, and other CNS depressants.

Some of the more serious central nervous system (CNS) reactions noted with the tricyclic antidepressants have occurred in short-term studies of cyclobenzaprine for indications other than muscle spasm associated with acute musculoskeletal conditions, and usually at doses somewhat greater than those recommended for skeletal muscle spasm. If clinically significant CNS symptoms develop, consider discontinuation of cyclobenzaprine hydrochloride extended-release capsules.

### **5.3 Use in the Elderly**

As a result of a 40% increase in cyclobenzaprine plasma levels and a 56% increase in plasma half-life following administration of cyclobenzaprine hydrochloride extended-release capsules in elderly subjects as compared to young adults, use of cyclobenzaprine hydrochloride extended-release capsules is not recommended in the elderly [*see Clinical Pharmacology (12.3)*].

### **5.4 Use in Patients with Hepatic Impairment**

As a result of two-fold higher cyclobenzaprine plasma levels in subjects with mild hepatic impairment, as compared to healthy subjects, following administration of immediate-release cyclobenzaprine and because there is limited dosing flexibility with cyclobenzaprine hydrochloride extended-release capsules, use of cyclobenzaprine hydrochloride extended-release capsules is not recommended in patients with mild, moderate, or severe hepatic impairment [*see Clinical Pharmacology (12.3)*].

### **5.5 Atropine-like Action**

Because of its atropine-like action, cyclobenzaprine hydrochloride extended-release capsules should be used with caution in patients with a history of urinary retention, angle-closure glaucoma, increased intraocular pressure, and in patients taking anticholinergic medication.

## 6 ADVERSE REACTIONS

The following clinically significant reactions are described in greater detail, in other sections.

- Serotonin Syndrome [see *Warnings and Precautions (5.1)*]
- Adverse Cardiovascular Effects [see *Warnings and Precautions (5.2)*]

### 6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

The data described below reflect exposure to cyclobenzaprine hydrochloride extended-release capsules in 253 patients in 2 clinical trials. Cyclobenzaprine hydrochloride extended-release capsules were studied in two double-blind, parallel-group, placebo-controlled, active-controlled trials of identical design [see *Clinical Studies (14)*]. The study population was composed of patients with muscle spasms associated with acute painful musculoskeletal conditions. Patients received 15 mg or 30 mg of cyclobenzaprine hydrochloride extended-release capsules taken orally once daily, cyclobenzaprine immediate-release (IR) 10 mg three times a day, or placebo for 14 days.

The most common adverse reactions (incidence  $\geq 3\%$  in any treatment group and greater than placebo) were dry mouth, dizziness, fatigue, constipation, nausea, dyspepsia, and somnolence (see Table 1).

**Table 1: Incidence of the Most Common Adverse Reactions Occurring in  $\geq 3\%$  of Patients in any Treatment Group\* and Greater Than Placebo in the Two Phase 3, Double-Blind Cyclobenzaprine Hydrochloride Extended-Release Capsules Trials**

	<b>Placebo</b>  <b>N=128</b>	<b>Cyclobenzaprine Hydrochloride Extended-Release Capsules 15 mg</b>  <b>N=127</b>	<b>Cyclobenzaprine Hydrochloride Extended-Release Capsules 30 mg</b>  <b>N=126</b>
Dry mouth	2%	6%	14%
Dizziness	2%	3%	6%
Fatigue	2%	3%	3%
Constipation	0%	1%	3%
Somnolence	0%	1%	2%
Nausea	1%	3%	3%
Dyspepsia	1%	0%	4%

\*Cyclobenzaprine Hydrochloride Extended-Release Capsules 15 mg QD, Cyclobenzaprine Hydrochloride Extended-Release Capsules 30 mg QD, or cyclobenzaprine IR tablets TID

### 6.2 Postmarketing Experience

The following adverse reactions have been reported in clinical studies or postmarketing experience with cyclobenzaprine hydrochloride extended-release capsules, cyclobenzaprine IR, or tricyclic drugs. Because some of these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

In a postmarketing surveillance program of cyclobenzaprine IR, the adverse reactions reported most

frequently were drowsiness, dry mouth, and dizziness and adverse reactions reported in 1% to 3% of the patients were: fatigue/tiredness, asthenia, nausea, constipation, dyspepsia, unpleasant taste, blurred vision, headache, nervousness, and confusion.

The following adverse reactions have been reported in postmarketing experience (cyclobenzaprine hydrochloride extended-release capsules or cyclobenzaprine IR), in clinical studies of cyclobenzaprine IR (incidence <1%), or in postmarketing experience with other tricyclic drugs:

*Body as a Whole:* Syncope; malaise; chest pain; edema.

*Cardiovascular:* Tachycardia; arrhythmia; vasodilatation; palpitation; hypotension; hypertension; myocardial infarction; heart block; stroke.

*Digestive:* Vomiting; anorexia; diarrhea; gastrointestinal pain; gastritis; thirst; flatulence; edema of the tongue; abnormal liver function and rare reports of hepatitis, jaundice, and cholestasis; paralytic ileus, tongue discoloration; stomatitis; parotid swelling.

*Endocrine:* Inappropriate ADH syndrome.

*Hematologic and Lymphatic:* Purpura; bone marrow depression; leukopenia; eosinophilia; thrombocytopenia.

*Hypersensitivity:* Anaphylaxis; angioedema; pruritus; facial edema; urticaria; rash.

*Metabolic, Nutritional, and Immune:* Elevation and lowering of blood sugar levels; weight gain or loss.

*Musculoskeletal:* Local weakness; myalgia.

*Nervous System and Psychiatric:* Seizures, ataxia; vertigo; dysarthria; tremors; hypertonia; convulsions; muscle twitching; disorientation; insomnia; depressed mood; abnormal sensations; anxiety; agitation; psychosis, abnormal thinking and dreaming; hallucinations; excitement; paresthesia; diplopia; serotonin syndrome; neuroleptic malignant syndrome; decreased or increased libido; abnormal gait; delusions; aggressive behavior; paranoia; peripheral neuropathy; Bell's palsy; alteration in EEG patterns; extrapyramidal symptoms.

*Respiratory:* Dyspnea.

*Skin:* Sweating; photosensitization; alopecia.

*Special Senses:* Ageusia; tinnitus.

*Urogenital:* Urinary frequency and/or retention; impaired urination; dilatation of urinary tract; impotence; testicular swelling; gynecomastia; breast enlargement; galactorrhea.

## **7 DRUG INTERACTIONS**

Based on its structural similarity to tricyclic antidepressants, cyclobenzaprine hydrochloride extended-release capsules may have life-threatening interactions with MAO inhibitors [see *Contraindications (4)*], may enhance the effects of alcohol, barbiturates, and other CNS depressants, may enhance the seizure risk in patients taking tramadol, or may block the antihypertensive action of guanethidine and similarly acting compounds.

Postmarketing cases of serotonin syndrome have been reported during combined use of cyclobenzaprine and other drugs, such as SSRIs, SNRIs, TCAs, tramadol, bupropion, meperidine, verapamil, or MAO inhibitors [see *Warnings and Precautions (5.1)*].

## **8 USE IN SPECIFIC POPULATIONS**

### **8.1 Pregnancy**

#### Risk Summary

Available data from case reports with cyclobenzaprine hydrochloride extended-release capsules use in pregnancy have not identified a drug-associated risk of major birth defects, miscarriage, or adverse maternal or fetal outcomes. In rats, decreased pup body weight and survival was noted at cyclobenzaprine doses  $\geq 10$  mg/kg/day (approximately  $\geq 3$  times the maximum recommended human dose (MRHD) of 30 mg/day), when administered orally during pregnancy and lactation (*see Data*).

The estimated background risk of major birth defects and miscarriage for the indicated populations is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the US general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively.

## Data

### *Animal Data*

No adverse embryofetal effects were reported following oral administration of cyclobenzaprine during organogenesis to mice and rabbits at maternal doses up to 20 mg/kg/day (approximately 3 and 15 times the MRHD, respectively, on a mg/m<sup>2</sup> basis). Maternal toxicity characterized by decreased body weight gain was observed only in mice at the highest tested dose of 20 mg/kg/day.

Decreased pup body weight and survival were reported in a prenatal and postnatal study where pregnant rats were treated orally with cyclobenzaprine during pregnancy and lactation with maternal doses of 10 and 20 mg/kg/day (approximately 3 and 6 times the MRHD on a mg/m<sup>2</sup> basis). Maternal toxicity, characterized by a decreased body weight gain, was observed only at the highest tested dose of 20 mg/kg/day.

## **8.2 Lactation**

### Risk Summary

There are no data on the presence of cyclobenzaprine in either human or animal milk, the effects on a breastfed infant, or the effects on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for cyclobenzaprine hydrochloride extended-release capsules and any potential adverse effects on the breastfed child from cyclobenzaprine hydrochloride extended-release capsules or from the underlying maternal condition.

## **8.4 Pediatric Use**

Safety and effectiveness of cyclobenzaprine hydrochloride extended-release capsules have not been studied in pediatric patients.

## **8.5 Geriatric Use**

Clinical studies of cyclobenzaprine hydrochloride extended-release capsules did not include sufficient numbers of patients aged 65 and over to determine the safety and efficacy of cyclobenzaprine hydrochloride extended-release capsules in the elderly population. The plasma concentration and half-life of cyclobenzaprine are substantially increased in the elderly when compared to the general patient population. Accordingly, use of cyclobenzaprine hydrochloride extended-release capsules is not recommended in the elderly [*see Warnings and Precautions (5.3) and Clinical Pharmacology (12.3)*].

## **8.6 Hepatic Impairment**

The use of cyclobenzaprine hydrochloride extended-release capsules is not recommended in patients with mild, moderate, or severe hepatic impairment [*see Warnings and Precautions (5.4) and Clinical Pharmacology (12.3)*].

# **9 DRUG ABUSE AND DEPENDENCE**

## **9.3 Dependence**

Pharmacologic similarities among the tricyclic drugs require that certain withdrawal symptoms be considered when cyclobenzaprine hydrochloride extended-release capsules are administered, even though they have not been reported to occur with this drug. Abrupt cessation of treatment after prolonged administration rarely may produce nausea, headache, and malaise. These are not indicative of addiction.

## 10 OVERDOSAGE

### 10.1 Manifestations

Although rare, deaths may occur from overdosage with cyclobenzaprine hydrochloride extended-release capsules. Multiple drug ingestion (including alcohol) is common in deliberate cyclobenzaprine overdose. As management of overdose is complex and changing, it is recommended that the physician contact a poison control center for current information on treatment. Signs and symptoms of toxicity may develop rapidly after cyclobenzaprine overdose; therefore, hospital monitoring is required as soon as possible.

The most common effects associated with cyclobenzaprine overdose are drowsiness and tachycardia. Less frequent manifestations include tremor, agitation, coma, ataxia, hypertension, slurred speech, confusion, dizziness, nausea, vomiting, and hallucinations. Rare but potentially critical manifestations of overdose are cardiac arrest, chest pain, cardiac dysrhythmias, severe hypotension, seizures, and neuroleptic malignant syndrome. Changes in the electrocardiogram, particularly in QRS axis or width, are clinically significant indicators of cyclobenzaprine toxicity. Other potential effects of overdosage include any of the symptoms listed under *Adverse Reactions* (6).

### 10.2 Management

#### General

As management of overdose is complex and changing, it is recommended that the physician contact a poison control center for current information on treatment.

In order to protect against the rare but potentially critical manifestations described above, obtain an ECG and immediately initiate cardiac monitoring. Protect the patient's airway, establish an intravenous line, and initiate gastric decontamination. Observation with cardiac monitoring and observation for signs of CNS or respiratory depression, hypotension, cardiac dysrhythmias and/or conduction blocks, and seizures is necessary. If signs of toxicity occur at any time during this period, extended monitoring is required. Monitoring of plasma drug levels should not guide management of the patient. Dialysis is probably of no value because of low plasma concentrations of the drug.

#### Gastrointestinal Decontamination

All patients suspected of an overdose with cyclobenzaprine hydrochloride extended-release capsules should receive gastrointestinal decontamination. This should include large volume gastric lavage followed by activated charcoal. If consciousness is impaired, the airway should be secured prior to lavage and emesis is contraindicated.

#### Cardiovascular

A maximal limb-lead QRS duration of 0.10 seconds may be the best indication of the severity of the overdose. Serum alkalization, to a pH of 7.45 to 7.55, using intravenous sodium bicarbonate and hyperventilation (as needed), should be instituted for patients with dysrhythmias and/or QRS widening. A pH >7.60 or a pCO<sub>2</sub> <20 mmHg is undesirable. Dysrhythmias unresponsive to sodium bicarbonate therapy/hyperventilation may respond to lidocaine, bretylium, or phenytoin. Type 1A and 1C antiarrhythmics are generally contraindicated (e.g., quinidine, disopyramide, and procainamide).

#### CNS



In patients with CNS depression, early intubation is advised because of the potential for abrupt deterioration. Seizures should be controlled with benzodiazepines or, if these are ineffective, other anticonvulsants (e.g., phenobarbital, phenytoin). Physostigmine is not recommended except to treat life-threatening symptoms that have been unresponsive to other therapies, and then only in close consultation with a poison control center.

### Psychiatric Follow-Up


Since overdose is often deliberate, patients may attempt suicide by other means during the recovery phase. Psychiatric referral may be appropriate.

### Pediatric Management

The principles of management of child and adult overdose are similar. It is strongly recommended that the physician contact the local poison control center for specific pediatric treatment.

## **11 DESCRIPTION**

Cyclobenzaprine hydrochloride extended-release capsules USP are skeletal muscle relaxant which relieves muscle spasm of local origin without interfering with muscle function. The active ingredient in cyclobenzaprine hydrochloride extended-release capsules is cyclobenzaprine hydrochloride, USP. Cyclobenzaprine hydrochloride (HCl) is a white, crystalline tricyclic amine salt with the empirical formula  $C_{20}H_{21}N \cdot HCl$  and a molecular weight of 311.9. It has a melting point of  $217^{\circ}C$ , and a  $pK_a$  of 8.47 at  $25^{\circ}C$ . It is freely soluble in water and alcohol, sparingly soluble in isopropanol, and insoluble in hydrocarbon solvents. If aqueous solutions are made alkaline, the free base separates. Cyclobenzaprine HCl is designated chemically as 3-(5H-dibenzo[a,d] cyclohepten-5-ylidene)-N,N-dimethyl-1-propanamine hydrochloride, and has the following structural formula:



Cyclobenzaprine hydrochloride extended-release capsules for oral administration are supplied in 15 and 30 mg strengths. Cyclobenzaprine hydrochloride extended-release capsules contain the following inactive ingredients: colloidal silicon dioxide, ethyl alcohol, ethylcellulose, FDC yellow #6, gelatin, hydroxypropyl cellulose, isopropyl alcohol, potassium hydroxide, sugar spheres (which contain sucrose and corn starch) and titanium dioxide. In addition, red iron oxide is also included in the 15 mg strength. The capsule is imprinted with black ink consisting of: black iron oxide, DC Yellow #10, ethanol, FDC Blue #1, FDC Blue #2, FDC Red #40, methanol, N-butyl alcohol, propylene glycol, and shellac.

## **12 CLINICAL PHARMACOLOGY**

### **12.1 Mechanism of Action**

Cyclobenzaprine relieves skeletal muscle spasm of local origin without interfering with muscle function. Cyclobenzaprine has not been shown to be effective in muscle spasm due to central nervous system disease. In animal models, cyclobenzaprine reduced or abolished skeletal muscle hyperactivity. Animal studies indicate that cyclobenzaprine does not act at the neuromuscular junction or directly on

skeletal muscle. Such studies show that cyclobenzaprine acts primarily within the central nervous system at the brain stem as opposed to the spinal cord level, although an overlapping action on the latter may contribute to its overall skeletal muscle relaxant activity. Evidence suggests that the net effect of cyclobenzaprine is a reduction of tonic somatic motor activity, influencing both gamma ( $\gamma$ ) and alpha ( $\alpha$ ) motor systems. Pharmacological studies in animals demonstrated a similarity between the effects of cyclobenzaprine and the structurally related tricyclic antidepressants, including reserpine antagonism, norepinephrine potentiation, potent peripheral and central anticholinergic effects, and sedation. Cyclobenzaprine caused slight to moderate increase in heart rate in animals.

## **12.3 Pharmacokinetics**

### **Absorption**

Following single-dose administration of cyclobenzaprine hydrochloride extended-release capsules 15 mg and 30 mg in healthy adult subjects ( $n=15$ ),  $C_{\max}$ ,  $AUC_{0-168h}$  and  $AUC_{0-\infty}$  increased in an approximately dose-proportional manner from 15 mg to 30 mg. The time to peak plasma cyclobenzaprine concentration ( $T_{\max}$ ) was 7 to 8 hours for both doses of cyclobenzaprine hydrochloride extended-release capsules.

A food effect study conducted in healthy adult subjects ( $n=15$ ) utilizing a single dose of AMRIX 30 mg demonstrated a statistically significant increase in bioavailability when cyclobenzaprine hydrochloride extended-release capsules 30 mg was given with food relative to the fasted state. There was a 35% increase in peak plasma cyclobenzaprine concentration ( $C_{\max}$ ) and a 20% increase in exposure ( $AUC_{0-168h}$  and  $AUC_{0-\infty}$ ) in the presence of food. No effect, however, was noted in  $T_{\max}$  or the shape of the mean plasma cyclobenzaprine concentration versus time profile. Cyclobenzaprine in plasma was first detectable in both the fed and fasted states at 1.5 hours.

When the contents of cyclobenzaprine hydrochloride extended-release capsules were administered by sprinkling on applesauce, it was found to be bioequivalent to the same dose when administered as an intact capsule.

In a multiple-dose study utilizing cyclobenzaprine hydrochloride extended-release capsules 30 mg administered once daily for 7 days in a group of healthy adult subjects ( $n=35$ ), a 2.5-fold accumulation of plasma cyclobenzaprine levels was noted at steady-state.

### **Metabolism and Excretion**

Cyclobenzaprine is extensively metabolized and is excreted primarily as glucuronides via the kidney. Cytochromes P-450 3A4, 1A2, and, to a lesser extent, 2D6, mediate N-demethylation, one of the oxidative pathways for cyclobenzaprine. Cyclobenzaprine has an elimination half-life of 32 hours (range 8-37 hours;  $n=18$ ); plasma clearance is 0.7 L/min following single-dose administration of cyclobenzaprine hydrochloride extended-release capsules.

### **Special Populations**

#### **Elderly**

Although there were no notable differences in  $C_{\max}$  or  $T_{\max}$ , cyclobenzaprine plasma AUC is increased by 40% and the plasma half-life of cyclobenzaprine is prolonged in elderly subjects greater than 65 years of age (50 hours) after dosing with cyclobenzaprine hydrochloride extended-release capsules compared to younger subjects 18 to 45 years of age (32 hours). Pharmacokinetic characteristics of cyclobenzaprine following multiple-dose administration of cyclobenzaprine hydrochloride extended-release capsules in the elderly were not evaluated.

#### **Hepatic Impairment**

In a pharmacokinetic study of immediate-release cyclobenzaprine in 16 subjects with hepatic impairment (15 mild, 1 moderate per Child-Pugh score), both AUC and  $C_{\max}$  were approximately double the values seen in the healthy control group. The pharmacokinetics of cyclobenzaprine in subjects with severe hepatic impairment is not known.

## 13 NONCLINICAL TOXICOLOGY

### 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

#### Carcinogenesis

Long-term studies were conducted in CD-1 mice and Sprague-Dawley rats with oral cyclobenzaprine to evaluate its carcinogenic potential. In an 81-week carcinogenicity study, metastatic hemangiosarcoma was seen in 3 of 21 male mice at 10 mg/kg/day (approximately 2 times the maximum recommended human dose (MRHD) of 30 mg/day on a mg/m<sup>2</sup> basis). In a 105-week carcinogenicity study, malignant astrocytoma was seen in 3 of 50 male rats at 10 mg/kg/day (approximately 3 times the MRHD on a mg/m<sup>2</sup> basis). There were no tumor findings in female mice or rats.

#### Mutagenesis

Cyclobenzaprine HCl was not mutagenic or clastogenic in the following assays: an *in vitro* Ames bacterial mutation assay, *in vitro* Chinese hamster ovary (CHO) cell chromosomal aberration test, and *in vivo* mouse bone marrow micronucleus assay.

#### Impairment of Fertility

Cyclobenzaprine HCl, when administered 70 and 14 days prior to mating to male and female rats, respectively, had no effects on fertility or reproductive performance at oral doses up to 20 mg/kg/day (approximately 6.5 times the MRHD on a mg/m<sup>2</sup> basis).

### 13.2 Animal Toxicology and/or Pharmacology

In a 67-week study with rats that received cyclobenzaprine at oral doses of 10, 20, or 40 mg/kg/day (3 to 15 times the MRHD on mg/m<sup>2</sup> basis), there were findings in the liver consisting of midzonal vacuolation with lipidosis for males and midzonal and centrilobular hepatocytic enlargement for females. In addition, there were findings of centrilobular coagulative necrosis. In the higher dose groups, these microscopic changes were seen after 26 weeks and even earlier in rats that died prior to 26 weeks; at lower doses, these changes were not seen until after 26 weeks.

In a 26-week study with Cynomolgus monkeys that received cyclobenzaprine at oral doses of 2.5, 5, 10, or 20 mg/kg/day, one monkey at 20 mg/kg/day (15 times the MRHD on mg/m<sup>2</sup> basis) was euthanized in week 17. Morbidity for this animal was attributed to findings of chronic pancreatitis, cholecystitis, cholangitis, and focal liver necrosis.

## 14 CLINICAL STUDIES

Efficacy was assessed in two double-blind, parallel-group, active-controlled, placebo-controlled studies of identical design of cyclobenzaprine hydrochloride extended-release capsules 15 mg and 30 mg taken once daily, between 6:00 and 7:00 PM, cyclobenzaprine 10 mg three times a day, or placebo for 14 days in patients with muscle spasms associated with acute painful musculoskeletal conditions.

There were significant differences in the primary efficacy analysis, the patient's rating of medication helpfulness, between the cyclobenzaprine hydrochloride extended-release capsules 15 mg group and the placebo group at Days 4 and 14 in one study and between the cyclobenzaprine hydrochloride extended-release capsules 30 mg group and the placebo group at Day 4 in the second study.

**Table 2: Patients' Rating of Medication Helpfulness - Study 1\***

	Day 4		Day 14	
	Number of Patients (%)		Number of Patients (%)	
	Placebo (N = 64)	Cyclobenzaprine Hydrochloride	Placebo (N = 64)	Cyclobenzaprine Hydrochloride

		<b>Extended-Release Capsules 30 mg (N = 64)</b>		<b>Extended-Release Capsules 30 mg (N = 64)</b>
Excellent	1 (2%)	3 (5%)	12 (19%)	15 (23%)
Very Good	5 (8%)	13 (20%)	9 (14%)	19 (30%)
Good	15 (23%)	22 (34%)	10 (16%)	15 (23%)
Fair	24 (38%)	20 (31%)	16 (25%)	10 (16%)
Poor	10 (16%)	5 (8%)	9 (14%)	4 (6%)
Missing	9 (14%)	1 (2%)	8 (13%)	1 (2%)

\* Percentages are rounded to the nearest whole percent.

**Table 3: Patients' Rating of Medication Helpfulness - Study 2\***

	<b>Day 4</b>		<b>Day 14</b>	
	<b>Number of Patients (%)</b>		<b>Number of Patients (%)</b>	
	<b>Placebo (N = 64)</b>	<b>Cyclobenzaprine Hydrochloride Extended-Release Capsules 15 mg (N = 63)</b>	<b>Placebo (N = 64)</b>	<b>Cyclobenzaprine Hydrochloride Extended-Release Capsules 15 mg (N = 63)</b>
Excellent	1 (2%)	2 (3%)	10 (16%)	13 (21%)
Very Good	10 (16%)	12 (19%)	12 (19%)	21 (33%)
Good	14 (22%)	21 (33%)	13 (20%)	9 (14%)
Fair	16 (25%)	17 (27%)	14 (22%)	10 (16%)
Poor	19 (30%)	6 (10%)	12 (19%)	5 (8%)
Missing	4 (6%)	5 (8%)	3 (5%)	5 (8%)

\* Percentages are rounded to the nearest whole percent.

In addition, one of the two studies demonstrated significant differences between the cyclobenzaprine hydrochloride extended-release capsules 30 mg group and the placebo group in terms of patient-rated relief from local pain due to muscle spasm at Day 4 and Day 8, in patient-rated restriction of movement at Day 4 and Day 8, and in patient-rated global impression of change at Day 4, Day 8, and Day 14.

In both studies, there were no significant treatment differences between the cyclobenzaprine hydrochloride extended-release capsules treatment groups and the placebo group in physician's global assessment, patient-rated restriction in activities of daily living, or quality of nighttime sleep.

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

### **16.1 How Supplied**

Cyclobenzaprine hydrochloride extended-release capsules USP are available in 15 and 30 mg strengths, packaged in bottles of 60 capsules.

Cyclobenzaprine hydrochloride extended-release capsules USP, 15 mg are available as hard gelatin capsule with dark orange opaque cap and red opaque body imprinted with "T035" in black ink. They are supplied as follows:

Bottle of 60 NDC 55700-857-60

Cyclobenzaprine hydrochloride extended-release capsules USP, 30 mg are available as hard gelatin capsule with orange opaque cap and white opaque body imprinted with "T036" in black ink. They are

supplied as follows:

Bottle of 60 NDC 55700-858-60

## 16.2 Storage and Handling

Dispense in a tight, light-resistant container as defined in the USP/NF.

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].

## 17 PATIENT COUNSELING INFORMATION

See FDA-approved patient labeling (Patient Information).

- Instruct patients to swallow cyclobenzaprine hydrochloride extended-release capsules intact or to sprinkle capsule contents on a tablespoon of applesauce and swallow immediately without chewing.
- Advise patients to stop taking cyclobenzaprine hydrochloride extended-release capsules and to notify their physician right away if they experience symptoms of an allergic reaction, such as difficulty breathing, hives, swelling of face or tongue, or itching.
- Advise patients that cyclobenzaprine hydrochloride extended-release capsules should not be taken with MAO inhibitors or within 14 days after their discontinuation.
- Caution patients about the risk of serotonin syndrome with concomitant use of cyclobenzaprine hydrochloride extended-release capsules and other drugs, such as SSRIs, SNRIs, TCAs, tramadol, bupropion, meperidine, verapamil, or MAO inhibitors. Advise patients of the signs and symptoms of serotonin syndrome [see *Warnings and Precautions (5.1)*] and instruct patients to seek medical care immediately if they experience these symptoms.
- Advise patients to stop taking cyclobenzaprine hydrochloride extended-release capsules and to notify their physician right away if they experience arrhythmias or tachycardia.
- Advise patients that cyclobenzaprine hydrochloride extended-release capsules may enhance the impairment effects of alcohol. These effects may also be seen if cyclobenzaprine hydrochloride extended-release capsules are taken with other CNS depressants.
- Caution patients about operating an automobile or other hazardous machinery until it is reasonably certain that cyclobenzaprine hydrochloride extended-release capsules therapy will not adversely affect their ability to engage in such activities.
- Advise patients to take cyclobenzaprine hydrochloride extended-release capsules at approximately the same time each day.

Manufactured for:

TWi Pharmaceuticals USA, Inc.

Paramus, NJ 07652

Manufactured by:

TWi Pharmaceuticals, Inc.

Taoyuan City, 32063, Taiwan

AMRIX is a trademark of Teva Pharmaceuticals International GmbH, or its affiliates.

## Patient Information

---

### PATIENT INFORMATION

Cyclobenzaprine Hydrochloride Extended Release Capsules USP

#### What are cyclobenzaprine hydrochloride extended-release capsules?

Cyclobenzaprine hydrochloride extended-release capsules are prescription medicine used along with rest and physical therapy to help treat muscle spasm due to acute, painful musculoskeletal problems. Cyclobenzaprine hydrochloride extended-release capsules should only be used for up to 2 or 3 weeks. It is not known if cyclobenzaprine hydrochloride extended-release capsules are effective when used

for longer periods.

It is not known if cyclobenzaprine hydrochloride extended-release capsules are safe and effective in children.

**Do not take cyclobenzaprine hydrochloride extended-release capsules if you:**

- are allergic to cyclobenzaprine or any of the ingredients in cyclobenzaprine hydrochloride extended-release capsules. See the end of this Patient Information leaflet for a complete list of ingredients in cyclobenzaprine hydrochloride extended-release capsules.  
Talk to your healthcare provider or get medical help right away if you have symptoms of an allergic reaction such as:
  - difficulty breathing
  - hives
  - swelling of your face or tongue
  - itching
- are taking certain antidepressants, known as monoamine oxidase (MAO) inhibitors or it has been 14 days or less since you stopped taking a MAO inhibitor. Ask your healthcare provider or pharmacist for a list of these medicines if you are not sure.
- have had a recent heart attack
- have heart rhythm problems (arrhythmias)
- have heart failure
- have an overactive thyroid (hyperthyroidism)

Talk to your healthcare provider before taking this medicine if you have any of the conditions listed above.

**Before taking cyclobenzaprine hydrochloride extended-release capsules, tell your healthcare provider about all of your medical conditions, including if you:**

- have a history of eye problems including glaucoma
- have heart problems or have had a heart attack
- have liver problems
- have trouble emptying your bladder (urinary retention)
- are pregnant or plan to become pregnant. It is not known if cyclobenzaprine hydrochloride extended-release capsules will harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if cyclobenzaprine hydrochloride extended-release capsules pass into your breast milk. Talk to your healthcare provider about the best way to to feed your baby if you take cyclobenzaprine hydrochloride extended-release capsules.

**Tell your healthcare provider about all the medicines you take** including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Especially tell your healthcare provider if you take:

- a medicine to treat depression, mood, anxiety, psychotic, or thought disorders
- a pain medicine called tramadol or meperidine
- barbiturates or other medicines that depress your central nervous system (CNS depressants)
- a medicine that prevents nerve impulses (anticholinergic medicines)
- a medicine to help quit smoking called bupropion
- a blood pressure medicine called verapamil

Know the medicines you take. Keep a list of your medicines and show it to your healthcare provider or pharmacist when you get a new medicine.

**How should I take cyclobenzaprine hydrochloride extended-release capsules?**

- Take cyclobenzaprine hydrochloride extended-release capsules exactly as your healthcare provider tells you to take it.

- Your healthcare provider will tell you how much cyclobenzaprine hydrochloride extended-release capsules to take and when to take it.
- Your healthcare provider may change your cyclobenzaprine hydrochloride extended-release capsules dose if needed.
- Take cyclobenzaprine hydrochloride extended-release capsules around the same time every day.
- Swallow cyclobenzaprine hydrochloride extended-release capsules whole.
- If you have difficulty swallowing cyclobenzaprine hydrochloride extended-release capsules, tell your healthcare provider. Your healthcare provider may recommend opening the cyclobenzaprine hydrochloride extended-release capsules and mixing the contents with applesauce.
- Cyclobenzaprine hydrochloride extended-release capsules should only be taken for short periods (up to two or three weeks).
- If you take too much cyclobenzaprine hydrochloride extended-release capsules, call your healthcare provider or go to the nearest hospital emergency room right away.

### **What should I avoid while taking cyclobenzaprine hydrochloride extended-release capsules?**

You should not drink alcohol until you know how cyclobenzaprine hydrochloride extended-release capsules affect you. Taking cyclobenzaprine hydrochloride extended-release capsules with alcohol or other medicines that depress your central nervous system can slow your thinking and physical response times.

Do not drive, operate machinery, or do other dangerous activities until you know how cyclobenzaprine hydrochloride extended-release capsules affect you.

### **What are the possible side effects of cyclobenzaprine hydrochloride extended-release capsules? Cyclobenzaprine hydrochloride extended-release capsules may cause serious side effects, including:**

- **Serotonin syndrome** is a serious medical condition that may happen when cyclobenzaprine hydrochloride extended-release capsules are taken with certain other medicines. Call your healthcare provider right away or go to the nearest hospital emergency room if you have some or all of these symptoms suggestive of serotonin syndrome:
  - agitation, hallucinations, coma, or other changes in mental status
  - coordination problems or muscle twitching (overactive reflexes)
  - fast heartbeat, high or low blood pressure
  - sweating or fever
  - nausea, vomiting, or diarrhea
  - muscle stiffness or tightness
- **Cyclobenzaprine hydrochloride extended-release capsules may cause serious side effects that may lead to heart attack or stroke.** Call your healthcare provider right away or go to the nearest hospital emergency room if you have:
  - irregular or abnormal heartbeats (arrhythmias)
  - fast heartbeat (tachycardia)

### **The most common side effects of cyclobenzaprine hydrochloride extended-release capsules include:**

- dry mouth
- dizziness
- fatigue
- constipation
- nausea
- upset stomach
- drowsiness

These are not all the possible side effects of cyclobenzaprine hydrochloride extended-release capsules.

Call your healthcare provider for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**How should I store cyclobenzaprine hydrochloride extended-release capsules?**

- Store cyclobenzaprine hydrochloride extended-release capsules at room temperature between 68°F to 77°F (20°C to 25°C).
- Keep cyclobenzaprine hydrochloride extended-release capsules in a tightly closed container, and keep cyclobenzaprine hydrochloride extended-release capsules out of light.
- **Keep cyclobenzaprine hydrochloride extended-release capsules and all medicines out of the reach of children.**

**General information about the safe and effective use of cyclobenzaprine hydrochloride extended-release capsules.**

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use cyclobenzaprine hydrochloride extended-release capsules for a condition for which it was not prescribed. Do not give cyclobenzaprine hydrochloride extended-release capsules to other people, even if they have the same symptoms you have. It may harm them.

You can ask your pharmacist or healthcare provider for information about cyclobenzaprine hydrochloride extended-release capsules that is written for healthcare professionals.

**What are the ingredients in cyclobenzaprine hydrochloride extended-release capsules?**

**Active Ingredient:** cyclobenzaprine hydrochloride USP

**Inactive Ingredients:** colloidal silicon dioxide, ethyl alcohol, ethylcellulose, FDC yellow #6, gelatin, hydroxypropyl cellulose, isopropyl alcohol, potassium hydroxide, sugar spheres (which contain sucrose and corn starch) and titanium dioxide. In addition, red iron oxide is also included in the 15 mg strength. The capsule is imprinted with black ink consisting of: black iron oxide, DC Yellow #10, ethanol, FDC Blue #1, FDC Blue #2, FDC Red #40, methanol, N-butyl alcohol, propylene glycol, and shellac.

Manufactured for::

TWi Pharmaceuticals USA, Inc.

Paramus, NJ 07652

Manufactured by:

TWi Pharmaceuticals, Inc.

Taoyuan City, 32063, Taiwan

Revised: 05/2019

AMRIX is a trademark of Teva Pharmaceuticals International GmbH, or its affiliates.

---

This Patient Information has been approved by the U.S. Food and Drug Administration

**INSTRUCTIONS FOR USE**

**INSTRUCTIONS FOR USE**

**Cyclobenzaprine Hydrochloride Extended-Release Capsules**

Read this Instructions for Use before you prepare your first dose of cyclobenzaprine hydrochloride extended-release capsules mixed with applesauce using the capsule sprinkle method, each time you get a refill, and as needed. There may be new information. Ask your healthcare provider or pharmacist if you have any questions about how to mix or give a dose of cyclobenzaprine hydrochloride extended-release capsules using the capsule sprinkle method.

**Important Information:**



- **Do not chew cyclobenzaprine hydrochloride extended-release capsules or the granules that are in the capsules.**
- The capsule sprinkle method for mixing the contents of cyclobenzaprine hydrochloride extended-release capsules with applesauce may be used for adults who cannot swallow capsules. **Do not use any other food in the place of applesauce.**

### **Preparing a dose of cyclobenzaprine hydrochloride extended-release capsules using the capsule sprinkle method.**

Before you prepare a dose of cyclobenzaprine hydrochloride extended-release capsules mixed with applesauce using the capsule sprinkle method, gather the following supplies:

- paper towels
- tablespoon
- applesauce
- cup of water

**Step 1:** Choose a clean, flat work surface. Place a clean paper towel on the work surface. Then place the other supplies on the paper towel.

**Step 2:** Wash and dry your hands well.

**Step 3:** Check the dose that was prescribed by your healthcare provider. Take out the number of cyclobenzaprine hydrochloride extended-release capsules needed to prepare your dose. Place them on the paper towel.

**Step 4:** Place enough applesauce to fill your tablespoon. Set the tablespoon down on the paper towel.



**Step 5:** Hold the cyclobenzaprine hydrochloride extended-release capsule in an upright position (vertical) directly over the tablespoon. Hold each end of the cyclobenzaprine hydrochloride extended-release capsule between your thumbs and index (pointer) fingers.



**Step 6:** Carefully twist both ends of the cyclobenzaprine hydrochloride extended-release capsule in opposite directions to open it. Be careful not to spill the capsule contents.



**Step 7:** Sprinkle the contents of the cyclobenzaprine hydrochloride extended-release capsule onto the applesauce.

- Check the capsule shells to make sure they are empty.
- Throw away the empty capsule shells.



**If the total prescribed dose is more than 1 capsule, repeat Steps 5 through 7 for each capsule. Do not add more applesauce. Then follow the rest of the steps below.**

**Step 8:** Pick up the tablespoon and swallow the cyclobenzaprine hydrochloride extended-release capsule contents and applesauce mixture right away. **Do not chew the cyclobenzaprine hydrochloride extended-release capsule contents and applesauce mixture.**



**Step 9:** Rinse your mouth with a sip of water and swallow to make sure that all of the cyclobenzaprine hydrochloride extended-release granules have been swallowed.



**Step 10:** Throw away any unused cyclobenzaprine hydrochloride extended-release capsule content and applesauce mixture. Do not keep any cyclobenzaprine hydrochloride extended-release capsule content and applesauce mixture for future use.

### **How should I store cyclobenzaprine hydrochloride extended-release capsules?**

- Store cyclobenzaprine hydrochloride extended-release capsules at room temperature between 68°F

to 77°F (20°C to 25°C).

**Keep cyclobenzaprine hydrochloride extended-release capsules and all medicines out of the reach of children.**

This Instructions for Use has been approved by the U.S. Food and Drug Administration.

Issued: 05/2019

**Packaged and Distributed by** *Quality Care Products, LLC* **1-800-337-8603**



Holland, OH 43528

**CYCLOBENZAPRINE HCl 15 MG**  
EXTENDED-RELEASE CAPSULES, USP

Warning: Keep out of children's reach.  
Store at 68 to 77 degrees F.  
Consult with a physician. See manufacturer's insert.



GTIN: 00355700857608

NDC: 55700-0857-60

Serial: **#60 CAPSULES**

Lot: Rx only

EXP: //


Each extended-release capsule contains Cyclobenzaprine hydrochloride USP, 15 mg

Mfr for: TWI Pharmaceuticals USA, Inc., Paramus, NJ 07652

Mfr by: TWI Pharmaceuticals, Inc., Taoyuan City, 32063, Taiwan

SOURCE CODE: 24979003504


**Packaged and Distributed by** *Quality Care Products, LLC* **1-800-337-8603**



Holland, OH 43528

**CYCLOBENZAPRINE HCl 30 MG**  
EXTENDED-RELEASE CAPSULES, USP

Warning: Keep out of children's reach.  
Store at 68 to 77 degrees F.  
Consult with a physician. See manufacturer's insert.



GTIN: 00355700858605

NDC: 55700-0858-60

Serial: **#60 CAPSULES**

Lot: Rx only

EXP: //

Each extended-release capsule contains Cyclobenzaprine hydrochloride USP, 30 mg

Mfr for: TWI Pharmaceuticals USA, Inc., Paramus, NJ 07652

Mfr by: TWI Pharmaceuticals, Inc., Taoyuan City, 32063, Taiwan

SOURCE CODE: 24979003504

## CYCLOBENZAPRINE HYDROCHLORIDE

cyclobenzaprine hydrochloride capsule, extended release

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:55700-857(NDC:24979-035)
Route of Administration	ORAL		

Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
CYCLOBENZAPRINE HYDRO CHLORIDE (UNII: 0VE05JYS2P) (CYCLOBENZAPRINE - UNII:69O5WQQ5TI)		CYCLOBENZAPRINE HYDROCHLORIDE	15 mg	
Inactive Ingredients				
Ingredient Name			Strength	
FERROSO FERRIC O XIDE (UNII: XM0M87F357)				
SILICON DIO XIDE (UNII: ETJ7Z6XBU4)				
ALCOHOL (UNII: 3K9958V90M)				
ETHYLCELLULOSE (10 MPA.S) (UNII: 3DYK7UYZ62)				
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)				
GELATIN (UNII: 2G86QN327L)				
HYDROXYPROPYL CELLULOSE (1200000 WAMW) (UNII: U3JF91U133)				
ISOPROPYL ALCOHOL (UNII: ND2M416302)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
POTASSIUM HYDRO XIDE (UNII: WZH3C48M4T)				
SHELLAC (UNII: 46N107B71O)				
SUCROSE (UNII: C151H8M554)				
TITANIUM DIO XIDE (UNII: 15FIX9V2JP)				
FERRIC O XIDE RED (UNII: 1K09F3G675)				
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)				
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)				
FD&C RED NO. 40 (UNII: WZB9127XOA)				
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)				
BUTYL ALCOHOL (UNII: 8PJ61P6TS3)				
METHYL ALCOHOL (UNII: Y4S76JWI15)				
STARCH, CORN (UNII: O8232NY3SJ)				
Product Characteristics				
Color	ORANGE (Dark Orange Opaque Cap; Red Opaque Body)	Score	no score	
Shape	capsule	Size	14mm	
Flavor		Imprint Code	T035	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55700-857-60	60 in 1 BOTTLE; Type 0: Not a Combination Product	04/17/2020	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA		ANDA091281	04/17/2020	

CYCLOBENZAPRINE HYDROCHLORIDE

cyclobenzaprine hydrochloride capsule, extended release

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:55700-858(NDC:24979-036)
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
CYCLOBENZAPRINE HYDROCHLORIDE (UNII: 0VE05JYS2P) (CYCLOBENZAPRINE - UNII:69O5WQQ5TI)	CYCLOBENZAPRINE HYDROCHLORIDE	30 mg

Inactive Ingredients	
Ingredient Name	Strength
FERROSO FERRIC OXIDE (UNII: XM0M87F357)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
ALCOHOL (UNII: 3K9958V90M)	
ETHYLCELLULOSE (10 MPAS) (UNII: 3DYK7UYZ62)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN (UNII: 2G86QN327L)	
HYDROXYPROPYL CELLULOSE (1200000 WAMW) (UNII: U3JF91U133)	
ISOPROPYL ALCOHOL (UNII: ND2M416302)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
SHELLAC (UNII: 46N107B71O)	
SUCROSE (UNII: C151H8M554)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
BUTYL ALCOHOL (UNII: 8PJ61P6TS3)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
METHYL ALCOHOL (UNII: Y4S76JWI15)	
STARCH, CORN (UNII: O8232NY3SJ)	

Product Characteristics			
Color	orange (Orange Opaque Cap; White Opaque Body)	Score	no score
Shape	capsule	Size	16mm
Flavor		Imprint Code	T036
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55700-858-60	60 in 1 BOTTLE; Type 0: Not a Combination Product	04/17/2020	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091281	04/17/2020	

**Labeler** - Quality Care Products, LLC (831276758)

## Establishment

Name	Address	ID/FEI	Business Operations
Quality Care Products, LLC		831276758	relabel(55700-857, 55700-858)

Revised: 4/2020

Quality Care Products, LLC